



STATE OF MARYLAND

DHMH

Maryland Department of Health and Mental Hygiene

201 W. Preston Street • Baltimore, Maryland 21201

Martin O'Malley, Governor – Anthony G. Brown, Lt. Governor – Joshua M. Sharfstein, M.D., Secretary

November 25, 2014

The Honorable Martin O'Malley
Governor
State of Maryland
Annapolis, MD 21401-1991

The Honorable Thomas V. Mike Miller, Jr.
President of the Senate
H-107 State House
Annapolis, MD 21401-1991

The Honorable Michael E. Busch
Speaker of the House
H-101 State House
Annapolis, MD 21401-1991

Re: Health-General Article, §13-1004(d), FY 2013 Biennial Tobacco Study
Cigarette Restitution Fund – Tobacco Use Prevention and Cessation Program

Dear Governor O'Malley, President Miller, and Speaker Busch:


Pursuant to Health-General Article, §13-1004(d), Annotated Code of Maryland, the Department of Health and Mental Hygiene (the Department) is directed to produce a biennial legislative report on the results of the Biennial Tobacco Study.

The enclosed legislative report summarizes trends related to tobacco use behaviors among underage middle and high school youth and adults ages 18 and older since 2000. Included findings are derived from the results of the biennial Youth Tobacco Survey (2000-2012) and the corresponding expanded Youth Tobacco and Risk Behavior Survey (2013). Findings on adult behaviors are derived from the results of the Behavioral Risk Factor Surveillance System (2000-2012). Data are presented for Maryland as a whole, as well as individually for each county and Baltimore City. Significant findings document continued reductions in tobacco use behaviors since program inception in Fiscal Year 2001.

This report was due September 1, 2013, and the Department apologizes for the lateness of this submission. The Department experienced a delay in its procurement process, which caused the data collection for the Maryland Youth Tobacco and Risk Behavior Survey (MYTRBS) to be delayed as well. The MYTRBS data serves as the basis for this report.

The Department appreciates your commitment to the progress we are making in reducing tobacco use in Maryland. If you have questions about this report, please contact Ms. Allison Taylor, Director of Governmental Affairs at (410) 767-6481.

Sincerely,

A handwritten signature in black ink, appearing to read "Josh M. Sharfstein". The signature is fluid and cursive, with the first name "Josh" being more prominent.

Joshua M. Sharfstein, M.D.
Secretary

Enclosure

cc: Allison W. Taylor, Director, Office of Governmental Affairs
Laura Herrera, Deputy Secretary for Public Health Services
Michelle Spencer, Director, Prevention and Health Promotion Administration
Sarah Albert, MSAR #2866

Monitoring Changing Tobacco Use Behaviors Underage Maryland Youth: 2000–2013 Adults Ages 18 and Older: 2000–2012

A Report to the Maryland Governor and General Assembly

Maryland Department of Health and Mental Hygiene

Cigarette Restitution Fund's Tobacco Use Prevention and Cessation Program

Fiscal Year 2013



Martin O'Malley

Governor
State of Maryland

Anthony G. Brown

Lieutenant Governor
State of Maryland

Joshua M. Sharfstein, MD

Secretary
Department of Health and Mental Hygiene

Statutory Authority and Requirements

Maryland's Health-General Article, Title 13, Subtitle 10, requires the Maryland Department of Health and Mental Hygiene (DHMH) to conduct a biennial tobacco study and report specific findings to the Maryland Governor and the General Assembly. The appendices to this Part 1 of the Report provide the detailed data for indicators DHMH is required to report in its biennial tobacco study for underage youth.

This space was left blank intentionally.

Sources

Unless otherwise explicitly stated, all data come from the Maryland Youth Tobacco Surveys (2000, 2002, 2006, 2008, 2010) and the Maryland Youth Tobacco and Risk Behavior Survey (2013).

Suggested Citation

Monitoring Changing Tobacco Use Behaviors: A Report to the Maryland Governor and the General Assembly, Fiscal Year 2013. (2014, August). Baltimore, MD: Maryland Department of Health and Mental Hygiene.

This space was left blank intentionally.

Table of Contents

PART 1—UNDERAGE YOUTH TOBACCO USE BEHAVIORS

	<u>Pages</u>
Key Findings	1–2
Introduction	3–4
Underage Use of Tobacco Products	5–10
Cigarette Smoking by Grade in School.....	6
Initiation of Tobacco Products in the Past 12 Months	7
Concurrent Use of More Than One Tobacco Product	7
Current Use of Flavored Tobacco Products	8
Underage Access to Tobacco Products.....	9
Underage Smoking and Use of Alcohol and Drugs	9
Underage Youth Attitudes and Beliefs	11–12
School-Based Tobacco Use Prevention Curricula	11
Underage Youth Exposure to Secondhand Smoke	13–14

PART 2—ADULT TOBACCO USE BEHAVIORS

Key Findings	17–18
Introduction	19–21
Adult Tobacco Use and Health.....	22–26
Tobacco-Related Cancers and Disease	23
Cost of Treating Tobacco-Related Cancers and Disease.....	25
Preventable Hospitalizations—Resulting from Tobacco-Related Cancers and Disease.....	26
Adult Current Tobacco Use	27–30
Adults Quitting Tobacco Use.....	31–34
Quitting Smoking.....	31
Adult Initiation of Tobacco Use.....	35–37
Adult Exposure to Secondhand Smoke	38

APPENDICES

Appendix A—Underage Youth Prevalence, Initiation, and Cessation	39–90
Appendix B—Adult Prevalence, Initiation, and Cessation	91–142
Appendix C—Where to Find More Information	143

This space was left blank intentionally.

PART 1—UNDERAGE YOUTH TOBACCO USE BEHAVIORS

This space was left blank intentionally.

KEY FINDINGS—UNDERAGE YOUTH

In 2013

- Cigarette smoking by Maryland high school youth younger than 18 years old has decreased substantially since 2000.
- Eleven percent (11.0%) of youth reported smoking a cigarette during the past 30 days—a 52.2% decrease from the 2000 rate of 23.0%.
- In 2013, cigar smoking by underage high school youth (11.5%) became just as popular as cigarette smoking (11.0%). Among underage tobacco users, 47.9% used more than one type of tobacco product.
- The use of smokeless tobacco by underage high school youth increased by 43.8% between 2010 and 2013.
- Underage youth who use cigars are also likely to smoke cigarettes (38.0% of underage youth tobacco users reported smoking both cigars and cigarettes).
- Seventy-one percent (71.4%) of underage youth who smoke cigars reported using flavored cigars in 2013. Fifty-seven percent (57.1%) of underage youth who use smokeless tobacco also use flavored smokeless tobacco products.
- Asking for photo identification reduces cigarette sales to underage youth. When tobacco retailers request to see photo identification when youth try to buy cigarettes, 32.7% of underage youth succeed in buying cigarettes—as compared with a 78.7% success rate when photo identification is not requested (Federal law requires retailers to ask for photo identification).
- Cigarette smoking by underage youth is highly correlated with underage use of alcohol, marijuana, abuse of prescription drugs, and use of other illegal drugs. The younger a person starts smoking, the more likely he or she is to engage in the risk behaviors listed below.
- Forty percent (40.0%) of underage high school smokers believe smoking helps them to look cool or fit in, an increase of 50.0% since 2000.

Table 1.
Alcohol, Marijuana, and Other Drug Use
Maryland Underage High School Youth, 2013

Smoking Status	Currently Drink Alcohol	Currently Use Marijuana	Currently Abuse Prescription Drugs	Ever Use Other Illegal Drugs
Nonsmokers	23.7%	12.6%	4.2%	8.9%
Smokers	79.4%	67.0%	37.9%	51.1%
Increased Likelihood of Smokers Engaging in Behavior*	3x	5x	9x	6x

* The increased likelihood represents the relative risk of a smoker engaging in the behavior versus that of a nonsmoker.

KEY FINDINGS—UNDERAGE YOUTH

- Twenty percent (20.6%) of underage high school nonsmokers believe smoking helps youth to look cool and fit in, an increase of 63.5% since 2000. Most of the increase occurred between 2010 and 2013.
- In households with a resident adult and underage children present, voluntary bans on smoking have increased 29.5% from 48.5% in 2000 to 62.8% in 2013.

Moving Forward

Flavored Tobacco Products and Cigar Package Size

- In 2012, increases in Maryland's excise tax on the cheap (nonpremium) cigars preferred by youth were seen as a means to deter use of these popular products.
- However, possibly as a result of Maryland's excise tax on those cheap cigars being increased to more than twice that of the excise tax on smokeless tobacco products in 2012, smokeless tobacco use by youth is now increasing.
- Unlike cigarettes which cannot be flavored and must be sold in packs of at least 20 cigarettes, both cigars and smokeless tobacco are sold in fruit and candy flavors and have no minimum package size requirements.
- Care should be taken to avoid unintentionally favoring one tobacco product over another when adopting tobacco control policies. Youth have proven ready to move to the products most accessible and least expensive.

Introduction

In Fiscal Year (FY) 2001, using nontax proceeds from participation in the national Master Settlement Agreement with the tobacco industry, the Maryland Department of Health and Mental Hygiene (DHMH) implemented a tobacco control program modeled on the best practice recommendations of the Centers for Disease Control and Prevention (CDC). In addition, by 2008 Maryland increased its excise tax on cigarettes to \$2.00 per pack and implemented the Clean Indoor Air Act, a comprehensive statutory smoking ban that applies to all worksites, restaurants, and bars.

DHMH is mandated by Health—General Article Title 13, Subtitle 10 to monitor changing tobacco use behaviors of youth and adults, and to report its findings biennially to the Maryland Governor and the General Assembly. This document fulfills that requirement with Part 1 focusing on the tobacco use behaviors of underage youth, while Part 2 focuses on the tobacco use behaviors of adults.

About the Data

The vast majority of the data presented in this report are from the Maryland Youth Tobacco and Risk Behavior Survey (MYTRBS), a biennial survey that combines the CDC's Youth Risk Behavior Survey (YRBS) and the Maryland Youth Tobacco Survey (MYTS), and provides both state-level and county-level data. The first MYTRBS was scheduled for the fall of 2012 (and in the fall of every year thereafter), although delayed, the inaugural administration of this survey was conducted in the spring of 2013. MYTRBS results are treated by the CDC as an official YRBS for the State of Maryland, each county, and Baltimore City.

The MYTRBS and the MYTS use identical survey methodologies and produce comparable data. However, due to some small differences in analysis, the results may not be identical. Caution should be exercised when comparing 2010 MYTS estimates to the 2013 MYTRBS estimates.

The YRBS was previously administered in Maryland by MSDE biennially (2005–2011) and provided state-level data only. The MYTS was previously administered in Maryland by DHMH biennially (2000–2010) and provided both state-level and county-level data. The new combined MYTRBS provides both state-level and county-level data biennially.

This report focuses on tobacco use behaviors previously monitored through the MYTS and fulfills the statutory data reporting requirement. Data on the broader range of risk behaviors addressed by the MYTRBS as an official CDC YRBS will be included in a separate report to be issued by the DHMH Center for Chronic Disease Prevention and Control.

Data on underage youth tobacco use behaviors is collected because early initiation of tobacco use, coupled with the addictive qualities of the nicotine found in all tobacco products, often leads to several decades or lifelong use of such products. Tobacco use significantly increases a smoker's risk for a long list of cancers, cardiovascular conditions, and respiratory diseases later in adult life. At the same time, the use of

INTRODUCTION TO YOUTH TOBACCO USE BEHAVIORS

tobacco products can have an immediate and detrimental impact on the health and well-being of underage youth.

All references to “youth” in this report refer only to youth less than 18 years old, unless otherwise stated. Most of the data presented in the charts in the main body of this report apply to public high school youth only, unless otherwise stated. Detailed data on middle school tobacco use behaviors can be found in Appendix A—Underage Youth.

Important Terms Defined

Statistically significant change—refers to observed differences between two estimates that are not likely the result of chance. A conservative approach to determining statistically significant change is used in this report—the confidence intervals around the estimates cannot overlap to be statistically significant.

Confidence levels and intervals—refers to estimates made using a 95% confidence level, that is, if the survey had been repeated 100 times, 95 of the 100 estimates would fall within the stated confidence interval. The stated confidence interval is an amount above and below the estimate, with the estimate at the midpoint of the interval. For example, the 2012 estimate for current cigarette smoking is 16.2% with a confidence interval of 15.0% to 17.4%. This means that we are 95% confident that the estimated smoking rate for adults in Maryland would have fallen between 15.0% and 17.4% in 95 out of 100 times if the survey had been repeated. The 16.2% estimate is the midpoint of the confidence interval.


UNDERAGE USE OF TOBACCO PRODUCTS

With respect to youth, the CDC defines “current use” as any use of a tobacco product within 30 days before survey administration (even a single puff of a cigarette). The percentage of public high school youth who currently use one or more types of tobacco product is significantly higher than the percentage of middle school youth (17.7% and 6.7%, respectively). The greater popularity of tobacco among high school students as compared with middle school students has remained consistent since 2000, when data collection began. Between 2000 and 2013, there has, however, been a significant decrease in current tobacco use among high school youth (from 29.4% to 17.7%).

Cigars have now surpassed cigarettes as the most popular tobacco product among public high school youth **who used any tobacco product during the past 30 days**. Cigars are followed by cigarettes and smokeless tobacco products. In 2013, cigars were used by 11.5% of Maryland public high school youth. Cigarettes and smokeless tobacco products were used by 11.0% and 6.9% of these youth, respectively. Decreases in cigarette and cigar smoking between 2010 and 2013 are statistically significant. In contrast, use of smokeless tobacco by underage youth saw a statistically significant increase between 2010 and 2013 (see Figure 1).

Reported cigar smoking increased significantly in 2008 after years of steady decline. In the absence of data that pinpoint the reasons, possible explanations for the increase include: (1) the significant increase in the state excise tax on cigarettes in FY 2009—perhaps prompting youth to switch to cheaper little cigar products that can be sold individually because there is no minimum package size requirement for cigars as there is for cigarettes; (2) implementation of the U.S. Food and Drug Administration (FDA) ban on flavored cigarettes but not flavored cigars; and (3) the clarification of the survey question in 2008 as to which products are considered cigars—perhaps resulting in more youth who answer “yes” when asked if they smoked cigars. The subsequent decrease in cigar smoking since 2008 is statistically significant. Reasons for this decrease are most likely: (1) implementation of Maryland’s educational campaign (“The Cigar Trap”) and (2) the FY 2012 increase in the excise tax on the nonpremium cigars favored by youth. A recent statistically significant increase in the use of smokeless tobacco products may be the result of higher excise taxes on cigarettes and cheap cigars.

Figure 1.
Current Use of Cigarettes, Cigars, and Smokeless Tobacco Products (%)
Underage Maryland Public High School Youth, 2000–2013




Sources: Maryland Youth Tobacco Surveys, 2000–2010, Maryland Youth Tobacco and Risk Behavior Survey, 2013.

UNDERAGE USE OF TOBACCO PRODUCTS

Cigarette Smoking by Grade in School

Cigarette smoking increases by school grade, a pattern seen in every survey since the first MYTS administered in the fall of 2000. The percentage of youth in any single grade who currently smoke cigarettes is significantly less in 2013 than in 2000. Further, between 2010 and 2013, there were significant decreases in cigarette use among 9th-, 10th-, 11th-, and 12th-graders (see Figure 2).


Figure 2.
Current Cigarette Smoking by Grade (%)
Underage Maryland Public Middle and High School Youth, 2000–2013



Sources: Maryland Youth Tobacco Surveys, 2000–2010, Maryland Youth Tobacco and Risk Behavior Survey, 2013.

Figure 3 presents the percentage of underage high school youth who were currently using any type of tobacco product in spring of 2013. It is evident that there is considerable variation between jurisdictions (ranging from 10.2% to 33.1%), and that the prevalence of “any use of tobacco” in many jurisdictions is considerably higher than the statewide figure of 15.7%.

Figure 3.
Current Use of Any Type of Tobacco Product (%)
Underage Maryland Public High School Youth, 2013




Source: Maryland Youth Tobacco and Risk Behavior Survey, 2013.

Initiation of Tobacco Products in the Past 12 Months

Among all underage middle and high school youth, 12.3% reported initiating the use of any kind of tobacco product in the 12 months preceding the 2013 survey, compared with 18.1% of underage youth who reported doing so in 2010. (Note: In 2013, the questionnaire items that assess initiation of tobacco use were modified. Comparisons with previous survey years should be interpreted with caution.) Figure 4 presents the proportion of youth within a given grade who reported first starting tobacco use in the preceding year. There are significant increases in the initiation of tobacco use between each grade, except for grades 11 and 12. An estimated 36,294 underage high school youth reported first using tobacco products during the preceding 12 months, as did 10,721 middle school youth. This averages to 129 new underage tobacco users daily in Maryland.

Figure 4.
Initiation of Use of Tobacco Products in the Past Year by Grade (%)
Underage Maryland Public High School Youth, Spring 2013 Survey




Source: Maryland Youth Tobacco and Risk Behavior Survey, 2013.

Concurrent Use of More Than One Tobacco Product

In 2013, 17.7% of underage high school youth used some form of tobacco product in the preceding 30 days. As shown in Figure 5, of these tobacco users, 47.9% used more than one tobacco product in the preceding 30 days, compared with 42.6% in 2010 (this change is statistically significant). While cigars were the most prevalent tobacco product used among underage youth overall (11.5%), among underage youth who used multiple tobacco products during the past 30 days, cigarettes were the most popular tobacco product, with 68.4% smoking cigarettes (followed by cigars at 62.4% and smokeless tobacco at 33.5%). Of those using multiple tobacco products, most (38.0%) used both cigarettes and cigars. The overall proportion of multiple product users increased between 2010 and 2013.

UNDERAGE USE OF TOBACCO PRODUCTS

Figure 5.
Single Tobacco Product vs. Multiple Tobacco Product Use by Current Tobacco Users (100%)
Underage Maryland Public High School Current Tobacco Users, 2010 and 2013




Source: Maryland Youth Tobacco Survey, 2010.

Source: Maryland Youth Tobacco and Risk Behavior Survey, 2013.

Current Use of Flavored Tobacco Products

Between 2010 and 2013, there has been a significant increase in the use of flavored tobacco products among underage public high school youth who use cigars or smokeless tobacco (72.1% and 77.1%, respectively). Figure 6 presents (1) the percentage of underage middle and high school youth who smoked cigars or used smokeless tobacco and who used a flavored tobacco product (either flavored cigars or flavored smokeless tobacco, (2) the percentage who currently smoke cigars and used only flavored cigars, and (3) the percentage who currently use smokeless tobacco and used only flavored smokeless tobacco. Although there has been a significant decrease in the use of flavored cigars and smokeless tobacco among middle and high school cigar and smokeless tobacco users, nearly three-quarters of high school cigar users reported flavored cigar use in 2013.

Figure 6.
Use of Flavored Tobacco Products Among Those Who Currently Use Cigars or Smokeless Tobacco
Maryland Underage Public Middle and High School Cigar or Smokeless Tobacco Users, 2010 and 2013




Sources: Maryland Youth Tobacco Survey, 2010, the Maryland Youth Tobacco and Risk Behavior Survey, 2013.

Underage Access to Tobacco Products

In the United States, it is illegal to sell cigarettes or other tobacco products to youth under 18 years old. The FDA's regulations restricting the sale and distribution of cigarettes and smokeless tobacco require retailers to check photo identification (ID) of anyone purchasing cigarettes or smokeless tobacco who appears to be less than 27 years old.

Among underage public high school smokers who reported attempting to purchase cigarettes from a retail outlet during the preceding 30 days, 37.5% reported being asked for photo ID. Of the underage high school smokers who were not asked for photo ID at a retail outlet, 78.7% were able to purchase cigarettes despite being underage. In contrast, of the youth who were asked for photo ID, significantly fewer (32.7%) were sold cigarettes (see Figure 7). Between 2010 and 2013, there was no significant change in the percentage of underage high school youth who were asked for proof of age when attempting to purchase cigarettes.

Figure 7.
Refusal of Sales—With and Without Request for Proof of Age
Maryland Underage Public High School Cigarette Smokers, 2013



Source: Maryland Youth Tobacco and Risk Behavior Survey, 2013.

Underage Smoking and Use of Alcohol and Drugs

Underage high school youth who report smoking cigarettes also are more likely to report drinking alcohol, using marijuana, abusing prescription drugs, or using other illegal drugs compared with nonsmoking youth. Approximately 79.4% of underage high school cigarette smokers report they currently use alcohol (i.e., have had one or more drinks of alcohol in the past 30 days), compared with 23.7% of nonsmokers (see Figure 8). Similarly, 67.0% of underage high school cigarette smokers reported they currently use marijuana (i.e., have smoked marijuana during the past 30 days), compared with 12.6% of nonsmokers. Despite decreased smoking prevalence, the proportion of cigarette smokers who use alcohol has not changed.

UNDERAGE USE OF TOBACCO PRODUCTS

Figure 8.
Alcohol, Marijuana, and Other Drug Use
Maryland Underage Public High School Youth, 2013


Smoking Status	Currently Drink Alcohol	Currently Use Marijuana	Currently Abuse Prescription Drugs	Ever Use Other Illegal Drugs
Nonsmokers	23.7%	12.6%	4.2%	8.9%
Smokers	79.4%	67.0%	37.9%	51.1%
Increased Likelihood of Smokers Engaging in Behavior*	3x	5x	9x	6x

* The increased likelihood represents the relative risk of a smoker engaging in the behavior vs. that of a nonsmoker. Note that the data above represent observed correlations in risk behaviors and are not presented as evidence of causality.

Source: Maryland Youth Tobacco and Risk Behavior Survey, 2013.

The age at which youth initiate cigarette smoking also is correlated with their likelihood to engage in other risk behaviors. Figure 9 presents the percentage of underage high school cigarette smokers who currently use alcohol, use marijuana, abuse prescription drugs, or used another illegal drug by the age at which they smoked their first cigarette. The younger a youth begins smoking, the more likely he or she is to engage in one of these risk behaviors.

Figure 9.
Use of Alcohol, Marijuana, and Other Drugs by Age When Smoked First Whole Cigarette
Underage Maryland Public High School Cigarette Smokers, 2013




Source: Maryland Youth Tobacco and Risk Behavior Survey, 2013.

School-Based Tobacco Use Prevention Curricula

In 2013, 81.6% of Maryland public middle school youth and 70.4% of public high school youth reported that they had been taught about the dangers of tobacco use during the preceding school year (2011–2012). Figure 10 documents trends in exposure to tobacco use prevention curricula in Maryland public schools from the 1999–2000 school year through the 2011–2012 school year. There has been a statistically significant increase in middle and high school youth exposure to such curricula since the Cigarette Restitution Fund (CRF) Tobacco Use Prevention and Control Program was first implemented in 2001 and a significant increase between calendar years 2010 and 2013.

Figure 10.
Taught About Dangers of Tobacco Use, by School Year
Underage Maryland Public Middle and High School Youth, 1999–2012 School Years




Sources: Maryland Youth Tobacco Surveys, 2000–2010, Maryland Youth Tobacco and Risk Behavior Survey, 2013. No surveys were conducted during the 2003–2004 school year.

School-based health education, community-based prevention activities, restrictions on youth access to tobacco, statewide prevention campaigns, mass media prevention messages, tobacco industry marketing, promotion, and advertising, and perceived social norms all influence the attitudes of underage youth toward smoking and tobacco products. As presented in Figures 11 and 12, between 2010 and 2013, there were significant increases in the percentage of underage high school youth (smoking and nonsmoking) who believe that smoking helps youth to “fit in or look cool” and believe that smokers have more friends.


UNDERAGE YOUTH ATTITUDES AND BELIEFS

Figure 11.
Believe Smoking Helps Youth to “Fit In” or “Look Cool,” by Smoking Status
Underage Maryland Public High School Youth, 2000–2013



Sources: Maryland Youth Tobacco Surveys 2000–2010, Maryland Youth Tobacco and Risk Behavior Survey, 2013.

Figure 12.
Believe Smokers Have More Friends, by Smoking Status
Underage Maryland Public High School Youth, 2000–2013




Sources: Maryland Youth Tobacco Surveys, 2000–2010, Maryland Youth Tobacco and Risk Behavior Survey, 2013.

UNDERAGE YOUTH EXPOSURE TO SECONDHAND SMOKE


Underage youth exposure to secondhand smoke can occur at home, in the car, or in other indoor locations. In addition to the Maryland Clean Indoor Air Act's mandatory smoking ban in almost all indoor public areas, local and state health department tobacco control programs have encouraged voluntary in-home and in-car prohibitions on smoking to reduce this exposure. Since 2000, there has been a significant decrease in the percentage of underage youth who report being exposed to secondhand smoke (see Figure 13). In households without a resident smoker, there has also been an increase in youth reporting that they have not been exposed to secondhand smoke. The 2013 data, however, suggest that this progress may be eroding—the steady upward trend (2000–2008) having slowed (2008–2010) and then decreased in 2013.

Figure 13.
Underage Youth NOT Exposed to Secondhand Smoke Indoors
Maryland Public Middle and High School Youth, 2000–2013



Sources: Maryland Youth Tobacco Surveys, 2000–2010, Maryland Youth Tobacco and Risk Behavior Survey, 2013.

Figure 14.
Voluntary Smoking Prohibition Inside the Home
Maryland Public Middle and High School Youth (All Ages), 2006–2013



Sources: Maryland Youth Tobacco Surveys, 2000–2010, Maryland Youth Tobacco and Risk Behavior Survey, 2013.

UNDERAGE YOUTH EXPOSURE TO SECONDHAND SMOKE

The critical program elements identified in the CDC Best Practice Recommendations (2014) as essential to changing the behaviors of adults about exposure to secondhand smoke include: dissemination of information about the dangers smoke from tobacco products poses to those who inhale it secondhand, creation and enforcement of smoke-free policies, and changes in attitudes and beliefs about secondhand smoke. Together, these elements can create a social norm in which involuntary exposure to secondhand smoke is not acceptable. One measure of the success of such initiatives is the extent to which underage youth are routinely exposed to secondhand smoke.


PART 2—ADULT TOBACCO USE BEHAVIORS

This space was left blank intentionally.

In 2012

- By every measure, cigarette smoking by Maryland adults has decreased substantially since 2000.
- Sixteen percent (16.2%) of Maryland adults reported that they currently smoked cigarettes, with 74.0% of these smokers wanting to quit smoking.
- Four percent (4.2%) of Maryland adults were committed cigarette smokers (currently smoke and do not want to quit smoking now or in the future).
- Smokers' desire to quit smoking was universal—no statistically significant difference was found between racial/ethnic groups, gender, income, or highest educational attainment.
- Smokers who want to quit will reach out for assistance in quitting when made aware of available support services such as Maryland's telephone Tobacco Quitline 1-800-QUIT-NOW (see Figure 15).

Figure 15.
Monthly Registered Callers/Web Users of Maryland Quitline, 2011–2013



Source: Maryland Tobacco Quitline, 1-800-QUIT-NOW

- Never starting to smoke is increasingly common. In 2012 almost 61.0% of Maryland adults report that they have never smoked cigarettes.
- Decreased smoking is improving health and reducing health care costs. Hospital admissions to treat tobacco-related cancers decreased 11.0% between 2000 and 2011, saving more than \$102 million in hospital charges in 2011.

Moving Forward

Tobacco Use Cessation

- DHMH will continue to support and operate the Maryland Tobacco Quitline, 1-800-QUIT-NOW, and ensure universal accessibility to smokers who want to quit.
- DHMH and local health departments will continue to support and implement the Pregnancy and Tobacco Cessation Help (PATCH) initiative in local communities, aiming to reduce maternal and paternal smoking during pregnancy, as well as smoking among women of childbearing age, and improve health outcomes for newborns.
- DHMH and its partners will continue to implement and expand collaborative efforts to institutionalize smoking cessation services for the behavioral health – mental health and substance abuse – populations, both of which have extremely high rates of smoking.
- DHMH will monitor e-cigarette use to determine whether these products are being used (1) to successfully quit all nicotine products; (2) instead of tobacco products, primarily where smoking is prohibited; or (3) to start use of these devices and never start smoking cigarettes.

Preventing the Initiation of Tobacco Use

- DHMH, local health departments, and their partners will continue to support efforts at the state and community level to discourage underage youth and young adults from initiating tobacco use.
- Flavored tobacco, particularly candy- and fruit-flavored cigars and smokeless tobacco, continue to provide an easy path to nicotine addiction. DHMH will monitor use of these products, as well as public health proposals to restrict their availability.
- DHMH will monitor e-cigarette use to determine if candy and fruit flavors play a role in their popularity and how they are used—and if they are an entry point for nicotine addiction among young adults who had not previously used tobacco regularly.

Clean Indoor Air

- DHMH, local health departments, and their partners will continue to support community efforts promoting the adoption of Clean Indoor Air rules on school properties at all times, in multi-unit housing, public parks, and beaches—and encourage voluntary adoption of such rules in private homes.
- DHMH and local health departments will monitor the impact of e-cigarette use on the ability to easily enforce existing Clean Indoor Air regulations.

INTRODUCTION—ADULT TOBACCO USE BEHAVIORS

Introduction

The focus of this part of the report is on adult tobacco use behaviors at the state-level, while county-level data can be found in Appendix B—Adults at the end of the report. In FY 2001, using non-tax proceeds from participation in the national Master Settlement Agreement with the tobacco industry, Maryland implemented a comprehensive tobacco control program modeled on the best practice recommendations of the CDC. Today, Maryland has a comprehensive statutory smoking ban (applying to all worksites, restaurants, and bars); the Maryland Tobacco Quitline (1-800-QUIT-NOW); assistance for those who use tobacco to succeed in quitting; and a robust tobacco program to encourage underage youth and young adults not to start using tobacco.

About the Data

Most of the data presented in Part 2 of the report are from the CDC's Behavioral Risk Factor Surveillance System (BRFSS). The BRFSS is a random-digit-dial telephone survey sponsored jointly by the CDC and DHMH, and is conducted throughout each calendar year. The objective of the BRFSS is not specific to tobacco-behaviors, but rather to collect uniform, state-specific data on preventive health practices and risk behaviors that are linked to chronic diseases, injuries, and preventable infectious diseases that affect the adult population in Maryland and across the nation. Risk factors monitored by the BRFSS include tobacco use, health care coverage, HIV/AIDS knowledge and prevention, physical activity, and fruit and vegetable consumption.

Data are collected from a random sample of adults age 18 and older who live in noninstitutional settings. The BRFSS survey has been conducted each calendar year since 1987 with an annual sample size in Maryland that has varied from an average of 4,400 to more than 11,000 interviews annually. In each state, BRFSS data collection/survey administration operations are managed by state health departments that follow guidelines provided by the CDC while the actual surveys are conducted using private survey contractors.

The survey data are transmitted by the survey contractor to the CDC's Office of Surveillance, Epidemiology, and Laboratory Services, Behavioral Risk Factor Surveillance Branch for editing, processing, weighting, and analysis. An edited and weighted data file is provided to each participating health department for each calendar year of data collection.

The CDC BRFSS protocol requires the use of a statewide sampling strategy; it is not designed to produce county-specific estimates. Therefore, the single-year county-specific estimates in this report (see Appendix B) should be used cautiously, as those estimates may or may not be representative of that specific county. Estimates of behaviors or conditions with fewer than 50 respondents in a sample are not reported because they are considered inherently unreliable.

With respect to adults, 'current use' generally refers to use of a tobacco product during the 30 days immediately before completing the BRFSS survey or the respondent stating that he or she now uses the tobacco product every day or on some days. In the case of cigarettes, the respondent must also indicate that they have smoked at least 100 cigarettes in his or her lifetime (about five packs of cigarettes).

INTRODUCTION—ADULT TOBACCO USE BEHAVIORS

Changes in CDC Methodology

1987–2010. The BRFSS survey is a random sample of Maryland adults, so sample results must be weighted by the CDC to match the statewide population of Maryland. Between 1987 and 2010, the CDC used the same methodology for weighting the BRFSS data. During that time period, the CDC used a statistical method called post stratification to weight BRFSS survey data across four categories: (1) age, (2) race and ethnicity, (3) gender, and (4) geographic region within a population.

Changes in 2011. A major change was made with the 2011 CDC BRFSS that prevents data from 2011 and 2012 from being compared with data from 2000–2010. The change was in how the data are weighted—the calculations that make the survey results applicable to the general adult population instead of only survey participants. The new methodology, “iterative proportional fitting” (or “raking”), replaced the post stratification weighting method previously used for the BRFSS. Raking adjusts the data so that groups underrepresented in the survey sample can be more accurately represented in the final dataset than previously possible. The practical impact was that estimates of current cigarette smoking showed an apparent increase from 15.1% in 2010 to 19.1% in 2011.

Changes in 2012. In 2012 the CDC made a much smaller change to its weighting methodology. In 2011, the cell phone sample included those households that only used cell phones; they had no landline phone. In 2012, composition of this group was expanded to include those who reported using cell phones for 90.0% of their calls or more. The net result is that as compared with 2011, more Marylanders were eligible for inclusion in the cell-phone sample and it is likely that the cell phone sample in 2012 had more older adults as compared with 2011. The Maryland BRFSS website made a slight change in how estimates were calculated—consequently, historical statewide and county-level estimates in this report may vary from those previously published. This report presents all data, including historical data, as calculated using this revised methodology.

Data Compatibility

Data collected from 2000 through 2010 can be compared with each other and the observed differences in estimated smoking behaviors between years can be assessed for real change and program and policy impact. That data, however, cannot be compared with data from 2011 and 2012.

The 2012 data can only be compared with 2011 data, and even then the change should be viewed cautiously due to the modification by the CDC to the new weighting methodology it implemented in 2011.

Changes in Maryland Public Health Policies


Since starting its CRF-funded comprehensive tobacco control program in 2001, Maryland has concurrently moved forward and adopted public health policies that furthered the goals of reducing tobacco use, and tobacco-related morbidity, mortality, and health care costs. The more significant advances in state public health policy that are recognized as “*Best Practice*” policies adopted in Maryland during this period include:

- Establishing a minimum package size for cigarettes—prohibiting the sale of single cigarettes (2000)
- Increasing the excise tax on cigarettes so that by 2008 it was \$2.00 per pack
- Adopting comprehensive Clean Indoor Air legislation in 2008 that prohibits smoking indoors at all worksites, restaurants, and bars
- Adding licensure requirements for the sale of tobacco products other than cigarettes (2010)
- Imposing an excise tax on tobacco products other than cigarettes, and increasing it in 2012 to parity with 2008 cigarette excise taxes for nonpremium cigars

TOBACCO USE AND HEALTH

Smoking and the use of tobacco products have short-term and long-term effects on the health of the tobacco user and those who inhale tobacco smoke. Cigarette smoking negatively affects the general health of adults. Figure 16 documents the relationship between smoking status and self-reported general health status at the two ends of the health status spectrum.

Figure 16.
Adult General Health Status by Smoking Status, 2012



Source: CDC Behavioral Risk Factor Surveillance System, Maryland, 2012.

A significantly higher proportion of adults who never smoked cigarettes report “Very Good to Excellent Health” (61.9%), as compared with current cigarette smokers (40.1%). At the other end of the spectrum, a significantly smaller proportion of adults who never smoked cigarettes (11.8%) report “Poor to Fair Health” as compared with current cigarette smokers (24.3%). Data on former smokers fall between the never smoked and current smokers at both ends of the health status spectrum (no adjustment made for length of time since last smoked cigarettes). Adults who have never smoked cigarettes feel the best about their current health status, and those who quit smoking feel better about their health status than those who continue to smoke. This relationship between self-reported health status and smoking status remains across age groups, and income groups, educational attainment, and gender.

Tobacco-Related Cancers and Disease

Tobacco use, including cigarette smoking, causes a substantial number of cancers and disease and/or deaths from those cancers and diseases. Not every person who dies from a tobacco-related cancer or disease does so because of tobacco use or smoking—even lung cancer. Table 3 on the following page details the latest estimated percentage of deaths attributable to cigarette smoking for each of the major tobacco-related cancers and diseases.

Contrary to what many smokers believe, premature death from smoking is not just losing a few years of life when already “old.” A substantial percentage (27.7%) of cigarette smoking-related deaths occur between the ages of 35 and 64. Listed below are the cancers and conditions that account for 50.0% of the deaths among adults ages 35-64 estimated to be the result of cigarette smoking.

Table 2.
Percent of Deaths That Are Attributable to Smoking Cigarettes by Condition
Maryland Adults Ages 35–64

	<u>Females</u>	<u>Males</u>
Bronchitis and Emphysema	80.9%	86.8%
Cancers of the Trachea, Lung, and Bronchus	72.8%	85.3%
Chronic Airway Obstruction	76.5%	79.6%
Cancer of the Larynx	74.2%	78.8%
Cancers of the Lip, Oral Cavity, and Pharynx	69.8%	67.6%
Cancer of the Esophagus	59.8%	65.4%
Aortic Aneurysm	54.8%	58.8%

Sources: Update of Smoking Attributable Fractions from the CDC’s Smoking Attributable Mortality, Morbidity, and Economic Costs (SAMMEC) online estimation tool. Update by the Department of Health and Mental Hygiene, Center for Tobacco Prevention and Control, with technical assistance provided by the CDC SAMMEC staff.

During the 5-year period from 1999–2003, an estimated 6,900 Maryland adults died prematurely every year as a result of current or past cigarette smoking. In contrast, during the 5-year period from 2006–2010 (latest available data), an estimated 6,342 died prematurely every year. This reduction in 558 people annually represents 2,790 premature deaths avoided during 2006–2010.

Still, the number of premature deaths due to cigarette smoking in Maryland is almost three times greater than the combined number of deaths in Maryland for adults ages 35 or older (2,169 in 2007) from unintentional injuries (car accidents, accidental drowning, poisonings), homicides, and cases in which intent could not be established.

ADULT TOBACCO USE AND HEALTH

Table 3.
Percent of Deaths Due to Cigarette Smoking
Maryland Smoking Attributable Fractions, 2006–2010


Cancers and Disease Mortality Causally Related to Cigarette Smoking	FEMALES		MALES	
	Ages 35 to 64	Ages 65+	Ages 35 to 64	Ages 65+
MALIGNANT NEOPLASMS—Cancer Sites				
Lip, Oral Cavity, and Pharynx	48.6%	44.0%	69.8%	67.6%
Esophagus	59.8%	54.1%	65.4%	70.0%
Stomach	11.6%	12.4%	22.3%	24.9%
Pancreas	24.6%	22.8%	20.7%	15.7%
Larynx	74.2%	70.9%	78.8%	79.9%
Trachea, Lung, and Bronchus	72.8%	68.7%	85.3%	85.6%
Cervix Uteri	11.2%	8.9%	-	-
Kidney and Renal Pelvis	5.5%	4.0%	32.6%	34.8%
Urinary Bladder	28.6%	29.3%	40.2%	43.5%
Acute Myeloid Leukemia	9.8%	12.5%	18.9%	19.8%
CARDIOVASCULAR DISEASES				
Ischemic Heart Disease	28.8%	10.6%	32.2%	13.4%
Other Heart Disease	9.9%	8.2%	16.0%	15.4%
Cerebrovascular Disease	35.4%	4.8%	28.3%	6.7%
Atherosclerosis	11.7%	6.4%	24.9%	22.7%
Aortic Aneurysm	54.8%	46.6%	58.8%	60.6%
Other Arterial Diseases	17.6%	12.1%	15.5%	8.3%
RESPIRATORY DISEASES				
Pneumonia and Influenza	17.3%	11.6%	18.2%	20.3%
Bronchitis and Emphysema	80.9%	82.3%	86.8%	90.2%
Chronic Airway Obstruction	76.5%	75.1%	75.9%	79.6%

Sources: Update of Smoking Attributable Fractions from the CDC's Smoking Attributable Mortality, Morbidity, and Economic Costs (SAMMEC) online estimation tool. Update by the Department of Health and Mental Hygiene, Center for Tobacco Prevention and Control, with technical assistance provided by the CDC SAMMEC staff.

Cost of Treating Tobacco-Related Cancers and Disease

Even as the number of Maryland adults suffering from cancers and disease caused by cigarette smoking declines, total medical expenditures to treat these conditions has continued to grow. The cost of medical treatment in 2000 is estimated at \$1.4 billion and at \$2.7 billion in 2009. This is a 93.0% increase (without adjustment for inflation). The estimated annual total cost of treating cancers and disease caused by smoking (2000–2009) is presented in Figure 17 below, along with a trend-line through 2020 based upon the growth during 2000–2009. By 2015, the cost is estimated to be \$3.5 billion and, by 2020, \$4.5 billion.

Figure 17.
Estimated Maryland Smoking-Related Medical Expenditures, 2000–2009
Projected Maryland Smoking-Related Medical Expenditures, 2010–2020



Sources: Update of Smoking Attributable Medical Expenditures from the CDC's SAMMEC online estimation tool. Update by the Department of Health and Mental Hygiene, Center for Tobacco Prevention and Control, with technical assistance provided by the CDC SAMMEC staff.

The increasing cost of treating cancers of the lung, trachea, and bronchus is illustrative of how, even as the number of cases decreases, the costs of treatment continue to climb. In 2000, there were 1,362 males ages 35+ admitted to Maryland hospitals with a primary diagnosis of these cancers. By 2011, the number had decreased to 1,207. In 2000, the average cost per admission with this primary diagnosis was just over \$14,000 (males). In 2011, the average cost was almost \$23,000, an increase of almost 63%. The total cost for treating these cases increased from \$19.2 million to \$28.5 million.

Aside from initiatives to control the cost of health care generally, the only available means of reducing tobacco and smoking-related health care expenditures is to prevent youth and young adults from starting to smoke, and to encourage current smokers to quit and to assist them in doing so successfully.

Preventable Hospitalizations Resulting from Tobacco-Related Cancers and Disease

As noted in the previous section, cancers and disease caused by smoking are preventable. Maryland has made a long-term commitment to reducing the initiation of tobacco use and assisting those who use tobacco to successfully quit permanently. Success is beginning to result in fewer tobacco-related inpatient hospitalizations.

In 2000, there were 62,569 such admissions as compared with 55,672 in 2011, a decrease of 6,897 or 11.0% in tobacco-related primary and secondary diagnosis admissions. More than 53.0% of the decrease was in primary diagnosis admissions. This decrease in tobacco-related primary and secondary admissions is estimated to have saved more than \$102 million in hospital charges in 2011.¹


This space was left blank intentionally.

¹Data from the Maryland Health Services Cost Review Commission (HSCRC) for 2000 and 2011 were analyzed for the tobacco-related cancers and diseases in Table 2. First, only admissions with those cancers and diseases listed in the primary diagnosis field were analyzed. The associated smoking-attributable fraction (SAF) for each was applied, resulting in an estimate of tobacco-related primary admissions. Second, only admissions with those cancers and diseases listed in the secondary diagnosis field were analyzed, and the appropriate SAF applied, resulting in an estimate of tobacco-related secondary admissions.

Adult Current Tobacco Use

Most Maryland adults do not smoke or use other tobacco products. In 2011, an estimated 19.4% of Maryland adults used some form of tobacco product. Cigarettes were the most popular tobacco product (16.2%), followed by cigars (4.4%), smokeless tobacco (2.0%), and any other tobacco products (1.7%) such as pipes, bidis, kreteks, etc. Cigarettes are the only tobacco product with annual data from the CDC. In 2012, Maryland was able to include for the first time cigar smoking as a state-added variable to the CDC's BRFSS.

Figure 18.
Use of Tobacco Products
Maryland Adults Ages 18 and Older, 2000–2012



Source: CDC Behavioral Risk Factor Surveillance System (Maryland), 2000–2012.

Because the CDC changed its methodology for weighting BRFSS data beginning in 2011, data collected before 2011 cannot be compared to data collected in 2011 and after. It is believed that the new methodology more accurately estimates the overall prevalence of cigarette smoking than did the previous method. The break in Figure 18 between 2010 and 2011 highlights this fact. In 2012, the CDC made a minor adjustment to its methodology, but states that the 2012 data are comparable to that of 2011 (Note: This report provides estimates for adult tobacco use through 2012 only. The 2013 BRFSS adult data were not publicly released until August 2014, after production of this report).


The decrease in adult cigarette smoking in Maryland between 2011 and 2012 is statistically significant and coincides with (1) a 3-month intensive national media campaign by the CDC to encourage smokers to quit and nonsmokers to never start, and (2) sharp increases in gasoline prices that may have impacted cigarette purchases by reducing consumer funds available for such purchases. Data from the Maryland

ADULT CURRENT TOBACCO USE

Comptroller's Office show decreased cigarette sales between 2011 and 2012 which provides an independent perspective for the decrease shown in Figure 18.


In Figure 19, between 2011 and 2012, current cigarette smoking by both adult females and males decreased (not statistically significant). In 2012, the difference in smoking between the genders is not statistically significant. In Figure 20, between 2011 and 2012, current cigarette smoking by non-Hispanic Blacks and non-Hispanic White adults decreased, although that decrease was not statistically significant. In 2012, the difference in the estimated prevalence of cigarette smoking between races was not statistically significant.

Figure 19.
Current Cigarette Smoking
Maryland Adults Ages 18 and Older—by Gender, 2000–2012



Source: CDC Behavioral Risk Factor Surveillance System (Maryland), 2000–2012.

Figure 20.
Current Cigarette Smoking
Maryland Adults Ages 18 and Older—by Race, 2000–2012




Source: CDC Behavioral Risk Factor Surveillance System (Maryland), 2000–2012.

ADULT CURRENT TOBACCO USE

As shown in Figure 21, educational attainment is inversely correlated with cigarette smoking—the lower the educational attainment status, the higher the prevalence of cigarette smoking. Adults whose highest educational attainment fell short of a high school diploma or GED have the highest rates of cigarette smoking in 2012 (29.1%) and those who graduated with a four-year college degree or higher have the lowest rates of smoking (6.2%). Between 2011 and 2012, none of the decreases within an educational attainment category were statistically significant.


Figure 21.
Current Cigarette Smoking
Maryland Adults Ages 18 and Older—by Highest Educational Attainment, 2011 and 2012



Source: CDC Behavioral Risk Factor Surveillance System (Maryland), 2011 and 2012.

As shown in Figure 22 on the next page, annual household income is inversely correlated with cigarette smoking—the lower the annual household income, the higher the prevalence of cigarette smoking. Adults whose annual household income was less than \$15,000 had the highest rates of cigarette smoking in 2012 (30.7%) and those whose annual income was \$50,000 or more had the lowest rates of smoking (11.6%). Between 2011 and 2012 none of the decreases within annual household income categories were statistically significant.

Figure 22.
Current Cigarette Smoking
Maryland Adults Ages 18 and Older—by Annual Household Income, 2011 and 2012




Source: CDC Behavioral Risk Factor Surveillance System (Maryland), 2011–2012.

QUITTING TOBACCO USE

Quitting Smoking

Though many Maryland adults are addicted to smoking, most actually want to quit. Asked whether they seriously planned to quit smoking, 74.0% of adult cigarette smokers said that they want to quit smoking, and 60.1% reported a serious quit attempt during the past 12 months. Figure 23 details the timeframes these smokers gave for quitting smoking—88.5% stated that they want to quit within the next 12 months.

Figure 23.
Timeline for Quitting Smoking
Among Adult Maryland Cigarette Smokers Who Want to Quit, 2012



Source: CDC Behavioral Risk Factor Surveillance System (Maryland), 2012.

There is no statistically significant difference in the overall desire of current smokers to quit by gender, although males (27.0%) were significantly more likely to state that they wanted to quit smoking during the next 30 days than were females (14.7%). Similarly there is no statistically significant difference in the overall desire of current smokers to quit between non-Hispanic Whites and non-Hispanic Blacks—and no significant differences in their timelines for quitting. Likewise, there are no statistically significant differences in the overall desire of current smokers to quit based on income or highest educational attainment.


It is not easy for a smoker to overcome his or her addiction to nicotine. The average smoker has to make multiple serious attempts to quit before succeeding, and even then, is subject to relapse. Many of the smokers stating a desire to quit smoking in the short-term may be stating what they know that they “should

ADULTS QUITTING TOBACCO USE

do” as opposed to what they actually will do to quit smoking during the next 30 days or within the next year. Supporting smokers with services such as the Maryland Tobacco Quitline (1-800-QUIT-NOW) dramatically increases success in quitting.

In 2012, the highest rate of quitting cigarette smoking (former smokers divided by those who ever smoked) were non-Hispanic White adults (61.5%), followed by adult males (60.5%), then adult females (57.1%), and then by non-Hispanic Black adults (54.3%)—although there has been sharp improvement in recent years. Non-Hispanic Black adults have both the highest rate of adults who never smoked cigarettes and a significantly higher rate of serious quit attempts in the past year (69.4%, 2012), compared with non-Hispanic Whites (55.1%, 2012). At the same time, as shown in Figure 24, the percentage of non-Hispanic Black adult smokers who have quit has remained consistently lower than the percentage of non-Hispanic White adult smokers over time.

Figure 24.
Percent of Maryland Adults Who Ever Smoked Cigarettes and Have Quit, by Race, 2000–2012



Source: CDC Behavioral Risk Factor Surveillance System (Maryland), 2000–2012.

It is not unusual for a “former” smoker to relapse and start smoking again. Therefore, it is important to also examine how long former smokers have abstained. The longer the period of nonsmoking the more likely it will continue.


Figure 25 documents the critical impact that mass media campaigns can have in motivating current smokers who want to quit (74.0% in Maryland) to take action and seek help. The CDC’s Tips from Former Smokers campaign (TIPS) that ran in Maryland media markets March through June 2012 and 2013 prompted a sizable increase in registered users of the Maryland Tobacco Quitline (1-800-QUIT-NOW) and its online counterpart. The TIPS campaign consisted of media spots running in local markets combined with national

ADULTS QUITTING TOBACCO USE

media buys that ran alternate weeks. The weeks when both national and local media spots were aired resulted in the highest call volume to the Quitline.

In evaluations of the Maryland Tobacco Quitline (1-800-QUIT-NOW), 92.0% of respondents stated that it met their expectations, and 97.5% would recommend the Maryland Tobacco Quitline to others. Of those calling the Maryland Tobacco Quitline, almost 82.0% reported making a serious attempt to quit smoking as a result. Of those participating in the evaluation, almost 30.0% reported being smoke-free at 30 days after completion of the Maryland Tobacco Quitline program. This is in contrast to national averages of 4–7% of smokers successfully quitting without assistance. More than 78.0% of calls to the Maryland Tobacco Quitline are from persons with a high school education or higher. Callers are evenly split between non-Hispanic White (46.9%) and non-Hispanic Black (47.0%). Just over 60% of callers are female (60.6%).

Figure 25.
Monthly Registered Callers/Web Users of Maryland Quitline, 2011–2013




Source: Maryland 1-800-QUIT-NOW Registered callers (including those accessing via Internet beginning December 2012).

Maryland’s health messaging component (titled in statute as the “Counter-marketing and Media” component) of Maryland’s comprehensive tobacco control program could significantly increase the number of Maryland residents that it helps to succeed in quitting smoking, saving many more lives and further decreasing expenditures on tobacco-related health care.

As shown in Figure 26, in 2012 a significant majority (60.0%) of adult former cigarette smokers reported that they had not smoked a cigarette, not even a puff, for 10 years or more. Almost 14.0% of adult former smokers reported that their last cigarette was sometime during the past 12 months—they are at the highest risk of relapse.


ADULTS QUITTING TOBACCO USE

Figure 26.
Years Since Smoked Last Cigarette, Former Adult Smokers, 2012



Source: CDC Behavioral Risk Factor Surveillance System (Maryland), 2000–2012.

Figure 27.
Smoking Status of Maryland Adults, 2000–2012




Source: CDC Behavioral Risk Factor Surveillance System (Maryland), 2000–2012.

INITIATION OF TOBACCO USE

Increasing the number and proportion of people who have never smoked or used tobacco is an essential component of Maryland's objective of reducing smoking and tobacco use. In Figure 27, the smoking status of Maryland adults is tracked over time, and it is clear that preventing underage youth and young adults from ever initiating smoking should be a high priority.

As the proportion of adults who have never smoked increases, it can create a social norm where nonsmoking is the predominant behavior. Since 2000, there has been an increase in the proportion of Maryland adults reporting that they have never smoked cigarettes—moving nonsmoking closer to being the normative behavior (see Figure 28).

Figure 28.
Maryland Adults Who Have Never Smoked Cigarettes, 2000–2012




Source: CDC Behavioral Risk Factor Surveillance System (Maryland), 2000–2012.

This changing social norm, however, is not evenly distributed among subpopulations or even across the state. Adult females are significantly more likely than adult males never to have smoked cigarettes. Adult non-Hispanic Blacks are significantly more likely than adult non-Hispanic Whites never to have smoked cigarettes.

These differences are also present when considering highest educational attainment and household income. As shown in Figure 29, adults who have four or more years of college are significantly more likely to have never smoked cigarettes as compared with adults with some college or post-high school learning, adults with a high school diploma or GED, or those adults without a high school diploma or GED. Households with annual income of \$75,000 a year or more are significantly more likely to never have smoked cigarettes as compared with households with less annual income.

ADULT INITIATION OF TOBACCO USE

Figure 29.
Maryland Adults Who Have Never Smoked Cigarettes, 2000–2012
 By Gender, Race, Highest Educational Attainment, Annual Household Income



Source: CDC Behavioral Risk Factor Surveillance System (Maryland), 2000–2012.

ADULT INITIATION OF TOBACCO USE

In 2012, for the first time, BRFSS asked questions about the age at which smokers, age 30 or under, had (1) smoked a cigarette for the first time, and (2) started smoking cigarettes regularly. Overall, 66.8% reported that they had first smoked a cigarette before they were age 18. Approximately one-half (49.0%) of those who reported first trying cigarette smoking before they were 18 said that they actually had tried cigarette smoking before they were 15.


In 2012, 0.5% of Maryland adults reported that they had started smoking cigarettes during the past 12 months. Most were between the ages of 18 to 24. This suggests that efforts to reduce the initiation of cigarette smoking and tobacco use should therefore focus on underage youth and young adults between the ages of 18 and 24.

ADULT EXPOSURE TO SECONDHAND SMOKE

In previous reports, adult secondhand smoke data were presented that originated from the Maryland Adult Tobacco Survey (MATS). However, funding for that survey was discontinued after the fall 2008 survey, leaving only the BRFSS survey as a mechanism to collect adult smoking behavior data. From 2009 through 2011, the BRFSS did not include questions that would support presentation of secondhand smoke data.

In 2012, however, the Center for Tobacco Prevention and Control was able to insert a very limited number of questions into the BRFSS survey and the data collected are presented in Figure 30. Please note that apparent differences by race are not statistically significant, while differences by rent/ownership are statistically significant.

Figure 30.
Percentage Maryland Households with Children and Resident Adult Smokers, 2012



Source: CDC Behavioral Risk Factor Surveillance System (Maryland), 2012.

Table of Contents

APPENDIX A—UNDERAGE YOUTH PREVALENCE, INITIATION, AND CESSATION

	<u>Pages</u>
Maryland	41
Allegany County	43
Anne Arundel County	45
Baltimore City	47
Baltimore County	49
Calvert County	51
Caroline County	53
Carroll County	55
Cecil County	57
Charles County	59
Dorchester County	61
Frederick County	63
Garrett County	65
Harford County	67
Howard County	69
Kent County	71
Montgomery County	73
Prince George’s County	75
Queen Anne’s County	77
Somerset County	79
St. Mary’s County	81
Talbot County	83
Washington County	85
Wicomico County	87
Worcester County	89

APPENDIX A—UNDERAGE YOUTH PREVALENCE, INITIATION, AND CESSATION

Statewide and Jurisdiction Data


The data presented in this report from 2000 through 2010 are from the Maryland Youth Tobacco Survey (MYTS). The MYTS was conducted biennially in the fall of even calendar years except 2004 when no survey was conducted due to budget constraints. The MYTS was designed to produce county-level and statewide estimates of key tobacco use behaviors by students in public middle schools (grades 6 through 8) and by students in public high schools (grades 9 through 12).

The MYTS was one of three risk behavior surveys being conducted—the Maryland Adolescent Survey (MAS) which was a county-level survey of students in grades 6, 8, 10, and 12 conducted biennially in the spring, the MYTS, and beginning in 2005 the Youth Risk Behavior Survey (YRBS) conducted biennially in the spring at the statewide level. DHMH and MSDE began a collaborative process with the goal of reducing the survey burden on Maryland. It was essential that key indicators of risk behavior in grades 6 through 12 were available at both the county and state levels.

The result is a single survey, the MYTRBS which combines the MYTS and the YRBS; the MAS was discontinued. The first administration of this survey, conducted in the spring of 2013, produced the 2013 data in this report and in Appendix A—Underage Youth Prevalence, Initiation, And Cessation. The MYTRBS uses the same methodology and survey model as the MYTS since 2000, and the data can be compared across all years reported. Beginning with the fall of 2014, the MYTRBS will be administered in the fall of even calendar years.

The MYTRBS uses core and optional questions from the CDC’s state and national Youth Tobacco Surveys, state and national Youth Risk Behavior Surveys, and a very few select Maryland-specific questions to address unique indicators required by the Cigarette Restitution Fund Tobacco Use Prevention and Cessation Program legislation (CRF Tobacco Indicators, CRF-TI). Whenever possible the MYTRBS uses 2000-10 MYTS questions with respect to CRF-TI.

It was not possible to use the 2010 MYTS questions with respect to (1) past 12-month initiation of any tobacco use and (2) past 12-month cessation of tobacco use. The 2000–2010 MYTS asked seven separate questions about past 12-month initiation (one for cigarettes, cigars, smokeless, pipe tobacco, bidis, and kreteks), and seven questions with respect to past 12-month cessation. Due to survey protocol requirements established by the CDC limiting the number of survey questions to 99, the survey instrument had no room for 14 questions to create data for two indicators. Instead, one question asked youth whether they had initiated use of any kind of tobacco for the first time during the past 12 months, and a second question asked the same about cessation during the past 12 months. Therefore, the data in this report with respect to those two indicators may reflect both actual changes in tobacco use behavior and changes due to the fact that the questions asked about the behaviors differently. The new questions were cognitively tested by the CDC’s YRBS survey contractor before being used to ensure that students reading the questions understood what was being asked and that they could provide appropriate responses. Separate data specific to each product (pipe tobacco, bidis, and kreteks) was not collected in 2010 due to space limitations in the survey instrument. Instead a question was asked about the use of any of those products.



Maryland

Tobacco Use by Youth: 2000–2013

Maryland Department of Health and Mental Hygiene
Center for Tobacco Prevention and Control

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Any Tobacco Product (All Youth Populations)	21.4 $\pm 1.2\%$ 87,963	18.5 $\pm 1.0\%$ 80,831	15.6 $\pm 0.6\%$ 70,611	16.2 $\pm 0.5\%$ 68,959	17.1 $\pm 0.5\%$ 74,173	*12.9 $\pm 0.6\%$ 52,477
Any Tobacco Product (Minority Youth)	18.9 $\pm 1.4\%$ 33,913	16.9 $\pm 1.2\%$ 35,514	14.4 $\pm 0.8\%$ 34,560	16.3 $\pm 0.7\%$ 35,129	17.9 $\pm 0.7\%$ 43,110	*12.6 $\pm 0.8\%$ 27,567
Cigarette	15.7 $\pm 1.1\%$ 61,808	12.5 $\pm 0.8\%$ 51,993	10.0 $\pm 0.5\%$ 43,749	10.2 $\pm 0.4\%$ 42,578	9.6 $\pm 0.4\%$ 40,945	*7.9 $\pm 0.4\%$ 30,365
Cigar	8.8 $\pm 0.6\%$ 35,268	7.5 $\pm 0.5\%$ 31,622	6.5 $\pm 0.4\%$ 28,323	10.6 $\pm 0.4\%$ 44,064	9.5 $\pm 0.4\%$ 40,752	8.3 $\pm 0.4\%$ 33,066
Pipe (tobacco)	3.9 $\pm 0.4\%$ 15,670	4.9 $\pm 0.4\%$ 20,509	4.7 $\pm 0.3\%$ 20,847	5.9 $\pm 0.4\%$ 24,263	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2013
Bidi	5.3 $\pm 0.5\%$ 21,077	5.4 $\pm 0.4\%$ 22,554	4.8 $\pm 0.4\%$ 21,247	4.0 $\pm 0.3\%$ 16,201	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2013
Kretek	3.8 $\pm 0.3\%$ 15,285	4.3 $\pm 0.4\%$ 18,163	5.1 $\pm 0.4\%$ 22,766	3.6 $\pm 0.3\%$ 14,787	KRETEK SPECIFIC DATA NOT AVAILA- BLE FOR 2010	KRETEK SPECIFIC DATA NOT AVAILA- BLE FOR 2013
Smokeless Tobacco	3.5 $\pm 0.3\%$ 14,109	3.7 $\pm 0.3\%$ 15,511	3.6 $\pm 0.3\%$ 16,605	3.7 $\pm 0.2\%$ 14,910	3.3 $\pm 0.2\%$ 13,746	*5.2 $\pm 0.4\%$ 20,934

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	19.5 $\pm 1.0\%$ 80,050	16.2 $\pm 0.8\%$ 70,879	15.0 $\pm 0.6\%$ 67,969	14.8 $\pm 0.4\%$ 62,909	18.1 $\pm 0.5\%$ 78,621	*12.4 $\pm 0.6\%$ 48,067
Minority Youth	16.7 $\pm 1.0\%$ 29,886	13.7 $\pm 0.9\%$ 28,873	12.4 $\pm 0.7\%$ 29,729	13.2 $\pm 0.5\%$ 28,346	18.9 $\pm 0.6\%$ 45,605	*12.2 $\pm 0.8\%$ 25,157

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	35.1 $\pm 1.9\%$ 16,304	36.0 $\pm 1.7\%$ 12,657	40.2 $\pm 1.9\%$ 11,518	38.0 $\pm 1.8\%$ 13,498	45.0 $\pm 1.8\%$ 12,442	*45.1 $\pm 1.4\%$ 32,768
Minority Youth	44.5 $\pm 4.0\%$ 6,629	43.4 $\pm 3.1\%$ 5,496	47.7 $\pm 3.1\%$ 5,960	44.9 $\pm 2.7\%$ 7,882	49.7 $\pm 2.5\%$ 7,402	48.9 $\pm 1.8\%$ 19,291

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (\pm) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined; between 2000 and 2013 preceded by an asterisk (*).


CURRENT CIGARETTE SMOKING — School Type, Gender, and Minority Populations


Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Middle School (MS)	7.2% $\pm 0.9\%$ 13,134	5.2% $\pm 0.5\%$ 10,110	3.7% $\pm 0.5\%$ 7,029	3.5% $\pm 0.4\%$ 6,388	3.5% $\pm 0.4\%$ 6,294	*3.9% $\pm 0.4\%$ 6,717
MS Females	7.2% $\pm 1.1\%$ 6,319	5.1% $\pm 0.7\%$ 4,798	3.2% $\pm 0.6\%$ 2,936	2.9% $\pm 0.5\%$ 2,584	3.1% $\pm 0.5\%$ 2,929	*3.5% $\pm 0.4\%$ 2,955
MS Males	7.2% $\pm 1.2\%$ 6,679	5.4% $\pm 0.7\%$ 5,233	4.2% $\pm 0.7\%$ 4,017	4.1% $\pm 0.6\%$ 3,793	3.8% $\pm 0.5\%$ 3,328	*4.2% $\pm 0.6\%$ 3,632
MS Minorities	6.5% $\pm 1.3\%$ 5,221	5.3% $\pm 0.8\%$ 5,057	3.9% $\pm 0.7\%$ 3,924	3.9% $\pm 0.5\%$ 3,625	4.1% $\pm 0.6\%$ 4,179	*4.5% $\pm 0.6\%$ 4,151
High School (HS)	23.0% $\pm 0.9\%$ 48,674	18.7% $\pm 0.8\%$ 41,822	14.7% $\pm 0.7\%$ 36,720	15.3% $\pm 0.5\%$ 36,190	14.1% $\pm 0.6\%$ 34,651	*11.0% $\pm 0.6\%$ 23,586
HS Females	23.4% $\pm 1.3\%$ 24,702	18.7% $\pm 1.0\%$ 21,121	13.7% $\pm 0.9\%$ 17,315	13.2% $\pm 0.6\%$ 15,589	11.7% $\pm 0.7\%$ 14,966	*9.4% $\pm 0.6\%$ 10,098
HS Males	22.4% $\pm 1.2\%$ 23,551	18.4% $\pm 1.0\%$ 20,113	15.6% $\pm 0.8\%$ 19,299	17.4% $\pm 0.7\%$ 20,444	16.6% $\pm 0.8\%$ 19,580	*12.2% $\pm 0.6\%$ 12,753
HS Minorities	16.0% $\pm 1.3\%$ 14,204	13.5% $\pm 0.9\%$ 13,884	11.1% $\pm 0.8\%$ 14,390	12.7% $\pm 0.6\%$ 14,782	12.2% $\pm 0.6\%$ 16,446	*8.9% $\pm 0.6\%$ 10,109

Maryland—Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Allegany County

Tobacco Use by Youth: 2000–2013

Maryland Department of Health and Mental Hygiene
Center for Tobacco Prevention and Control

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Any Tobacco Product (All Youth Populations)	30.1% $\pm 3.7\%$ 1,688	26.8% $\pm 3.8\%$ 1,468	22.5% $\pm 3.0\%$ 1,145	20.1% $\pm 2.6\%$ 964	19.1% $\pm 2.5\%$ 874	*17.8% $\pm 3.3\%$ 829
Any Tobacco Product (Minority Youth)	30.8% $\pm 7.3\%$ 154	33.7% $\pm 7.7\%$ 186	29.1% $\pm 6.3\%$ 174	32.1% $\pm 5.8\%$ 160	25.1% $\pm 5.3\%$ 157	19.0% $\pm 11.4\%$ 213
Cigarette	24.6% $\pm 3.4\%$ 1,320	21.3% $\pm 3.6\%$ 1,132	16.6% $\pm 2.5\%$ 826	14.9% $\pm 2.2\%$ 705	13.5% $\pm 2.1\%$ 606	*13.2% $\pm 2.7\%$ 590
Cigar	9.3% $\pm 1.5\%$ 511	8.0% $\pm 1.8\%$ 433	9.4% $\pm 1.7\%$ 470	10.9% $\pm 1.9\%$ 520	9.1% $\pm 1.7\%$ 410	9.3% $\pm 2.0\%$ 426
Pipe (tobacco)	4.7% $\pm 1.4\%$ 260	4.4% $\pm 1.2\%$ 238	4.1% $\pm 1.0\%$ 205	5.3% $\pm 1.3\%$ 252	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Bidi	4.4% $\pm 1.1\%$ 243	5.6% $\pm 1.3\%$ 302	4.1% $\pm 1.1\%$ 209	3.5% $\pm 0.9\%$ 166	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010	BIDI SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Kretek	3.3% $\pm 1.0\%$ 181	3.6% $\pm 1.1\%$ 197	4.1% $\pm 1.1\%$ 208	3.1% $\pm 0.8\%$ 145	KRETEK SPECIFIC DATA NOT AVAILABLE FOR 2010	KRETEK SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Smokeless Tobacco	9.1% $\pm 1.6\%$ 503	8.6% $\pm 2.0\%$ 460	6.1% $\pm 1.3\%$ 305	7.0% $\pm 1.3\%$ 327	6.6% $\pm 1.3\%$ 292	9.3% $\pm 2.0\%$ 427

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	27.4% $\pm 2.8\%$ 1,536	21.9% $\pm 2.6\%$ 1,204	22.6% $\pm 2.8\%$ 1,151	19.5% $\pm 2.3\%$ 933	19.1% $\pm 2.3\%$ 876	*16.6% $\pm 3.1\%$ 745
Minority Youth	24.5% $\pm 5.6\%$ 122	22.7% $\pm 6.4\%$ 125	17.8% $\pm 4.5\%$ 107	23.6% $\pm 5.3\%$ 117	24.8% $\pm 4.6\%$ 155	14.0% $\pm 8.8\%$ 152

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	31.2% $\pm 4.7\%$ 336	26.2% $\pm 6.4\%$ 205	30.1% $\pm 5.5\%$ 165	30.6% $\pm 6.6\%$ 174	34.8% $\pm 7.8\%$ 129	33.4% $\pm 4.5\%$ 364
Minority Youth	20.5% $\pm 15.2\%$ 23	Not Available	Not Available	25.1% $\pm 11.6\%$ 25	18.6% $\pm 11.2\%$ 12	31.4% $\pm 9.8\%$ 76

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (\pm) appear immediately following prevalence estimates.
- Statistically significant change between survey years is underlined; between 2000 and 2013 preceded by an asterisk (*).


CURRENT CIGARETTE SMOKING — School Type, Gender, and Minority Populations

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Middle School (MS)	9.9% \pm 2.9% 229	8.3% \pm 2.3% 195	6.9% \pm 2.5% 140	3.9% \pm 1.4% 77	6.4% \pm 2.2% 123	5.5% \pm 2.4% 121
MS Females	8.5% \pm 3.2% 94	9.4% \pm 3.2% 105	7.7% \pm 4.4% 76	3.8% \pm 1.9% 37	6.0% \pm 2.3% 60	5.7% \pm 2.4% 57
MS Males	11.3% \pm 3.9% 135	7.3% \pm 2.8% 89	6.2% \pm 2.9% 64	4.1% \pm 1.5% 41	6.8% \pm 3.0% 62	5.3% \pm 3.1% 62
MS Minorities	15.3% \pm 7.3% 40	11.9% \pm 6.1% 35	7.0% \pm 4.7% 20	10.7% \pm 5.1% 25	12.6% \pm 6.7% 40	5.9% \pm 6.1% 43
High School (HS)	35.6% \pm 3.7% 1,092	31.6% \pm 4.9% 937	23.4% \pm 2.5% 686	22.8% \pm 3.6% 627	18.7% \pm 3.0% 483	*20.7% \pm 2.7% 469
HS Females	37.7% \pm 4.5% 575	34.8% \pm 5.9% 525	24.8% \pm 3.5% 347	22.5% \pm 4.4% 299	16.5% \pm 3.2% 223	*17.4% \pm 2.9% 197
HS Males	33.2% \pm 4.4% 505	28.0% \pm 5.7% 405	22.0% \pm 3.1% 337	23.0% \pm 4.4% 328	21.1% \pm 4.7% 258	*23.4% \pm 3.9% 263
HS Minorities	37.0% \pm 10.1% 78	44.9% \pm 13.9% 95	26.5% \pm 6.3% 72	40.5% \pm 7.8% 101	28.6% \pm 6.7% 82	32.5% \pm 6.3% 112

Allegany County—Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Anne Arundel County

Tobacco Use by Youth: 2000–2013

Maryland Department of Health and Mental Hygiene
Center for Tobacco Prevention and Control

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Any Tobacco Product (All Youth Populations)	24.8% ± 4.5% 9,349	20.4% ± 3.4% 8,030	16.5% ± 2.1% 6,465	18.4% ± 2.1% 6,910	18.2% ± 3.2% 7,016	*12.7% ± 2.4% 4,705
Any Tobacco Product (Minority Youth)	24.9% ± 5.5% 2,254	18.7% ± 4.7% 2,044	16.9% ± 3.0% 2,256	21.4% ± 3.3% 2,559	21.9% ± 3.0% 3,048	*14.2% ± 2.9% 1,894
Cigarette	19.8% ± 4.0% 7,203	14.7% ± 3.0% 5,588	11.1% ± 1.6% 4,253	12.2% ± 1.5% 4,517	10.4% ± 2.2% 3,928	*8.5% ± 1.8% 3,014
Cigar	10.7% ± 2.3% 3,946	9.0% ± 1.9% 3,445	7.3% ± 1.1% 2,773	12.0% ± 1.5% 4,431	9.6% ± 2.2% 3,664	8.0% ± 1.6% 2,934
Pipe (tobacco)	4.4% ± 1.1% 1,620	5.5% ± 1.0% 2,100	5.4% ± 1.1% 2,096	6.6% ± 1.1% 2,424	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Bidi	5.2% ± 1.2% 1,895	5.2% ± 1.0% 1,989	5.3% ± 1.1% 2,052	4.4% ± 0.8% 1,615	BIDI SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	BIDI SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Kretek	4.7% ± 1.1% 1,730	4.7% ± 1.0% 1,803	5.0% ± 1.0% 1,938	3.9% ± 0.8% 1,441	KRETEK SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	KRETEK SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Smokeless Tobacco	3.5% ± 1.1% 1,306	4.0% ± 0.9% 1,499	3.9% ± 0.8% 1,500	3.5% ± 0.7% 1,256	3.3% ± 0.8% 1,222	4.5% ± 1.0% 1,674

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	23.5% ± 3.2% 8,883	18.1% ± 2.7% 7,116	16.5% ± 1.9% 6,485	15.4% ± 1.5% 5,807	18.0% ± 3.2% 6,931	*12.3% ± 2.2% 4,349
Minority Youth	21.8% ± 3.9% 1,963	15.3% ± 3.2% 1,671	14.9% ± 2.3% 1,989	15.2% ± 2.2% 1,812	20.0% ± 3.4% 2,792	*13.0% ± 2.7% 1,641

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	31.0% ± 4.7% 1,684	32.3% ± 5.6% 1,159	40.3% ± 5.1% 1,144	34.6% ± 4.9% 1,234	41.3% ± 6.3% 1,053	*45.0% ± 3.7% 2,960
Minority Youth	37.7% ± 9.5% 410	28.7% ± 10.8% 165	44.8% ± 9.7% 408	37.9% ± 9.3% 483	44.3% ± 9.5% 402	43.8% ± 5.3% 1,141

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (± %) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2013 preceded by an asterisk (*).


CURRENT CIGARETTE SMOKING — School Type, Gender, and Minority Populations


Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Middle School (MS)	9.4% $\pm 3.1\%$ 1,556	6.0% $\pm 2.0\%$ 1,055	3.7% $\pm 1.2\%$ 602	4.4% $\pm 2.0\%$ 703	3.3% $\pm 1.5\%$ 540	*3.6% $\pm 1.6\%$ 583
MS Females	11.1% $\pm 4.3\%$ 873	6.6% $\pm 2.6\%$ 568	3.3% $\pm 1.7\%$ 263	4.2% $\pm 2.9\%$ 333	3.4% $\pm 2.0\%$ 291	*3.4% $\pm 1.8\%$ 281
MS Males	7.7% $\pm 3.0\%$ 667	5.3% $\pm 2.4\%$ 465	4.0% $\pm 1.7\%$ 330	4.6% $\pm 1.8\%$ 370	3.1% $\pm 1.5\%$ 249	3.7% $\pm 2.2\%$ 295
MS Minorities	9.4% $\pm 4.6\%$ 343	5.6% $\pm 3.0\%$ 266	4.5% $\pm 2.4\%$ 243	7.4% $\pm 3.6\%$ 381	5.0% $\pm 2.1\%$ 308	4.4% $\pm 2.5\%$ 262
High School (HS)	28.5% $\pm 3.2\%$ 5,647	22.5% $\pm 2.3\%$ 4,533	16.7% $\pm 1.7\%$ 3,651	18.3% $\pm 2.1\%$ 3,814	15.9% $\pm 1.9\%$ 3,388	*12.7% $\pm 1.8\%$ 2,431
HS Females	28.4% $\pm 3.7\%$ 2,769	23.2% $\pm 2.8\%$ 2,378	16.2% $\pm 2.2\%$ 1,797	16.0% $\pm 2.3\%$ 1,674	13.7% $\pm 2.1\%$ 1,496	*11.2% $\pm 2.0\%$ 1,067
HS Males	28.6% $\pm 4.2\%$ 2,864	21.1% $\pm 3.0\%$ 2,131	17.0% $\pm 2.1\%$ 1,826	20.3% $\pm 2.6\%$ 2,116	18.2% $\pm 2.5\%$ 1,892	*13.7% $\pm 2.2\%$ 1,287
HS Minorities	23.8% $\pm 5.1\%$ 1,152	15.9% $\pm 2.9\%$ 875	15.6% $\pm 2.4\%$ 1,149	16.0% $\pm 2.6\%$ 1,022	15.3% $\pm 2.2\%$ 1,132	*13.4% $\pm 2.4\%$ 871

Anne Arundel County—Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Baltimore City

Tobacco Use by Youth: 2000–2013

Maryland Department of Health and Mental Hygiene
Center for Tobacco Prevention and Control

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Any Tobacco Product (All Youth Populations)	20.3% $\pm 3.0\%$ 8,961	18.5% $\pm 2.6\%$ 8,321	15.5% $\pm 2.4\%$ 6,350	<u>20.7%</u> $\pm 2.1\%$ 6,507	21.1% $\pm 2.9\%$ 8,192	16.5% $\pm 2.2\%$ 6,005
Any Tobacco Product (Minority Youth)	19.5% $\pm 3.1\%$ 7,450	17.0% $\pm 2.5\%$ 6,757	15.0% $\pm 2.4\%$ 5,744	20.4% $\pm 4.6\%$ 5,810	20.8% $\pm 2.9\%$ 7,472	16.1% $\pm 2.0\%$ 5,132
Cigarette	10.8% $\pm 2.1\%$ 4,423	9.5% $\pm 1.6\%$ 3,954	7.7% $\pm 1.4\%$ 2,985	10.2% $\pm 1.3\%$ 3,072	9.5% $\pm 1.7\%$ 3,641	7.7% $\pm 1.4\%$ 2,575
Cigar	8.8% $\pm 1.8\%$ 3,638	7.7% $\pm 1.6\%$ 3,164	6.3% $\pm 1.1\%$ 2,357	<u>16.2%</u> $\pm 2.1\%$ 4,796	13.5% $\pm 2.1\%$ 5,067	12.4% $\pm 1.8\%$ 4,302
Pipe (tobacco)	4.2% $\pm 1.2\%$ 1,724	6.6% $\pm 1.4\%$ 2,707	5.4% $\pm 1.6\%$ 2,118	<u>9.8%</u> $\pm 1.8\%$ 2,767	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Bidi	6.5% $\pm 2.0\%$ 2,600	8.1% $\pm 1.9\%$ 3,350	6.1% $\pm 1.5\%$ 2,372	7.2% $\pm 1.3\%$ 2,017	BIDI SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	BIDI SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Kretek	4.5% $\pm 1.4\%$ 1,830	6.3% $\pm 1.7\%$ 2,611	6.5% $\pm 1.6\%$ 2,494	6.1% $\pm 1.3\%$ 1,711	KRETEK SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	KRETEK SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Smokeless Tobacco	3.6% $\pm 1.1\%$ 1,490	4.3% $\pm 1.2\%$ 1,768	3.1% $\pm 0.9\%$ 1,187	4.2% $\pm 0.9\%$ 1,141	2.9% $\pm 0.9\%$ 1,011	*<u>7.3%</u> $\pm 1.8\%$ 2,583

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	19.2% $\pm 2.9\%$ 8,425	14.5% $\pm 1.7\%$ 6,532	14.2% $\pm 1.8\%$ 5,786	16.9% $\pm 1.4\%$ 5,294	<u>21.7%</u> $\pm 2.3\%$ 8,423	*<u>14.4%</u> $\pm 1.6\%$ 4,807
Minority Youth	17.4% $\pm 2.1\%$ 6,597	13.5% $\pm 1.8\%$ 5,369	13.9% $\pm 1.8\%$ 5,312	16.5% $\pm 1.4\%$ 4,686	<u>21.6%</u> $\pm 2.3\%$ 7,758	<u>14.4%</u> $\pm 1.6\%$ 4,220

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	45.8% $\pm 7.3\%$ 1,978	44.8% $\pm 8.7\%$ 1,185	47.8% $\pm 9.4\%$ 1,027	43.7% $\pm 5.2\%$ 1,282	43.5% $\pm 6.2\%$ 1,088	49.2% $\pm 4.1\%$ 3,858
Minority Youth	48.8% $\pm 9.8\%$ 1,538	48.5% $\pm 10.9\%$ 985	51.1% $\pm 10.5\%$ 940	46.7% $\pm 5.5\%$ 1,219	45.1% $\pm 6.7\%$ 987	49.7% $\pm 4.1\%$ 3,476

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals ($\pm\%$) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2013 preceded by an asterisk (*).


CURRENT CIGARETTE SMOKING — School Type, Gender, and Minority Populations

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Middle School (MS)	9.0% \pm 3.0% 1,819	7.7% \pm 1.9% 1,524	6.6% \pm 2.5% 1,153	7.0% \pm 1.7% 972	5.8% \pm 1.0% 950	4.9% \pm 1.4% 748
MS Females	8.9% \pm 3.9% 874	6.4% \pm 1.7% 628	5.5% \pm 3.1% 468	5.5% \pm 2.2% 383	5.5% \pm 1.2% 454	3.9% \pm 1.8% 303
MS Males	9.2% \pm 4.7% 945	9.1% \pm 3.6% 896	7.3% \pm 3.5% 657	8.5% \pm 2.1% 585	6.0% \pm 1.7% 476	5.9% \pm 2.2% 445
MS Minorities	7.7% \pm 2.9% 1,304	7.2% \pm 2.2% 1,273	5.5% \pm 2.3% 894	6.7% \pm 1.5% 839	5.3% \pm 0.9% 794	4.5% \pm 1.6% 592
High School (HS)	12.5% \pm 2.9% 2,605	11.1% \pm 2.0% 2,430	8.6% \pm 1.3% 1,832	12.9% \pm 2.0% 2,101	12.3% \pm 2.4% 2,692	10.1% \pm 1.8% 1,827
HS Females	12.0% \pm 4.1% 1,278	9.7% \pm 2.6% 1,152	6.9% \pm 1.5% 791	10.0% \pm 1.9% 840	7.6% \pm 1.9% 861	6.8% \pm 1.8% 645
HS Males	13.1% \pm 4.5% 1,316	12.5% \pm 3.0% 1,235	10.4% \pm 2.2% 1,032	15.7% \pm 2.7% 1,243	17.1% \pm 2.9% 1,809	12.9% \pm 3.1% 1,107
HS Minorities	11.2% \pm 2.8% 2,063	8.6% \pm 2.0% 1,647	8.1% \pm 1.4% 1,634	11.7% \pm 2.1% 1,730	11.6% \pm 2.2% 2,364	9.2% \pm 1.8% 1,479

Baltimore City—Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Baltimore County

Tobacco Use by Youth: 2000–2013

Maryland Department of Health and Mental Hygiene
Center for Tobacco Prevention and Control

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Any Tobacco Product (All Youth Populations)	21.6% ^{± 3.9%} 11,445	18.4% ^{± 4.0%} 10,268	16.1% ^{± 1.9%} 9,059	18.8% ^{± 2.0%} 9,391	18.2% ^{± 3.4%} 9,519	*12.8% ^{± 2.4%} 6,319
Any Tobacco Product (Minority Youth)	18.6% ^{± 3.5%} 3,388	15.2% ^{± 3.7%} 3,575	13.2% ^{± 1.9%} 3,819	17.2% ^{± 1.8%} 4,274	17.7% ^{± 3.5%} 4,919	*10.8% ^{± 2.0%} 2,796
Cigarette	15.8% ^{± 3.6%} 8,046	12.8% ^{± 3.2%} 6,815	10.4% ^{± 1.5%} 5,628	12.1% ^{± 1.7%} 5,866	10.8% ^{± 2.5%} 5,548	*8.1% ^{± 1.8%} 3,843
Cigar	9.0% ^{± 2.4%} 4,650	8.1% ^{± 1.9%} 4,345	6.6% ^{± 1.0%} 3,565	12.7% ^{± 1.5%} 6,172	11.1% ^{± 2.5%} 5,730	8.2% ^{± 1.6%} 3,981
Pipe (tobacco)	3.4% ^{± 1.1%} 1,799	4.6% ^{± 1.2%} 2,435	4.6% ^{± 0.8%} 2,547	7.6% ^{± 1.5%} 3,634	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Bidi	5.3% ^{± 1.2%} 2,731	4.9% ^{± 1.3%} 2,631	4.8% ^{± 0.9%} 2,633	4.5% ^{± 0.9%} 2,150	BIDI SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	BIDI SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Kretek	4.0% ^{± 0.9%} 2,103	3.9% ^{± 1.2%} 2,086	5.2% ^{± 0.9%} 2,866	4.7% ^{± 1.1%} 2,233	KRETEK SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	KRETEK SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Smokeless Tobacco	2.6% ^{± 0.7%} 1,358	2.9% ^{± 0.8%} 1,565	3.4% ^{± 0.6%} 1,860	3.9% ^{± 1.0%} 1,808	3.0% ^{± 1.0%} 1,477	3.9% ^{± 0.8%} 1,892

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	18.5% ^{± 3.4%} 9,829	16.9% ^{± 2.7%} 9,431	15.4% ^{± 1.6%} 8,663	15.8% ^{± 1.9%} 7,861	19.5% ^{± 3.2%} 10,181	13.1% ^{± 2.4%} 6,204
Minority Youth	15.2% ^{± 2.9%} 2,775	12.2% ^{± 2.8%} 2,890	11.1% ^{± 1.6%} 3,207	12.3% ^{± 1.9%} 3,038	19.9% ^{± 3.4%} 5,541	11.1% ^{± 2.0%} 2,742

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	34.2% ^{± 6.7%} 2,014	32.5% ^{± 5.6%} 1,501	33.5% ^{± 4.6%} 1,243	32.6% ^{± 6.1%} 1,534	42.4% ^{± 7.3%} 1,550	44.6% ^{± 4.1%} 4,044
Minority Youth	54.1% ^{± 9.6%} 667	45.4% ^{± 10.9%} 623	47.1% ^{± 8.9%} 584	42.0% ^{± 7.7%} 766	50.1% ^{± 10.7%} 809	51.5% ^{± 5.3%} 2,161

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals ([±]%) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2013 preceded by an asterisk (*).


CURRENT CIGARETTE SMOKING — School Type, Gender, and Minority Populations

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Middle School (MS)	6.5% $\pm 3.3\%$ 1,536	4.6% $\pm 1.9\%$ 1,121	3.0% $\pm 1.2\%$ 696	4.6% $\pm 2.0\%$ 996	3.0% $\pm 1.0\%$ 670	4.0% $\pm 1.0\%$ 865
MS Females	8.2% $\pm 3.9\%$ 955	4.9% $\pm 2.3\%$ 586	1.8% $\pm 1.1\%$ 198	4.1% $\pm 2.5\%$ 444	2.3% $\pm 1.2\%$ 258	*2.6% $\pm 1.2\%$ 277
MS Males	4.9% $\pm 3.3\%$ 574	4.4% $\pm 1.9\%$ 536	4.2% $\pm 1.8\%$ 497	5.0% $\pm 2.0\%$ 552	3.8% $\pm 2.2\%$ 412	5.2% $\pm 1.2\%$ 565
MS Minorities	3.8% $\pm 1.8\%$ 321	4.7% $\pm 2.2\%$ 528	2.6% $\pm 1.5\%$ 316	3.4% $\pm 1.2\%$ 386	3.5% $\pm 1.3\%$ 423	3.5% $\pm 1.4\%$ 393
High School (HS)	23.7% $\pm 3.0\%$ 6,510	19.7% $\pm 3.4\%$ 5,694	15.8% $\pm 1.6\%$ 4,933	18.3% $\pm 2.6\%$ 4,870	16.7% $\pm 2.5\%$ 4,878	*11.5% $\pm 2.5\%$ 2,978
HS Females	25.6% $\pm 4.5\%$ 3,501	20.3% $\pm 3.7\%$ 2,973	15.1% $\pm 2.0\%$ 2,399	16.1% $\pm 3.0\%$ 2,151	15.1% $\pm 2.8\%$ 2,262	*10.9% $\pm 3.3\%$ 1,432
HS Males	21.5% $\pm 3.1\%$ 2,932	18.9% $\pm 4.2\%$ 2,655	16.5% $\pm 2.1\%$ 2,526	20.4% $\pm 2.6\%$ 2,711	18.4% $\pm 3.0\%$ 2,609	*11.6% $\pm 2.2\%$ 1,461
HS Minorities	16.2% $\pm 2.7\%$ 1,445	11.9% $\pm 4.1\%$ 1,315	9.8% $\pm 1.6\%$ 1,496	12.8% $\pm 1.6\%$ 1,629	11.5% $\pm 1.8\%$ 1,733	*6.7% $\pm 1.6\%$ 888

Baltimore County—Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Calvert County

Tobacco Use by Youth: 2000–2013

Maryland Department of Health and Mental Hygiene
Center for Tobacco Prevention and Control

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Any Tobacco Product (All Youth Populations)	24.2% \pm 2.7% 1,932	22.5% \pm 3.8% 1,979	17.2% \pm 2.4% 1,677	14.7% \pm 1.7% 1,382	17.4% \pm 2.2% 1,615	*14.7% \pm 1.8% 1,264
Any Tobacco Product (Minority Youth)	24.8% \pm 4.4% 433	27.4% \pm 6.8% 527	21.0% \pm 3.4% 519	17.7% \pm 3.4% 350	18.3% \pm 3.0% 415	17.4% \pm 3.1% 356
Cigarette	19.6% \pm 2.7% 1,502	17.9% \pm 3.5% 1,505	13.3% \pm 2.0% 1,270	10.6% \pm 1.4% 985	11.6% \pm 1.7% 1,060	*10.7% \pm 1.6% 880
Cigar	10.0% \pm 1.8% 786	10.1% \pm 2.3% 856	6.4% \pm 1.2% 612	7.2% \pm 1.1% 671	7.8% \pm 1.3% 716	8.2% \pm 1.4% 697
Pipe (tobacco)	3.6% \pm 0.9% 286	6.5% \pm 1.9% 551	3.8% \pm 0.9% 367	3.7% \pm 0.8% 344	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Bidi	3.9% \pm 0.9% 306	5.5% \pm 1.5% 467	3.8% \pm 1.0% 363	2.5% \pm 0.7% 232	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010	BIDI SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Kretek	3.5% \pm 0.8% 274	5.4% \pm 2.1% 455	4.1% \pm 1.0% 391	2.3% \pm 0.6% 211	KRETEK SPECIFIC DATA NOT AVAILA- BLE FOR 2010	KRETEK SPECIFIC DATA NOT AVAILA- BLE FOR 2013
Smokeless Tobacco	2.8% \pm 0.7% 220	4.0% \pm 1.4% 340	3.4% \pm 0.8% 320	4.1% \pm 0.8% 374	3.7% \pm 0.9% 338	*5.6% \pm 1.2% 480

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	23.3% \pm 2.1% 1,857	19.5% \pm 2.9% 1,717	17.5% \pm 2.1% 1,707	14.4% \pm 1.5% 1,353	18.4% \pm 2.1% 1,708	*14.5% \pm 1.8% 1,203
Minority Youth	20.5% \pm 3.4% 358	19.4% \pm 4.4% 373	16.2% \pm 2.6% 398	12.9% \pm 2.6% 256	20.9% \pm 3.2% 472	17.7% \pm 2.9% 338

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	31.2% \pm 4.8% 367	26.7% \pm 5.6% 275	36.1% \pm 5.6% 311	38.5% \pm 5.9% 299	39.2% \pm 5.8% 279	*40.1% \pm 3.9% 670
Minority Youth	27.6% \pm 9.4% 61	29.7% \pm 13.3% 62	37.9% \pm 11.8% 80	36.7% \pm 10.7% 72	39.0% \pm 11.2% 67	*46.2% \pm 7.1% 213

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (\pm %) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2013 preceded by an asterisk (*).


CURRENT CIGARETTE SMOKING — School Type, Gender, and Minority Populations

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Middle School (MS)	9.6% \pm 3.4% 337	6.1% \pm 2.4% 238	3.8% \pm 1.5% 152	2.9% \pm 1.2% 114	3.0% \pm 1.2% 115	*3.1% \pm 1.2% 112
MS Females	12.2% \pm 5.0% 199	5.3% \pm 2.9% 99	3.8% \pm 1.9% 75	2.7% \pm 1.5% 52	3.3% \pm 1.9% 64	*4.1% \pm 2.2% 73
MS Males	7.2% \pm 3.3% 137	6.8% \pm 2.8% 139	3.7% \pm 2.1% 77	3.1% \pm 1.8% 62	2.7% \pm 1.5% 51	*1.9% \pm 1.0% 36
MS Minorities	11.8% \pm 6.6% 97	7.4% \pm 3.9% 69	7.3% \pm 3.4% 74	5.3% \pm 3.6% 45	4.2% \pm 2.2% 40	4.9% \pm 2.4% 41
High School (HS)	28.3% \pm 3.6% 1,164	28.1% \pm 5.8% 1,267	20.3% \pm 2.1% 1,119	16.2% \pm 2.2% 871	17.7% \pm 2.3% 945	*16.8% \pm 2.5% 768
HS Females	29.2% \pm 4.3% 604	26.1% \pm 6.5% 610	20.8% \pm 2.6% 587	15.7% \pm 2.7% 421	15.7% \pm 2.7% 435	*15.4% \pm 3.3% 354
HS Males	27.3% \pm 4.3% 557	30.4% \pm 7.2% 657	19.8% \pm 3.0% 532	16.7% \pm 2.8% 445	19.9% \pm 3.2% 509	*17.9% \pm 2.9% 401
HS Minorities	23.2% \pm 5.2% 196	33.5% \pm 10.5% 290	19.8% \pm 3.8% 268	17.4% \pm 3.1% 193	19.6% \pm 3.6% 247	17.3% \pm 4.1% 182

Calvert County—Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Caroline County

Tobacco Use by Youth: 2000–2013

Maryland Department of Health and Mental Hygiene
Center for Tobacco Prevention and Control

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Any Tobacco Product (All Youth Populations)	31.3% ± 3.9% 870	25.7% ± 2.4% 747	22.7% ± 2.7% 663	21.6% ± 2.5% 599	20.9% ± 2.7% 548	*16.7% ± 2.4% 432
Any Tobacco Product (Minority Youth)	29.6% ± 5.7% 205	24.5% ± 4.6% 184	24.3% ± 4.0% 196	21.5% ± 3.5% 160	24.1% ± 4.1% 206	*17.0% ± 3.3% 124
Cigarette	25.8% ± 3.6% 691	19.8% ± 2.0% 554	16.9% ± 2.4% 842	17.0% ± 2.3% 460	14.3% ± 2.3% 366	*13.1% ± 2.2% 324
Cigar	13.2% ± 2.3% 362	10.0% ± 1.4% 285	8.6% ± 1.5% 246	12.0% ± 1.8% 328	10.5% ± 1.9% 273	*8.7% ± 1.8% 222
Pipe (tobacco)	5.0% ± 1.5% 138	5.0% ± 1.2% 143	4.8% ± 1.1% 138	6.2% ± 1.2% 167	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Bidi	5.5% ± 1.3% 149	4.8% ± 1.3% 138	3.8% ± 1.0% 110	3.8% ± 1.0% 104	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010	BIDI SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Kretek	5.9% ± 1.5% 162	4.1% ± 1.0% 117	4.1% ± 1.1% 119	3.4% ± 1.0% 93	KRETEK SPECIFIC DATA NOT AVAILA- BLE FOR 2010	KRETEK SPECIFIC DATA NOT AVAILA- BLE FOR 2013
Smokeless Tobacco	6.3% ± 1.8% 172	5.0% ± 1.2% 140	5.6% ± 1.1% 161	6.0% ± 1.3% 159	6.6% ± 1.4% 167	8.1% ± 1.8% 207

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	25.2% ± 3.0% 700	22.7% ± 2.2% 661	23.5% ± 2.5% 689	19.9% ± 2.1% 551	19.8% ± 2.6% 517	*15.2% ± 2.2% 376
Minority Youth	20.1% ± 5.1% 139	22.8% ± 3.4% 172	20.2% ± 3.3% 163	15.7% ± 3.0% 116	23.2% ± 4.1% 198	14.5% ± 3.3% 98

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	26.0% ± 6.4% 124	28.2% ± 5.1% 96	35.1% ± 6.5% 117	31.7% ± 5.5% 109	41.4% ± 8.3% 96	36.4% ± 5.3% 198
Minority Youth	22.5% ± 11.2% 24	41.4% ± 13.8% 26	36.2% ± 12.0% 29	45.9% ± 12.3% 38	36.9% ± 13.6% 23	41.4% ± 9.0% 61

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2013 preceded by an asterisk (*).


CURRENT CIGARETTE SMOKING — School Type, Gender, and Minority Populations


Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Middle School (MS)	13.5% $\pm 3.9\%$ 165	10.6% $\pm 2.5\%$ 141	7.5% $\pm 1.5\%$ 89	6.0% $\pm 2.1\%$ 70	3.4% $\pm 1.4\%$ 38	*4.7% $\pm 2.0\%$ 55
MS Females	11.7% $\pm 6.3\%$ 69	10.5% $\pm 3.2\%$ 68	7.9% $\pm 2.5\%$ 44	4.7% $\pm 2.4\%$ 26	2.9% $\pm 1.4\%$ 16	5.6% $\pm 2.9\%$ 32
MS Males	15.4% $\pm 4.4\%$ 96	10.7% $\pm 2.9\%$ 73	7.2% $\pm 2.0\%$ 45	7.3% $\pm 2.7\%$ 44	4.0% $\pm 2.1\%$ 22	*3.9% $\pm 2.5\%$ 22
MS Minorities	13.9% $\pm 6.0\%$ 46	11.8% $\pm 4.4\%$ 43	9.8% $\pm 3.3\%$ 31	6.1% $\pm 3.0\%$ 20	3.2% $\pm 1.9\%$ 13	*4.4% $\pm 2.9\%$ 15
High School (HS)	36.0% $\pm 5.0\%$ 527	28.3% $\pm 3.5\%$ 413	23.7% $\pm 2.9\%$ 393	25.1% $\pm 3.5\%$ 390	22.6% $\pm 3.4\%$ 328	*20.6% $\pm 3.5\%$ 269
HS Females	35.5% $\pm 5.9\%$ 249	28.9% $\pm 4.2\%$ 207	23.1% $\pm 3.7\%$ 191	22.5% $\pm 4.2\%$ 172	19.1% $\pm 3.5\%$ 149	*15.3% $\pm 3.7\%$ 100
HS Males	36.3% $\pm 6.0\%$ 274	27.1% $\pm 5.0\%$ 198	24.0% $\pm 3.6\%$ 199	27.6% $\pm 4.8\%$ 217	26.7% $\pm 5.1\%$ 178	25.6% $\pm 4.9\%$ 165
HS Minorities	30.0% $\pm 7.8\%$ 96	22.5% $\pm 6.7\%$ 76	21.4% $\pm 4.5\%$ 93	19.9% $\pm 4.5\%$ 76	24.6% $\pm 5.5\%$ 105	*15.4% $\pm 4.7\%$ 54

Caroline County—Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Carroll County

Tobacco Use by Youth: 2000–2013

Maryland Department of Health and Mental Hygiene
Center for Tobacco Prevention and Control

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Any Tobacco Product (All Youth Populations)	21.1% ± 3.4% 3,014	18.9% ± 3.4% 2,842	17.0% ± 2.2% 2,764	15.3% ± 1.6% 2,372	15.0% ± 1.6% 2,263	*12.1% ± 1.4% 1,706
Any Tobacco Product (Minority Youth)	34.8% ± 7.7% 463	26.9% ± 6.1% 466	23.0% ± 4.2% 436	23.9% ± 4.1% 327	18.0% ± 3.0% 319	*17.4% ± 3.7% 257
Cigarette	16.4% ± 3.0% 2,263	13.9% ± 2.9% 2,024	12.1% ± 1.7% 1,932	11.2% ± 1.2% 1,709	9.5% ± 1.2% 1,419	*8.4% ± 1.2% 1,143
Cigar	7.6% ± 1.9% 1,076	7.9% ± 1.7% 1,169	7.9% ± 1.2% 1,263	7.8% ± 1.1% 1,196	7.2% ± 1.0% 1,084	6.9% ± 1.0% 964
Pipe (tobacco)	3.1% ± 0.8% 438	4.7% ± 1.3% 688	3.4% ± 0.6% 551	3.4% ± 0.7% 525	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Bidi	4.8% ± 1.1% 674	4.9% ± 1.4% 728	2.9% ± 0.6% 477	2.5% ± 0.6% 384	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010	BIDI SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Kretek	3.3% ± 0.8% 469	4.0% ± 1.2% 595	3.3% ± 0.6% 527	1.9% ± 0.4% 292	KRETEK SPECIFIC DATA NOT AVAILA- BLE FOR 2010	KRETEK SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Smokeless Tobacco	3.2% ± 0.9% 448	4.0% ± 1.1% 587	3.8% ± 0.7% 614	4.7% ± 0.8% 707	4.3% ± 0.7% 638	*5.5% ± 1.0% 780

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	17.7% ± 2.3% 2,522	16.2% ± 2.6% 2,433	18.0% ± 2.3% 2,940	15.7% ± 1.6% 2,425	16.2% ± 1.7% 2,451	*11.6% ± 1.4% 1,591
Minority Youth	22.3% ± 5.3% 293	15.4% ± 5.4% 267	15.7% ± 3.5% 298	14.2% ± 3.6% 194	17.9% ± 3.1% 318	17.0% ± 3.7% 233

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	34.2% ± 6.1% 558	27.8% ± 8.2% 352	36.0% ± 5.1% 385	30.6% ± 5.0% 410	43.0% ± 6.3% 372	37.5% ± 4.1% 830
Minority Youth	26.0% ± 15.2% 49	Not Available	41.3% ± 17.0% 298	22.3% ± 11.4% 33	45.9% ± 15.4% 46	44.5% ± 10.0% 138

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2013 preceded by an asterisk (*).


CURRENT CIGARETTE SMOKING — School Type, Gender, and Minority Populations


Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Middle School (MS)	4.9% \pm 2.8% 302	3.1% \pm 1.6% 208	3.5% \pm 1.3% 233	1.6% \pm 0.8% 103	1.8% \pm 1.0% 109	2.4% \pm 1.2% 143
MS Females	4.5% \pm 3.2% 134	3.3% \pm 2.4% 108	3.4% \pm 1.5% 111	1.8% \pm 1.8% 54	1.6% \pm 1.1% 50	1.5% \pm 1.2% 43
MS Males	5.2% \pm 2.9% 168	2.5% \pm 1.6% 86	3.6% \pm 2.0% 122	1.5% \pm 0.8% 49	2.0% \pm 1.3% 59	3.3% \pm 1.8% 100
MS Minorities	11.2% \pm 9.9% 71	8.0% \pm 5.5% 65	4.2% \pm 4.1% 36	6.0% \pm 3.9% 38	3.2% \pm 2.3% 27	5.6% \pm 3.9% 36
High School (HS)	25.9% \pm 3.7% 1,960	23.1% \pm 4.2% 1,816	18.1% \pm 1.8% 1,699	17.8% \pm 1.9% 1,606	14.8% \pm 1.7% 1,310	*13.0% \pm 1.8% 1,000
HS Females	26.2% \pm 4.8% 963	23.4% \pm 5.2% 916	17.7% \pm 2.3% 808	15.5% \pm 2.2% 693	13.2% \pm 2.0% 593	*10.9% \pm 2.0% 420
HS Males	25.6% \pm 3.9% 975	22.3% \pm 4.6% 860	18.5% \pm 2.2% 887	20.0% \pm 2.6% 902	16.5% \pm 2.4% 717	*14.5% \pm 2.5% 551
HS Minorities	42.8% \pm 9.0% 275	28.6% \pm 7.4% 223	20.9% \pm 4.7% 191	25.4% \pm 4.5% 173	17.6% \pm 3.7% 154	*18.4% \pm 4.5% 136

Carroll County—Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Cecil County

Tobacco Use by Youth: 2000–2013

Maryland Department of Health and Mental Hygiene
Center for Tobacco Prevention and Control

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Any Tobacco Product (All Youth Populations)	26.9% \pm 3.7% 2,016	22.9% \pm 2.9% 1,840	21.8% \pm 2.7% 1,914	20.4% \pm 1.8% 1,721	20.2% \pm 2.0% 1,658	<u>*15.9%</u> \pm 2.0% 1,234
Any Tobacco Product (Minority Youth)	29.2% \pm 6.5% 274	25.4% \pm 6.4% 287	26.1% \pm 4.1% 462	22.7% \pm 3.5% 321	20.4% \pm 3.2% 325	<u>*14.9%</u> \pm 2.7% 208
Cigarette	22.3% \pm 3.4% 1,613	17.6% \pm 2.7% 1,347	16.0% \pm 2.4% 1,360	15.0% \pm 1.5% 1,245	13.8% \pm 1.5% 1,114	<u>*11.6%</u> \pm 1.6% 864
Cigar	9.2% \pm 1.8% 679	7.3% \pm 1.5% 575	9.7% \pm 1.4% 829	12.3% \pm 1.5% 1,025	10.2% \pm 1.4% 831	8.9% \pm 1.2% 683
Pipe (tobacco)	3.3% \pm 0.9% 248	4.2% \pm 1.3% 327	5.1% \pm 0.9% 442	5.9% \pm 1.2% 490	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Bidi	6.2% \pm 1.4% 454	6.0% \pm 1.2% 474	4.0% \pm 1.0% 426	3.8% \pm 0.9% 309	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010	BIDI SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Kretek	2.8% \pm 0.7% 208	4.2% \pm 1.1% 329	5.0% \pm 1.0% 434	3.4% \pm 0.9% 283	KRETEK SPECIFIC DATA NOT AVAILA- BLE FOR 2010	KRETEK SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Smokeless Tobacco	3.3% \pm 0.9% 242	3.8% \pm 0.9% 292	5.6% \pm 1.0% 484	5.7% \pm 0.9% 462	5.3% \pm 1.1% 421	<u>*6.0%</u> \pm 1.0% 462

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	23.3% \pm 2.9% 1,748	18.9% \pm 2.6% 1,518	20.8% \pm 2.5% 1,831	19.3% \pm 1.6% 1,630	21.1% \pm 2.1% 1,737	<u>*15.9%</u> \pm 1.8% 1,191
Minority Youth	26.4% \pm 5.2% 248	21.3% \pm 5.1% 238	17.7% \pm 3.0% 313	17.9% \pm 3.2% 251	19.6% \pm 3.1% 312	18.0% \pm 3.3% 242

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	26.2% \pm 5.0% 297	32.5% \pm 5.0% 333	34.3% \pm 5.5% 335	35.1% \pm 4.5% 344	42.2% \pm 5.5% 317	<u>*38.4%</u> \pm 3.7% 655
Minority Youth	27.3% \pm 13.8% 39	25.8% \pm 12.6% 37	39.1% \pm 12.7% 69	41.2% \pm 10.2% 73	49.8% \pm 10.9% 68	43.3% \pm 8.0% 151

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (\pm %) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2013 preceded by an asterisk (*).


CURRENT CIGARETTE SMOKING — School Type, Gender, and Minority Populations


Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Middle School (MS)	11.3% $\pm 2.9\%$ 389	9.8% $\pm 2.7\%$ 359	6.1% $\pm 2.7\%$ 235	6.2% $\pm 1.9\%$ 222	5.0% $\pm 1.4\%$ 176	*5.3% $\pm 1.6\%$ 180
MS Females	11.6% $\pm 4.2\%$ 198	11.7% $\pm 3.6\%$ 205	7.3% $\pm 4.4\%$ 136	5.9% $\pm 2.2\%$ 103	4.9% $\pm 1.9\%$ 89	6.4% $\pm 2.5\%$ 106
MS Males	11.1% $\pm 3.2\%$ 191	8.0% $\pm 2.9\%$ 151	5.0% $\pm 2.5\%$ 99	6.4% $\pm 2.5\%$ 119	5.1% $\pm 2.0\%$ 87	*4.2% $\pm 1.6\%$ 72
MS Minorities	13.6% $\pm 6.4\%$ 63	6.5% $\pm 3.8\%$ 38	8.0% $\pm 4.5\%$ 66	7.2% $\pm 3.3\%$ 50	5.2% $\pm 3.1\%$ 38	*4.2% $\pm 2.7\%$ 24
High School (HS)	32.3% $\pm 4.1\%$ 1,224	24.7% $\pm 3.7\%$ 988	24.0% $\pm 2.1\%$ 1,125	21.8% $\pm 2.4\%$ 1,023	20.5% $\pm 2.1\%$ 938	*16.9% $\pm 2.2\%$ 683
HS Females	35.3% $\pm 4.5\%$ 656	28.3% $\pm 4.6\%$ 581	23.4% $\pm 2.7\%$ 550	21.0% $\pm 3.0\%$ 495	20.3% $\pm 2.8\%$ 479	*15.7% $\pm 2.9\%$ 314
HS Males	29.4% $\pm 5.8\%$ 562	21.0% $\pm 4.6\%$ 403	24.5% $\pm 2.9\%$ 575	22.5% $\pm 3.0\%$ 522	20.7% $\pm 2.9\%$ 460	*17.6% $\pm 2.7\%$ 360
HS Minorities	36.6% $\pm 8.9\%$ 156	33.0% $\pm 9.9\%$ 151	25.7% $\pm 4.6\%$ 216	24.7% $\pm 4.6\%$ 168	19.5% $\pm 3.8\%$ 164	*14.8% $\pm 3.5\%$ 107

Cecil County—Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Charles County

Tobacco Use by Youth: 2000–2013

Maryland Department of Health and Mental Hygiene
Center for Tobacco Prevention and Control

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Any Tobacco Product (All Youth Populations)	24.6% ± 3.2% 2,957	20.4% ± 2.6% 2,572	14.4% ± 1.9% 2,156	16.9% ± 1.6% 2,531	16.6% ± 1.9% 2,491	*12.6% ± 1.6% 1,738
Any Tobacco Product (Minority Youth)	21.9% ± 3.3% 913	17.0% ± 3.0% 955	13.8% ± 2.1% 1,264	16.6% ± 2.1% 1,491	15.9% ± 1.8% 1,542	*11.5% ± 1.6% 1,011
Cigarette	19.7% ± 3.0% 2,276	14.6% ± 2.2% 1,771	9.9% ± 1.7% 1,444	11.2% ± 1.3% 1,633	9.5% ± 1.3% 1,400	*7.9% ± 1.2% 1,041
Cigar	10.4% ± 2.0% 1,215	7.6% ± 1.3% 900	6.3% ± 1.0% 921	11.7% ± 1.4% 1,716	9.1% ± 1.2% 1,354	8.4% ± 1.2% 1,132
Pipe (tobacco)	4.8% ± 1.1% 569	5.0% ± 1.5% 589	4.0% ± 0.8% 593	5.7% ± 1.1% 831	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Bidi	6.0% ± 1.3% 694	5.2% ± 1.7% 621	4.1% ± 0.8% 608	3.5% ± 0.7% 505	BIDI SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	BIDI SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Kretek	4.6% ± 1.1% 544	4.6% ± 1.6% 548	4.6% ± 0.9% 680	3.4% ± 0.7% 494	KRETEK SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	KRETEK SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Smokeless Tobacco	3.9% ± 0.9% 242	4.1% ± 1.6% 292	3.2% ± 0.7% 484	3.4% ± 0.7% 462	3.3% ± 0.8% 471	4.4% ± 0.8% 597

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	21.6% ± 2.4% 2,597	18.4% ± 1.8% 2,331	14.6% ± 1.9% 2,187	16.0% ± 1.4% 2,398	18.9% ± 1.7% 2,834	*13.7% ± 1.4% 1,813
Minority Youth	19.5% ± 3.0% 819	14.9% ± 2.4% 838	12.8% ± 2.0% 1,175	14.8% ± 1.7% 1,324	18.4% ± 1.8% 1,782	*13.0% ± 1.6% 1,090

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	33.1% ± 5.7% 534	31.8% ± 6.6% 404	38.4% ± 5.8% 389	38.0% ± 5.3% 553	52.3% ± 5.8% 559	*44.5% ± 4.3% 1,177
Minority Youth	43.9% ± 9.7% 194	43.6% ± 14.3% 143	44.4% ± 8.4% 246	40.4% ± 7.7% 325	58.7% ± 7.5% 344	48.3% ± 5.3% 802

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (± %) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2013 preceded by an asterisk (*).


CURRENT CIGARETTE SMOKING — School Type, Gender, and Minority Populations


Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Middle School (MS)	9.3% \pm 2.8% 478	5.5% \pm 1.7% 303	1.6% \pm 0.8% 98	3.1% \pm 1.0% 183	3.7% \pm 1.3% 224	*3.1% \pm 1.2% 171
MS Females	8.9% \pm 4.2% 219	6.5% \pm 2.4% 180	1.0% \pm 1.2% 30	2.7% \pm 1.2% 78	3.3% \pm 1.6% 103	3.5% \pm 1.8% 92
MS Males	9.7% \pm 3.3% 259	4.6% \pm 2.0% 123	2.2% \pm 1.0% 67	3.2% \pm 1.4% 99	4.1% \pm 1.9% 120	*2.8% \pm 1.4% 78
MS Minorities	8.7% \pm 3.6% 166	5.1% \pm 2.2% 127	1.6% \pm 1.0% 63	3.2% \pm 1.1% 118	3.4% \pm 1.3% 133	4.0% \pm 1.8% 144
High School (HS)	28.0% \pm 4.0% 1,797	22.0% \pm 3.3% 1,469	16.0% \pm 1.9% 1,347	16.8% \pm 2.1% 1,450	13.6% \pm 1.8% 1,176	*11.5% \pm 1.6% 870
HS Females	26.9% \pm 4.4% 860	21.2% \pm 3.5% 729	15.1% \pm 2.0% 656	14.2% \pm 2.3% 615	12.3% \pm 2.1% 552	*9.5% \pm 2.0% 357
HS Males	28.9% \pm 5.0% 923	22.7% \pm 5.3% 728	16.9% \pm 2.4% 691	19.4% \pm 2.8% 834	14.8% \pm 2.4% 618	*13.1% \pm 2.4% 493
HS Minorities	22.5% \pm 4.1% 457	14.3% \pm 3.9% 401	14.0% \pm 2.3% 681	14.6% \pm 2.8% 735	10.3% \pm 1.7% 578	*8.7% \pm 1.6% 415

Charles County—Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Dorchester County

Tobacco Use by Youth: 2000–2013

Maryland Department of Health and Mental Hygiene
Center for Tobacco Prevention and Control

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Any Tobacco Product (All Youth Populations)	25.5% \pm 3.2% 663	22.1% \pm 2.7% 587	19.7% \pm 2.6% 499	18.1% \pm 2.5% 427	21.6% \pm 3.1% 491	*15.4% \pm 2.4% 334
Any Tobacco Product (Minority Youth)	25.3% \pm 4.1% 262	22.3% \pm 3.3% 236	19.6% \pm 3.4% 252	21.2% \pm 3.9% 236	24.6% \pm 3.9% 278	*16.0% \pm 3.3% 158
Cigarette	20.0% \pm 2.9% 495	15.9% \pm 2.3% 405	13.7% \pm 2.2% 335	13.6% \pm 2.1% 312	13.4% \pm 2.2% 298	*9.9% \pm 2.0% 201
Cigar	10.7% \pm 2.0% 272	9.6% \pm 1.7% 247	7.8% \pm 1.7% 188	10.9% \pm 2.0% 248	11.2% \pm 1.9% 252	8.2% \pm 1.8% 175
Pipe (tobacco)	5.9% \pm 1.6% 153	4.8% \pm 1.1% 121	6.7% \pm 1.4% 166	5.8% \pm 1.9% 132	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2013
Bidi	6.6% \pm 1.8% 164	5.5% \pm 1.3% 140	6.2% \pm 1.5% 153	4.1% \pm 1.2% 92	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2013
Kretek	5.1% \pm 1.4% 133	4.6% \pm 1.3% 117	7.0% \pm 1.7% 173	3.5% \pm 1.3% 78	KRETEK SPECIFIC DATA NOT AVAILA- BLE FOR 2010	KRETEK SPECIFIC DATA NOT AVAILA- BLE FOR 2013
Smokeless Tobacco	4.4% \pm 1.3% 112	4.1% \pm 1.2% 103	4.4% \pm 1.1% 106	3.0% \pm 1.0% 65	4.2% \pm 1.2% 91	*8.9% \pm 2.0% 191

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	23.8% \pm 2.5% 622	19.0% \pm 2.4% 504	18.7% \pm 2.4% 473	18.0% \pm 2.0% 423	21.2% \pm 2.9% 483	*14.5% \pm 2.4% 300
Minority Youth	23.4% \pm 3.5% 243	17.3% \pm 2.8% 192	16.0% \pm 2.8% 206	18.8% \pm 2.8% 209	23.6% \pm 3.4% 268	*14.1% \pm 3.1% 130

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	36.0% \pm 6.4% 126	32.5% \pm 6.6% 84	41.1% \pm 6.9% 93	39.8% \pm 7.1% 104	38.1% \pm 7.9% 69	38.7% \pm 6.7% 158
Minority Youth	39.7% \pm 13.9% 40	38.2% \pm 12.4% 32	40.3% \pm 11.9% 34	38.9% \pm 10.3% 44	33.2% \pm 9.8% 32	36.8% \pm 8.6% 65

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (\pm %) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2013 preceded by an asterisk (*).


CURRENT CIGARETTE SMOKING — School Type, Gender, and Minority Populations


Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Middle School (MS)	11.1% $\pm 2.9\%$ 122	8.2% $\pm 2.4\%$ 99	6.3% $\pm 2.1\%$ 63	5.3% $\pm 1.9\%$ 53	5.6% $\pm 2.2\%$ 53	*3.5% $\pm 1.6\%$ 33
MS Females	11.6% $\pm 4.0\%$ 60	8.1% $\pm 2.8\%$ 47	6.6% $\pm 2.4\%$ 33	5.6% $\pm 2.3\%$ 27	4.7% $\pm 2.3\%$ 24	*3.5% $\pm 1.8\%$ 17
MS Males	10.2% $\pm 3.7\%$ 58	8.3% $\pm 3.1\%$ 51	6.1% $\pm 2.7\%$ 31	5.1% $\pm 2.3\%$ 26	6.5% $\pm 3.2\%$ 29	*3.5% $\pm 1.8\%$ 17
MS Minorities	12.0% $\pm 4.6\%$ 54	9.4% $\pm 2.7\%$ 49	6.1% $\pm 2.5\%$ 32	7.4% $\pm 3.5\%$ 33	5.8% $\pm 2.2\%$ 28	5.8% $\pm 2.5\%$ 27
High School (HS)	27.2% $\pm 3.2\%$ 373	22.9% $\pm 3.7\%$ 306	18.7% $\pm 2.5\%$ 272	19.7% $\pm 3.5\%$ 259	19.4% $\pm 3.1\%$ 245	*15.4% $\pm 3.3\%$ 168
HS Females	28.5% $\pm 4.0\%$ 194	24.4% $\pm 4.4\%$ 167	18.4% $\pm 3.3\%$ 142	16.7% $\pm 3.7\%$ 116	14.6% $\pm 3.3\%$ 96	*11.5% $\pm 3.9\%$ 60
HS Males	24.9% $\pm 4.6\%$ 168	21.1% $\pm 4.8\%$ 136	19.1% $\pm 4.1\%$ 130	23.2% $\pm 5.5\%$ 144	24.6% $\pm 4.3\%$ 150	18.7% $\pm 4.7\%$ 105
HS Minorities	23.3% $\pm 5.3\%$ 119	19.4% $\pm 5.2\%$ 102	16.0% $\pm 3.8\%$ 112	19.4% $\pm 5.0\%$ 122	21.9% $\pm 4.3\%$ 134	15.3% $\pm 5.9\%$ 70

Dorchester County—Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Frederick County

Tobacco Use by Youth: 2000–2013

Maryland Department of Health and Mental Hygiene
Center for Tobacco Prevention and Control

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Any Tobacco Product (All Youth Populations)	25.1% \pm 3.2% 4,535	19.6% \pm 3.0% 3,866	17.5% \pm 2.3% 3,795	15.2% \pm 1.5% 3,245	15.0% \pm 3.1% 3,185	*13.5% \pm 2.4% 2,759
Any Tobacco Product (Minority Youth)	26.3% \pm 4.2% 839	26.1% \pm 5.4% 955	22.0% \pm 3.4% 1,216	17.1% \pm 3.8% 811	17.2% \pm 3.4% 1,066	*14.5% \pm 2.7% 882
Cigarette	19.5% \pm 2.8% 3,397	14.0% \pm 2.5% 2,649	11.8% \pm 1.8% 2,470	10.4% \pm 1.2% 2,188	9.2% \pm 2.2% 1,929	*9.0% \pm 1.8% 1,770
Cigar	10.0% \pm 1.5% 1,774	7.6% \pm 1.4% 1,471	7.0% \pm 1.1% 1,473	8.2% \pm 1.0% 1,722	7.9% \pm 2.0% 1,680	7.5% \pm 1.4% 1,520
Pipe (tobacco)	4.3% \pm 1.0% 779	4.9% \pm 1.1% 952	5.4% \pm 0.9% 1,155	4.1% \pm 0.7% 851	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2013
Bidi	5.2% \pm 1.1% 914	5.1% \pm 1.2% 989	5.3% \pm 0.9% 1,141	2.8% \pm 0.6% 580	BIDI SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2013
Kretek	4.5% \pm 1.1% 808	4.2% \pm 1.0% 803	5.4% \pm 0.9% 1,156	2.7% \pm 0.6% 547	KRETEK SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	KRETEK SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Smokeless Tobacco	4.7% \pm 1.1% 836	5.1% \pm 1.1% 980	4.9% \pm 0.9% 1,030	4.6% \pm 0.8% 943	4.1% \pm 1.2% 850	6.3% \pm 1.2% 1,280

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	22.4% \pm 2.4% 4,039	17.5% \pm 2.3% 3,441	16.6% \pm 2.1% 3,581	14.8% \pm 1.4% 3,151	16.9% \pm 3.2% 3,585	*11.8% \pm 2.4% 2,344
Minority Youth	22.9% \pm 3.9% 732	20.6% \pm 4.3% 754	15.4% \pm 2.5% 846	13.4% \pm 2.0% 630	20.3% \pm 3.7% 1,263	*12.4% \pm 2.7% 720

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	33.1% \pm 4.9% 797	36.0% \pm 6.1% 654	38.1% \pm 5.5% 562	31.6% \pm 5.3% 448	44.1% \pm 7.6% 550	41.5% \pm 4.5% 1,489
Minority Youth	32.9% \pm 12.6% 118	34.3% \pm 14.0% 117	41.0% \pm 11.2% 156	34.5% \pm 11.2% 103	57.3% \pm 11.5% 229	43.4% \pm 7.3% 475

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (\pm %) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2013 preceded by an asterisk (*).


CURRENT CIGARETTE SMOKING — School Type, Gender, and Minority Populations


Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Middle School (MS)	8.8% $\pm 2.3\%$ 699	4.2% $\pm 1.4\%$ 366	2.6% $\pm 1.2\%$ 232	2.6% $\pm 0.5\%$ 230	2.7% $\pm 1.0\%$ 237	*3.0% $\pm 0.6\%$ 256
MS Females	8.2% $\pm 3.1\%$ 314	3.4% $\pm 2.0\%$ 148	2.6% $\pm 1.6\%$ 112	2.2% $\pm 1.0\%$ 95	2.5% $\pm 1.4\%$ 114	*1.5% $\pm 1.0\%$ 62
MS Males	9.1% $\pm 2.2\%$ 370	4.8% $\pm 1.9\%$ 208	2.6% $\pm 1.5\%$ 120	2.9% $\pm 0.9\%$ 135	2.8% $\pm 1.1\%$ 123	*4.4% $\pm 1.0\%$ 194
MS Minorities	12.4% $\pm 5.7\%$ 191	7.1% $\pm 3.6\%$ 119	3.9% $\pm 2.7\%$ 88	4.1% $\pm 1.7\%$ 89	4.2% $\pm 1.6\%$ 110	*4.2% $\pm 1.4\%$ 105
High School (HS)	28.5% $\pm 3.4\%$ 2,697	22.4% $\pm 3.1\%$ 2,283	18.8% $\pm 1.9\%$ 2,239	16.2% $\pm 2.1\%$ 1,958	14.0% $\pm 2.1\%$ 1,692	*13.7% $\pm 1.6\%$ 1,514
HS Females	31.3% $\pm 4.2\%$ 1,448	23.6% $\pm 3.8\%$ 1,194	18.0% $\pm 2.2\%$ 1,082	14.9% $\pm 2.4\%$ 901	11.6% $\pm 2.3\%$ 739	*12.2% $\pm 2.4\%$ 670
HS Males	25.8% $\pm 3.7\%$ 1,227	21.0% $\pm 3.8\%$ 1,066	19.4% $\pm 2.4\%$ 1,146	17.3% $\pm 2.7\%$ 1,049	16.6% $\pm 2.8\%$ 949	*14.8% $\pm 2.0\%$ 815
HS Minorities	29.3% $\pm 5.8\%$ 417	27.4% $\pm 7.4\%$ 473	19.3% $\pm 3.0\%$ 553	18.0% $\pm 3.3\%$ 445	13.7% $\pm 2.7\%$ 469	*13.9% $\pm 2.4\%$ 453

Frederick County—Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Garrett County

Tobacco Use by Youth: 2000–2013

Maryland Department of Health and Mental Hygiene
Center for Tobacco Prevention and Control

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Any Tobacco Product (All Youth Populations)	28.1% $\pm 3.8\%$ 669	27.0% $\pm 3.9\%$ 639	23.9% $\pm 3.4\%$ 618	25.0% $\pm 3.1\%$ 596	28.2% $\pm 3.4\%$ 628	25.7% $\pm 3.3\%$ 520
Any Tobacco Product (Minority Youth)	42.1% $\pm 13.1\%$ 59	35.9% $\pm 11.4\%$ 49	32.2% $\pm 7.8\%$ 69	40.6% $\pm 8.2\%$ 66	37.3% $\pm 6.2\%$ 130	45.6% $\pm 7.4\%$ 122
Cigarette	22.1% $\pm 3.1\%$ 513	19.7% $\pm 3.2\%$ 450	17.0% $\pm 2.6\%$ 430	17.9% $\pm 2.7\%$ 416	19.1% $\pm 2.7\%$ 416	*14.8% $\pm 2.5\%$ 282
Cigar	9.0% $\pm 2.0\%$ 212	9.2% $\pm 2.0\%$ 214	8.4% $\pm 1.6\%$ 214	11.5% $\pm 2.0\%$ 273	10.6% $\pm 1.9\%$ 234	10.7% $\pm 1.8\%$ 214
Pipe (tobacco)	4.2% $\pm 1.4\%$ 99	4.8% $\pm 1.2\%$ 110	3.5% $\pm 0.9\%$ 90	6.2% $\pm 1.5\%$ 146	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2013
Bidi	5.3% $\pm 1.5\%$ 124	5.6% $\pm 1.5\%$ 131	3.1% $\pm 0.8\%$ 80	4.3% $\pm 1.3\%$ 102	BIDI SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2013
Kretek	4.2% $\pm 1.4\%$ 100	4.6% $\pm 1.2\%$ 107	3.6% $\pm 0.9\%$ 91	4.2% $\pm 1.1\%$ 99	KRETEK SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	KRETEK SPECIFIC DATA NOT AVAILA- BLE FOR 2013
Smokeless Tobacco	8.2% $\pm 2.0\%$ 194	10.0% $\pm 2.1\%$ 230	9.5% $\pm 1.8\%$ 242	10.7% $\pm 2.0\%$ 247	14.4% $\pm 2.5\%$ 307	*17.2% $\pm 2.9\%$ 345

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	24.7% $\pm 3.2\%$ 585	21.7% $\pm 2.8\%$ 513	23.5% $\pm 3.0\%$ 608	21.9% $\pm 2.7\%$ 522	27.5% $\pm 3.0\%$ 612	20.0% $\pm 2.7\%$ 383
Minority Youth	27.9% $\pm 13.4\%$ 38	20.2% $\pm 8.7\%$ 27	23.3% $\pm 6.0\%$ 50	23.7% $\pm 6.9\%$ 39	36.0% $\pm 6.2\%$ 125	31.1% $\pm 6.9\%$ 71

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	37.6% $\pm 7.7\%$ 147	29.6% $\pm 6.4\%$ 88	33.7% $\pm 6.5\%$ 82	30.0% $\pm 5.7\%$ 91	30.4% $\pm 7.0\%$ 67	29.9% $\pm 4.1\%$ 172
Minority Youth	Not Available	Not Available	Not Available	Not Available	35.3% $\pm 15.5\%$ 15	37.2% $\pm 10.4\%$ 38

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (\pm %) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2013 preceded by an asterisk (*).


CURRENT CIGARETTE SMOKING — School Type, Gender, and Minority Populations


Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Middle School (MS)	9.5% \pm 3.4% 101	10.9% \pm 3.8% 119	8.2% \pm 2.6% 93	6.5% \pm 2.3% 63	7.1% \pm 2.5% 64	6.4% \pm 2.4% 56
MS Females	9.5% \pm 4.4% 49	10.0% \pm 4.4% 51	10.2% \pm 3.7% 55	6.0% \pm 3.1% 28	7.5% \pm 3.5% 34	4.4% \pm 2.5% 19
MS Males	9.4% \pm 5.7% 51	11.6% \pm 4.1% 68	6.5% \pm 2.9% 39	7.1% \pm 2.8% 35	6.8% \pm 3.0% 30	8.4% \pm 3.5% 37
MS Minorities	18.4% \pm 9.8% 14	11.3% \pm 9.6% 8	11.3% \pm 5.9% 14	12.9% \pm 7.7% 11	11.7% \pm 5.5% 20	14.2% \pm 7.3% 17
High School (HS)	32.8% \pm 3.8% 413	27.7% \pm 4.1% 331	24.1% \pm 3.0% 336	25.8% \pm 4.2% 353	27.6% \pm 3.9% 352	*22.0% \pm 3.9% 226
HS Females	31.1% \pm 4.9% 184	26.7% \pm 5.1% 156	22.6% \pm 3.9% 154	25.3% \pm 5.3% 173	27.3% \pm 5.0% 178	*17.6% \pm 4.7% 89
HS Males	34.4% \pm 4.9% 228	28.8% \pm 5.5% 175	25.5% \pm 4.3% 182	26.3% \pm 6.0% 181	27.9% \pm 5.2% 172	26.2% \pm 4.5% 136
HS Minorities	41.6% \pm 15.1% 24	44.6% \pm 16.1% 27	36.4% \pm 10.9% 26	47.3% \pm 10.7% 35	44.3% \pm 9.0% 73	40.7% \pm 10.4% 46

Garrett County—Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Harford County

Tobacco Use by Youth: 2000–2013

Maryland Department of Health and Mental Hygiene
Center for Tobacco Prevention and Control

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Any Tobacco Product (All Youth Populations)	26.0% \pm 3.3% 5,071	<u>19.7%</u> \pm 2.5% 4,056	16.2% \pm 2.1% 3,433	15.7% \pm 1.8% 3,024	17.3% \pm 1.7% 3,525	* <u>13.3%</u> \pm 1.6% 2,524
Any Tobacco Product (Minority Youth)	25.0% \pm 4.5% 1,143	22.5% \pm 3.9% 1,184	15.7% \pm 3.1% 972	18.8% \pm 3.4% 930	18.3% \pm 2.3% 1,103	* <u>14.6%</u> \pm 2.4% 826
Cigarette	21.2% \pm 2.9% 3,933	<u>14.2%</u> \pm 2.1% 2,811	11.0% \pm 1.7% 2,271	11.0% \pm 1.3% 2,079	10.9% \pm 1.3% 2,196	* <u>8.8%</u> \pm 1.2% 1,604
Cigar	11.3% \pm 1.9% 2,153	8.3% \pm 1.4% 1,661	7.3% \pm 1.2% 1,506	9.3% \pm 1.4% 1,753	10.0% \pm 1.3% 2,021	9.0% \pm 1.2% 1,682
Pipe (tobacco)	4.7% \pm 1.1% 894	4.9% \pm 1.1% 997	3.6% \pm 0.7% 752	4.8% \pm 1.1% 889	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2013
Bidi	5.5% \pm 1.2% 1,028	5.1% \pm 1.0% 1,031	3.9% \pm 0.8% 820	3.4% \pm 0.9% 628	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2013
Kretek	4.2% \pm 1.0% 808	4.4% \pm 1.0% 880	4.4% \pm 0.8% 914	3.0% \pm 0.7% 562	KRETEK SPECIFIC DATA NOT AVAILA- BLE FOR 2010	KRETEK SPECIFIC DATA NOT AVAILA- BLE FOR 2013
Smokeless Tobacco	4.8% \pm 1.1% 927	3.7% \pm 0.8% 732	3.3% \pm 0.7% 681	3.5% \pm 0.7% 642	3.3% \pm 0.6% 660	4.7% \pm 0.8% 885

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	23.3% \pm 2.6% 4,534	<u>17.2%</u> \pm 2.1% 3,549	15.8% \pm 2.0% 3,343	15.7% \pm 1.6% 3,020	<u>19.2%</u> \pm 1.7% 3,909	* <u>12.5%</u> \pm 1.4% 2,277
Minority Youth	20.4% \pm 3.7% 930	17.5% \pm 3.0% 918	12.6% \pm 2.1% 781	14.2% \pm 2.1% 697	<u>20.0%</u> \pm 2.3% 1,206	* <u>12.6%</u> \pm 2.0% 670

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	30.0% \pm 4.9% 839	31.3% \pm 5.4% 521	32.1% \pm 5.1% 435	34.3% \pm 5.0% 558	40.7% \pm 5.9% 549	* <u>41.4%</u> \pm 3.5% 1,378
Minority Youth	34.3% \pm 10.2% 195	36.6% \pm 13.2% 132	27.8% \pm 10.0% 91	38.6% \pm 10.1% 172	46.4% \pm 9.5% 167	42.3% \pm 6.1% 430

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (\pm %) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2013 preceded by an asterisk (*).


CURRENT CIGARETTE SMOKING — School Type, Gender, and Minority Populations


Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Middle School (MS)	10.0% $\pm 2.6\%$ 874	5.5% $\pm 1.8\%$ 508	2.7% $\pm 1.4\%$ 241	3.5% $\pm 1.2\%$ 307	2.3% $\pm 0.8\%$ 201	*2.9% $\pm 1.2\%$ 240
MS Females	12.7% $\pm 3.9\%$ 540	5.9% $\pm 2.4\%$ 271	2.7% $\pm 1.6\%$ 120	1.9% $\pm 1.2\%$ 85	1.4% $\pm 0.7\%$ 61	*2.5% $\pm 1.4\%$ 97
MS Males	7.5% $\pm 2.6\%$ 334	5.1% $\pm 1.9\%$ 237	2.7% $\pm 1.6\%$ 121	5.1% $\pm 1.8\%$ 222	3.3% $\pm 1.5\%$ 136	3.3% $\pm 1.6\%$ 142
MS Minorities	9.1% $\pm 4.4\%$ 187	6.3% $\pm 2.6\%$ 155	3.1% $\pm 2.4\%$ 85	5.9% $\pm 2.7\%$ 135	3.3% $\pm 1.6\%$ 87	3.6% $\pm 2.0\%$ 89
High School (HS)	31.0% $\pm 3.1\%$ 3,059	21.9% $\pm 3.0\%$ 2,303	17.4% $\pm 1.9\%$ 2,030	17.3% $\pm 2.0\%$ 1,772	17.2% $\pm 2.0\%$ 1,995	*13.6% $\pm 1.8\%$ 1,364
HS Females	31.3% $\pm 4.2\%$ 1,533	22.9% $\pm 3.8\%$ 1,216	16.9% $\pm 2.3\%$ 997	16.5% $\pm 2.5\%$ 857	15.7% $\pm 2.2\%$ 920	*13.2% $\pm 2.4\%$ 665
HS Males	30.5% $\pm 3.9\%$ 1,504	20.5% $\pm 3.2\%$ 1,064	17.8% $\pm 2.4\%$ 1,033	18.0% $\pm 2.6\%$ 911	18.7% $\pm 2.6\%$ 1,066	*13.9% $\pm 2.2\%$ 689
HS Minorities	25.7% $\pm 4.7\%$ 555	21.9% $\pm 5.0\%$ 542	16.2% $\pm 3.1\%$ 510	16.1% $\pm 3.1\%$ 416	15.1% $\pm 2.5\%$ 498	*13.0% $\pm 2.5\%$ 366

Harford County—Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Howard County

Tobacco Use by Youth: 2000–2013

Maryland Department of Health and Mental Hygiene
Center for Tobacco Prevention and Control

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Any Tobacco Product (All Youth Populations)	17.9% ± 3.5% 3,970	17.1% ± 3.5% 4,139	13.3% ± 1.7% 3,603	12.0% ± 1.5% 3,285	12.7% ± 2.6% 3,484	*7.7% ± 1.6% 2,045
Any Tobacco Product (Minority Youth)	15.7% ± 3.8% 940	17.8% ± 3.5% 1,467	12.4% ± 2.0% 1,426	12.4% ± 1.7% 1,355	14.5% ± 2.8% 1,929	*7.2% ± 1.8% 951
Cigarette	13.5% ± 3.3% 2,913	11.6% ± 2.9% 2,702	8.0% ± 1.2% 2,119	7.8% ± 1.1% 2,094	7.0% ± 1.7% 1,887	*4.2% ± 1.0% 1,069
Cigar	7.4% ± 1.8% 1,626	7.4% ± 1.9% 1,749	5.6% ± 0.9% 1,462	7.7% ± 1.2% 2,090	7.0% ± 1.7% 1,911	5.3% ± 1.2% 1,397
Pipe (tobacco)	3.6% ± 1.2% 798	4.6% ± 1.2% 1,094	4.1% ± 0.7% 1,101	4.0% ± 0.8% 1,061	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2013
Bidi	4.6% ± 1.2% 992	5.3% ± 1.4% 1,252	4.2% ± 0.8% 1,128	2.2% ± 0.5% 602	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2013
Kretek	3.3% ± 1.0% 721	4.7% ± 1.3% 1,112	4.6% ± 0.8% 1,238	2.3% ± 0.5% 614	KRETEK SPECIFIC DATA NOT AVAILA- BLE FOR 2010	KRETEK SPECIFIC DATA NOT AVAILA- BLE FOR 2013
Smokeless Tobacco	4.6% ± 1.4% 1,021	4.1% ± 1.2% 972	3.3% ± 0.7% 876	2.5% ± 0.5% 656	2.3% ± 0.6% 604	2.6% ± 0.6% 693

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	18.1% ± 2.9% 4,028	15.7% ± 2.9% 3,787	12.9% ± 1.7% 3,497	12.7% ± 1.4% 3,473	14.2% ± 2.6% 3,895	*8.3% ± 1.8% 2,139
Minority Youth	15.4% ± 3.4% 923	13.4% ± 2.9% 1,107	9.9% ± 1.6% 1,132	10.5% ± 1.7% 1,151	15.9% ± 3.0% 2,110	*7.7% ± 1.8% 978

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	38.5% ± 7.6% 815	34.5% ± 5.9% 594	41.2% ± 6.2% 502	30.8% ± 5.2% 521	46.9% ± 7.8% 473	49.0% ± 3.7% 1,646
Minority Youth	39.3% ± 12.1% 194	35.3% ± 10.7% 179	45.1% ± 10.8% 172	31.4% ± 8.1% 216	45.3% ± 10.9% 266	51.4% ± 5.7% 833

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2013 preceded by an asterisk (*).


CURRENT CIGARETTE SMOKING — School Type, Gender, and Minority Populations

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Middle School (MS)	4.1% \pm 1.1% 407	3.4% \pm 1.0% 371	1.7% \pm 0.8% 195	1.4% \pm 0.8% 168	1.4% \pm 1.0% 155	*1.5% \pm 0.8% 168
MS Females	3.2% \pm 1.4% 154	3.6% \pm 1.0% 193	0.8% \pm 0.9% 44	1.4% \pm 0.9% 77	1.0% \pm 1.2% 59	*0.9% \pm 0.6% 50
MS Males	4.9% \pm 2.0% 253	3.2% \pm 1.3% 178	2.5% \pm 1.2% 152	1.5% \pm 1.1% 91	1.8% \pm 1.0% 97	2.0% \pm 1.2% 118
MS Minorities	3.6% \pm 1.6% 97	5.0% \pm 2.6% 186	1.2% \pm 1.1% 66	1.6% \pm 1.1% 81	2.3% \pm 1.7% 133	1.4% \pm 0.8% 84
High School (HS)	21.5% \pm 3.0% 2,506	18.7% \pm 2.4% 2,331	13.0% \pm 1.4% 1,924	12.6% \pm 1.8% 1,926	11.1% \pm 1.4% 1,732	*6.3% \pm 1.0% 901
HS Females	20.4% \pm 3.4% 1,176	18.4% \pm 2.9% 1,130	10.4% \pm 1.6% 765	10.4% \pm 1.9% 788	9.1% \pm 1.6% 757	*5.0% \pm 1.0% 359
HS Males	22.5% \pm 4.2% 1,310	18.1% \pm 2.9% 1,128	15.6% \pm 2.0% 1,150	14.5% \pm 2.5% 1,123	13.2% \pm 2.1% 962	*7.3% \pm 1.4% 526
HS Minorities	18.0% \pm 4.6% 540	16.8% \pm 3.0% 690	12.2% \pm 2.0% 708	11.7% \pm 2.3% 686	12.4% \pm 1.8% 900	*6.2% \pm 1.4% 433

Howard County—Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Kent County

Tobacco Use by Youth: 2000–2013

Maryland Department of Health and Mental Hygiene
Center for Tobacco Prevention and Control

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Any Tobacco Product (All Youth Populations)	32.0% \pm 3.9% 463	29.4% \pm 4.0% 424	24.6% \pm 4.5% 310	22.8% \pm 3.8% 257	23.9% \pm 5.0% 254	*17.4% \pm 3.9% 181
Any Tobacco Product (Minority Youth)	31.6% \pm 5.1% 145	28.0% \pm 5.2% 133	22.6% \pm 6.5% 101	23.3% \pm 5.4% 85	27.9% \pm 7.0% 114	*14.4% \pm 5.3% 51
Cigarette	23.0% \pm 3.8% 317	21.7% \pm 3.4% 298	19.1% \pm 4.0% 231	17.6% \pm 3.1% 195	14.7% \pm 3.9% 153	*13.3% \pm 3.7% 129
Cigar	12.4% \pm 2.5% 176	11.3% \pm 2.1% 158	9.5% \pm 2.3% 114	11.9% \pm 3.4% 133	12.4% \pm 3.2% 130	8.4% \pm 2.0% 87
Pipe (tobacco)	5.8% \pm 1.7% 83	8.0% \pm 2.0% 112	7.7% \pm 2.5% 95	6.2% \pm 2.2% 69	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Bidi	6.5% \pm 1.9% 92	8.2% \pm 2.3% 116	7.7% \pm 2.4% 96	4.5% \pm 2.0% 50	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010	BIDI SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Kretek	5.5% \pm 1.5% 78	6.8% \pm 2.0% 95	8.1% \pm 2.5% 100	4.2% \pm 2.0% 47	KRETEK SPECIFIC DATA NOT AVAILA- BLE FOR 2010	KRETEK SPECIFIC DATA NOT AVAILA- BLE FOR 2013
Smokeless Tobacco	9.3% \pm 2.1% 132	8.7% \pm 2.0% 121	5.7% \pm 1.8% 69	6.9% \pm 2.5% 75	7.3% \pm 2.1% 73	9.2% \pm 2.5% 95

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	28.7% \pm 3.2% 414	24.0% \pm 3.1% 348	22.9% \pm 3.8% 288	20.6% \pm 2.7% 232	23.9% \pm 4.3% 254	*16.6% \pm 3.1% 166
Minority Youth	28.0% \pm 4.6% 128	19.7% \pm 4.1% 94	19.2% \pm 3.2% 85	17.9% \pm 4.8% 65	25.9% \pm 5.9% 106	*14.2% \pm 4.5% 49

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	36.3% \pm 9.3% 84	32.0% \pm 7.3% 68	31.2% \pm 10.5% 41	29.8% \pm 7.8% 46	30.6% \pm 10.2% 30	38.3% \pm 8.4% 96
Minority Youth	36.0% \pm 13.1% 28	29.1% \pm 15.0% 17	Not Available	29.0% \pm 13.8% 15	28.7% \pm 15.7% 10	55.1% \pm 14.7% 42

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (\pm %) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2013 preceded by an asterisk (*).


CURRENT CIGARETTE SMOKING — School Type, Gender, and Minority Populations


Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Middle School (MS)	8.0% ± 2.6% 50	12.9% ± 5.0% 83	6.0% ± 2.7% 28	7.2% ± 3.2% 33	4.1% ± 3.1% 19	4.3% ± 2.9% 19
MS Females	5.7% ± 3.3% 18	13.6% ± 5.9% 46	3.3% ± 2.7% 8	5.1% ± 3.7% 11	3.4% ± 3.1% 8	4.1% ± 3.9% 8
MS Males	10.3% ± 3.8% 31	12.1% ± 5.7% 37	8.4% ± 5.3% 20	8.9% ± 4.7% 23	4.2% ± 3.6% 9	4.4% ± 2.7% 11
MS Minorities	9.3% ± 3.7% 19	12.1% ± 5.4% 25	4.0% ± 3.1% 6	8.0% ± 5.6% 12	2.7% ± 2.3% 5	4.0% ± 3.9% 7
High School (HS)	35.3% ± 5.5% 267	29.5% ± 4.1% 215	27.5% ± 3.7% 203	25.2% ± 4.7% 161	23.1% ± 5.5% 135	*21.0% ± 6.1% 110
HS Females	38.2% ± 6.9% 152	26.5% ± 5.5% 97	24.4% ± 5.9% 87	25.5% ± 5.9% 76	21.4% ± 6.0% 67	*13.5% ± 7.1% 32
HS Males	32.2% ± 6.6% 115	32.0% ± 5.0% 115	30.6% ± 5.0% 116	24.9% ± 5.3% 86	25.1% ± 7.0% 68	26.9% ± 8.4% 75
HS Minorities	41.3% ± 7.7% 88	28.8% ± 5.9% 64	22.3% ± 7.2% 58	25.6% ± 7.2% 49	27.6% ± 6.1% 59	*13.4% ± 7.8% 21

Kent County—Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Montgomery County

Tobacco Use by Youth: 2000–2013

Maryland Department of Health and Mental Hygiene
Center for Tobacco Prevention and Control

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Any Tobacco Product (All Youth Populations)	16.6% \pm 4.1% 10,604	14.9% \pm 2.7%	14.3% \pm 1.7%	11.5% \pm 1.4%	13.1% \pm 2.3%	*9.7% \pm 2.0%
Any Tobacco Product (Minority Youth)	15.9% \pm 3.4% 4,507	15.0% \pm 2.4%	14.5% \pm 1.8%	11.7% \pm 1.6%	14.5% \pm 2.3%	*10.1% \pm 2.0%
Cigarette	12.1% \pm 3.4% 7,455	9.5% \pm 2.1%	9.0% \pm 1.3%	6.9% \pm 0.8%	7.1% \pm 1.5%	*5.9% \pm 1.2%
Cigar	6.6% \pm 1.8% 4,107	6.1% \pm 1.3%	6.2% \pm 1.0%	7.7% \pm 1.0%	7.1% \pm 1.5%	6.3% \pm 1.4%
Pipe (tobacco)	2.9% \pm 0.7% 1,853	3.7% \pm 0.8%	5.0% \pm 0.9%	4.6% \pm 0.9%	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Bidi	5.0% \pm 1.6% 3,087	4.6% \pm 0.9%	5.2% \pm 0.8%	3.0% \pm 0.5% 2,136	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010	BIDI SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Kretek	3.4% \pm 0.8% 2,113	3.1% \pm 0.8%	5.8% \pm 0.9% 4,219	2.5% \pm 0.6% 1,755	KRETEK SPECIFIC DATA NOT AVAILA- BLE FOR 2010	KRETEK SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Smokeless Tobacco	2.9% \pm 1.1% 1,836	2.8% \pm 0.8%	3.3% \pm 0.7%	2.6% \pm 0.6% 1,849	2.2% \pm 0.6% 1,563	3.9% \pm 0.8% 2,727

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	15.8% \pm 3.4% 10,100	13.4% \pm 2.8%	13.1% \pm 1.7%	11.7% \pm 1.3%	15.0% \pm 2.5%	*9.8% \pm 2.2%
Minority Youth	14.9% \pm 2.7% 4,232	12.4% \pm 2.5%	11.8% \pm 1.6%	10.6% \pm 1.1%	16.3% \pm 2.4%	10.2% \pm 2.4%

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	36.9% \pm 6.9% 1,955	43.7% \pm 3.5%	45.0% \pm 6.2%	39.4% \pm 5.6%	50.8% \pm 6.4%	*50.9% \pm 4.9%
Minority Youth	46.7% \pm 12.1% 987	49.4% \pm 5.3%	47.9% \pm 7.9%	41.6% \pm 6.1%	52.3% \pm 6.2%	54.3% \pm 5.9%

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (\pm %) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2013 preceded by an asterisk (*).


CURRENT CIGARETTE SMOKING — School Type, Gender, and Minority Populations


Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Middle School (MS)	3.7% $\pm 1.5\%$ 1,065	3.1% $\pm 1.0\%$ 981	3.0% $\pm 1.2\%$ 916	1.3% $\pm 0.7\%$ 408	2.7% $\pm 1.2\%$ 816	3.5% $\pm 1.4\%$ 1,055
MS Females	3.3% $\pm 2.1\%$ 450	2.3% $\pm 0.9\%$ 346	2.1% $\pm 1.3\%$ 320	1.0% $\pm 0.7\%$ 146	2.3% $\pm 1.6\%$ 362	2.6% $\pm 1.2\%$ 387
MS Males	3.8% $\pm 2.5\%$ 558	4.0% $\pm 1.8\%$ 635	3.6% $\pm 1.8\%$ 556	1.7% $\pm 1.1\%$ 262	3.1% $\pm 1.5\%$ 454	4.0% $\pm 2.2\%$ 600
MS Minorities	4.4% $\pm 1.7\%$ 566	3.9% $\pm 1.2\%$ 685	3.3% $\pm 1.5\%$ 553	1.8% $\pm 0.9\%$ 316	3.7% $\pm 1.7\%$ 706	4.5% $\pm 1.8\%$ 874
High School (HS)	19.4% $\pm 2.5\%$ 6,389	14.9% $\pm 2.0\%$ 5,494	13.4% $\pm 1.5\%$ 5,498	11.1% $\pm 1.3\%$ 4,563	10.4% $\pm 1.5\%$ 4,366	*7.8% $\pm 1.4\%$ 2,981
HS Females	19.6% $\pm 4.4\%$ 3,217	14.1% $\pm 2.5\%$ 2,583	12.8% $\pm 2.0\%$ 2,573	8.9% $\pm 1.3\%$ 1,823	8.2% $\pm 1.7\%$ 1,798	*7.0% $\pm 1.6\%$ 1,312
HS Males	18.8% $\pm 3.9\%$ 3,101	15.4% $\pm 1.9\%$ 2,836	14.0% $\pm 1.9\%$ 2,899	13.1% $\pm 2.0\%$ 2,700	12.7% $\pm 1.9\%$ 2,568	*8.1% $\pm 1.8\%$ 1,545
HS Minorities	15.4% $\pm 2.9\%$ 2,222	14.0% $\pm 1.9\%$ 2,683	12.9% $\pm 1.7\%$ 3,006	11.2% $\pm 1.3\%$ 2,543	11.1% $\pm 1.3\%$ 2,927	*8.0% $\pm 1.8\%$ 1,894

Montgomery County—Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Prince George's County

Tobacco Use by Youth: 2000–2013

Maryland Department of Health and Mental Hygiene
Center for Tobacco Prevention and Control

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Any Tobacco Product (All Youth Populations)	17.1% $\pm 3.2\%$ 10,459	15.3% $\pm 2.5\%$ 10,097	11.9% $\pm 1.5\%$ 8,341	15.3% $\pm 1.5\%$ 10,089	17.4% $\pm 2.2\%$ 11,020	12.9% $\pm 1.6\%$ 6,735
Any Tobacco Product (Minority Youth)	15.7% $\pm 3.2\%$ 8,126	14.5% $\pm 2.3\%$ 8,236	11.3% $\pm 1.4\%$ 7,347	14.8% $\pm 1.5\%$ 9,105	17.2% $\pm 2.1\%$ 10,419	12.4% $\pm 1.4\%$ 6,058
Cigarette	10.6% $\pm 2.5\%$ 6,178	8.5% $\pm 1.7\%$ 5,269	6.3% $\pm 1.0\%$ 4,254	7.7% $\pm 0.9\%$ 4,964	7.6% $\pm 1.2\%$ 4,715	*6.4% $\pm 1.0\%$ 3,087
Cigar	7.4% $\pm 1.7\%$ 4,369	6.1% $\pm 1.4\%$ 3,784	4.4% $\pm 0.8\%$ 2,933	11.2% $\pm 1.4\%$ 7,170	9.9% $\pm 1.6\%$ 6,191	9.3% $\pm 1.4\%$ 4,698
Pipe (tobacco)	4.2% $\pm 1.5\%$ 2,487	4.8% $\pm 1.2\%$ 2,973	4.5% $\pm 0.9\%$ 3,100	6.7% $\pm 1.4\%$ 4,159	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2013
Bidi	5.8% $\pm 1.4\%$ 3,357	5.0% $\pm 1.2\%$ 3,174	4.6% $\pm 0.9\%$ 3,149	4.9% $\pm 1.0\%$ 2,998	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2013
Kretek	2.9% $\pm 1.1\%$ 1,743	4.2% $\pm 1.1\%$ 2,592	5.1% $\pm 1.0\%$ 3,467	4.4% $\pm 1.1\%$ 2,691	KRETEK SPECIFIC DATA NOT AVAILA- BLE FOR 2010	KRETEK SPECIFIC DATA NOT AVAILA- BLE FOR 2013
Smokeless Tobacco	2.4% $\pm 1.0\%$ 1,423	2.9% $\pm 0.9\%$ 1,808	2.6% $\pm 0.6\%$ 1,759	3.2% $\pm 0.6\%$ 1,947	2.8% $\pm 0.6\%$ 1,631	*5.3% $\pm 1.0\%$ 2,698

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	14.9% $\pm 2.0\%$ 9,115	13.3% $\pm 1.7\%$ 8,759	10.8% $\pm 1.3\%$ 7,547	12.7% $\pm 0.8\%$ 8,371	17.9% $\pm 1.9\%$ 11,346	12.3% $\pm 1.4\%$ 6,028
Minority Youth	14.0% $\pm 2.1\%$ 7,262	12.6% $\pm 1.6\%$ 7,149	10.4% $\pm 1.3\%$ 6,789	12.3% $\pm 1.0\%$ 7,545	17.7% $\pm 1.9\%$ 10,757	12.4% $\pm 1.4\%$ 5,725

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	39.1% $\pm 7.9\%$ 1,839	43.9% $\pm 6.4\%$ 1,686	51.6% $\pm 7.0\%$ 1,840	52.0% $\pm 6.0\%$ 2,974	55.3% $\pm 4.9\%$ 2,171	48.9% $\pm 4.3\%$ 4,852
Minority Youth	48.9% $\pm 11.2\%$ 1,639	47.5% $\pm 6.6\%$ 1,431	53.9% $\pm 7.5\%$ 1,731	55.1% $\pm 6.4\%$ 2,867	55.0% $\pm 5.2\%$ 2,064	48.7% $\pm 4.5\%$ 4,570

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals ($\pm \%$) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2013 preceded by an asterisk (*).


CURRENT CIGARETTE SMOKING — School Type, Gender and Minority Populations


Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Middle School (MS)	5.1% ± 3.0% 1,363	3.8% ± 1.3% 1,098	3.7% ± 1.4% 1,063	3.4% ± 1.3% 925	3.9% ± 1.2% 1,001	5.4% ± 1.4% 1,079
MS Females	3.1% ± 1.0% 411	3.7% ± 2.3% 536	3.4% ± 2.3% 486	2.2% ± 0.9% 300	4.1% ± 1.2% 548	*6.5% ± 1.6% 648
MS Males	6.9% ± 4.9% 925	3.7% ± 0.9% 541	3.9% ± 1.7% 577	4.6% ± 2.5% 625	3.6% ± 1.7% 453	4.1% ± 2.0% 415
MS Minorities	5.0% ± 3.4% 1,136	3.7% ± 1.4% 910	3.8% ± 1.5% 1,019	3.0% ± 1.3% 761	3.8% ± 1.3% 944	5.6% ± 1.6% 1,056
High School (HS)	15.3% ± 3.0% 4,816	12.8% ± 1.5% 4,172	8.3% ± 1.1% 3,191	10.9% ± 1.2% 4,039	10.2% ± 1.2% 3,714	*7.1% ± 1.4% 2,008
HS Females	16.0% ± 3.6% 2,577	12.1% ± 2.2% 2,033	6.4% ± 1.3% 1,250	8.0% ± 1.4% 1,475	6.7% ± 1.1% 1,268	*4.8% ± 1.2% 693
HS Males	14.1% ± 3.4% 2,155	12.8% ± 2.2% 2,011	10.3% ± 1.6% 1,941	13.9% ± 1.8% 2,558	14.0% ± 2.1% 2,423	8.6% ± 2.2% 1,172
HS Minorities	12.1% ± 3.0% 3,202	11.0% ± 1.4% 3,121	7.4% ± 1.0% 2,615	10.3% ± 1.4% 3,545	9.9% ± 1.3% 3,419	*6.5% ± 1.2% 1,751

Prince George's County—Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Queen Anne's County

Tobacco Use by Youth: 2000–2013

Maryland Department of Health and Mental Hygiene
Center for Tobacco Prevention and Control

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Any Tobacco Product (All Youth Populations)	25.7% ± 3.7% 896	22.4% ± 2.2% 833	21.4% ± 2.8% 899	19.1% ± 2.1% 802	20.5% ± 2.0% 819	*16.0% ± 2.2% 619
Any Tobacco Product (Minority Youth)	32.3% ± 6.2% 170	30.1% ± 4.8% 172	31.3% ± 4.9% 207	32.7% ± 5.1% 220	30.2% ± 4.2% 265	*21.2% ± 4.3% 123
Cigarette	20.4% ± 3.3% 683	17.4% ± 2.0% 622	15.7% ± 2.4% 641	14.7% ± 1.8% 610	13.9% ± 1.7% 547	*10.7% ± 1.8% 393
Cigar	10.0% ± 1.8% 342	7.8% ± 1.2% 283	9.4% ± 1.5% 381	10.8% ± 1.6% 450	10.7% ± 1.5% 426	9.1% ± 1.6% 348
Pipe (tobacco)	4.3% ± 1.1% 148	5.3% ± 1.2% 194	5.2% ± 1.1% 217	7.5% ± 1.3% 311	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2013
Bidi	4.6% ± 1.1% 157	6.3% ± 1.2% 230	5.7% ± 1.1% 236	5.0% ± 1.0% 208	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2013
Kretek	3.8% ± 1.0% 132	5.8% ± 1.3% 209	5.9% ± 1.2% 243	5.2% ± 1.0% 214	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2013
Smokeless Tobacco	4.3% ± 1.0% 149	5.3% ± 1.1% 192	5.0% ± 1.0% 204	5.2% ± 0.9% 208	5.0% ± 0.9% 191	*8.3% ± 1.8% 318

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	25.0% ± 3.4% 872	18.1% ± 1.7% 675	20.3% ± 2.5% 850	16.3% ± 1.6% 685	21.2% ± 1.9% 847	*14.4% ± 2.0% 531
Minority Youth	26.7% ± 7.3% 141	19.4% ± 3.9% 111	20.6% ± 3.7% 135	18.4% ± 3.3% 123	29.8% ± 3.6% 261	18.6% ± 3.7% 99

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	24.8% ± 5.3% 115	28.2% ± 5.4% 109	29.5% ± 5.8% 114	28.0% ± 5.3% 113	38.8% ± 5.8% 114	*40.7% ± 4.9% 316
Minority Youth	26.5% ± 11.0% 19	27.3% ± 12.2% 13	25.9% ± 11.5% 16	28.3% ± 11.8% 19	31.1% ± 12.8% 20	*48.7% ± 9.8% 61

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (±%) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2013 preceded by an asterisk (*).


CURRENT CIGARETTE SMOKING — School Type, Gender, and Minority Populations


Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Middle School (MS)	9.6% \pm 3.6% 152	5.3% \pm 1.8% 91	4.3% \pm 1.3% 74	2.8% \pm 1.0% 49	3.1% \pm 1.0% 52	4.7% \pm 2.0% 80
MS Females	8.3% \pm 3.7% 62	5.4% \pm 2.4% 44	4.1% \pm 1.6% 34	2.3% \pm 0.9% 19	2.2% \pm 1.1% 18	4.1% \pm 2.4% 34
MS Males	10.8% \pm 4.5% 91	5.2% \pm 10.9% 47	4.6% \pm 1.7% 40	3.3% \pm 1.6% 30	4.1% \pm 1.5% 34	5.4% \pm 2.5% 46
MS Minorities	15.3% \pm 7.9% 35	7.9% \pm 3.3% 20	11.0% \pm 4.2% 27	5.3% \pm 2.7% 14	5.2% \pm 2.5% 18	9.8% \pm 5.3% 25
High School (HS)	30.1% \pm 3.1% 531	28.6% \pm 3.1% 531	23.8% \pm 3.0% 567	23.5% \pm 2.9% 561	21.8% \pm 2.7% 495	*15.9% \pm 2.9% 313
HS Females	30.9% \pm 4.5% 261	29.1% \pm 3.8% 260	23.6% \pm 4.0% 289	20.7% \pm 3.4% 245	19.3% \pm 3.1% 227	*11.8% \pm 2.7% 117
HS Males	29.5% \pm 4.2% 268	27.8% \pm 4.6% 263	24.1% \pm 3.2% 278	26.3% \pm 3.7% 317	24.1% \pm 3.7% 261	*19.2% \pm 4.1% 186
HS Minorities	33.5% \pm 6.9% 85	33.0% \pm 7.4% 92	27.8% \pm 5.7% 101	35.2% \pm 7.0% 134	33.5% \pm 5.6% 161	22.2% \pm 5.9% 63

Queen Anne's County—Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Somerset County

Tobacco Use by Youth: 2000–2013

Maryland Department of Health and Mental Hygiene
Center for Tobacco Prevention and Control

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Any Tobacco Product (All Youth Populations)	33.9% \pm 4.1% 504	27.3% \pm 4.3% 390	20.4% \pm 3.4% 303	23.2% \pm 3.2% 326	25.5% \pm 3.9% 331	*16.5% \pm 4.1% 215
Any Tobacco Product (Minority Youth)	30.6% \pm 4.4% 202	27.3% \pm 5.2% 190	15.1% \pm 3.5% 114	24.6% \pm 4.7% 179	24.9% \pm 3.8% 178	*15.8% \pm 4.5% 107
Cigarette	29.2% \pm 4.5% 415	20.7% \pm 3.7% 275	13.5% \pm 3.1% 194	15.5% \pm 2.4% 213	17.3% \pm 3.2% 219	*13.5% \pm 2.9% 167
Cigar	14.6% \pm 3.4% 210	11.2% \pm 2.5% 152	8.5% \pm 1.9% 120	16.6% \pm 2.9% 228	14.7% \pm 3.0% 188	9.5% \pm 3.5% 123
Pipe (tobacco)	6.4% \pm 2.1% 94	8.3% \pm 2.3% 113	5.7% \pm 1.5% 82	9.9% \pm 2.5% 135	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Bidi	6.4% \pm 1.7% 91	7.6% \pm 2.0% 103	5.3% \pm 1.5% 76	7.0% \pm 1.6% 95	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010	BIDI SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Kretek	5.6% \pm 1.9% 82	7.3% \pm 2.3% 99	6.1% \pm 1.7% 89	5.7% \pm 1.7% 78	KRETEK SPECIFIC DATA NOT AVAILA- BLE FOR 2010	KRETEK SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Smokeless Tobacco	5.4% \pm 1.8% 78	6.5% \pm 2.8% 86	4.6% \pm 1.4% 65	5.2% \pm 1.5% 68	6.5% \pm 2.0% 79	8.4% \pm 3.1% 109

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	26.4% \pm 3.1% 393	23.3% \pm 3.4% 334	18.3% \pm 2.5% 272	21.4% \pm 3.5% 299	26.4% \pm 3.8% 343	*15.0% \pm 2.7% 183
Minority Youth	24.6% \pm 3.5% 162	22.1% \pm 4.1% 155	12.8% \pm 2.6% 97	23.0% \pm 4.3% 166	25.6% \pm 4.3% 183	*15.5% \pm 2.9% 96

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	38.7% \pm 7.8% 121	39.4% \pm 8.4% 67	28.1% \pm 9.6% 36	37.6% \pm 7.6% 64	37.2% \pm 7.5% 56	47.6% \pm 9.4% 124
Minority Youth	32.8% \pm 10.8% 30	44.4% \pm 18.1% 25	Not Available	42.8% \pm 11.9% 37	41.5% \pm 10.8% 31	54.2% \pm 12.3% 72

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (\pm %) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2013 preceded by an asterisk (*).


CURRENT CIGARETTE SMOKING — School Type, Gender, and Minority Populations


Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Middle School (MS)	17.5% \pm 4.3% 113	14.4% \pm 4.9% 88	6.9% \pm 2.3% 47	8.3% \pm 2.9% 53	7.6% \pm 2.9% 42	*6.1% \pm 2.4% 36
MS Females	13.1% \pm 6.9% 41	10.0% \pm 5.2% 31	8.9% \pm 3.6% 29	6.9% \pm 3.2% 20	5.0% \pm 2.5% 14	4.9% \pm 2.5% 15
MS Males	21.1% \pm 5.1% 68	18.6% \pm 6.2% 54	5.2% \pm 2.0% 19	9.3% \pm 4.5% 32	10.2% \pm 4.5% 28	*7.4% \pm 3.7% 21
MS Minorities	16.0% \pm 6.1% 48	12.1% \pm 5.6% 38	5.3% \pm 2.5% 18	11.6% \pm 4.6% 38	7.1% \pm 3.2% 23	*5.4% \pm 2.5% 17
High School (HS)	38.9% \pm 8.0% 302	26.1% \pm 5.4% 187	19.5% \pm 4.4% 147	21.5% \pm 3.4% 160	24.9% \pm 5.0% 177	*20.1% \pm 4.9% 131
HS Females	35.1% \pm 10.3% 142	21.4% \pm 6.0% 83	17.0% \pm 4.7% 69	14.7% \pm 4.1% 55	25.0% \pm 5.3% 93	*14.0% \pm 5.3% 44
HS Males	42.5% \pm 8.0% 158	31.2% \pm 7.2% 101	22.3% \pm 6.6% 78	28.3% \pm 5.9% 104	24.9% \pm 6.5% 84	*22.0% \pm 6.7% 69
HS Minorities	37.2% \pm 7.8% 117	22.3% \pm 6.9% 70	10.7% \pm 4.7% 41	16.6% \pm 5.0% 63	22.7% \pm 4.8% 83	*18.4% \pm 4.7% 58

Somerset County—Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





St. Mary's County

Tobacco Use by Youth: 2000–2013

Maryland Department of Health and Mental Hygiene
Center for Tobacco Prevention and Control

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Any Tobacco Product (All Youth Populations)	24.0% $\pm 3.0\%$ 1,776	23.7% $\pm 3.2\%$ 1,841	<u>14.7%</u> $\pm 1.9\%$ 1,282	16.1% $\pm 2.0\%$ 1,402	15.7% $\pm 1.9\%$ 1,369	*14.2% $\pm 1.8\%$ 1,174
Any Tobacco Product (Minority Youth)	22.6% $\pm 4.0\%$ 483	26.1% $\pm 5.3\%$ 583	<u>16.3%</u> $\pm 3.0\%$ 430	18.1% $\pm 3.1\%$ 418	18.2% $\pm 3.0\%$ 458	*15.2% $\pm 2.7\%$ 349
Cigarette	19.4% $\pm 2.9\%$ 1,363	18.0% $\pm 2.9\%$ 1,345	<u>10.2%</u> $\pm 1.6\%$ 870	11.8% $\pm 1.6\%$ 1,012	10.1% $\pm 1.4\%$ 864	*10.5% $\pm 1.4\%$ 819
Cigar	10.1% $\pm 1.6\%$ 729	8.0% $\pm 1.6\%$ 603	5.8% $\pm 1.0\%$ 490	<u>9.5%</u> $\pm 1.5\%$ 813	8.1% $\pm 1.4\%$ 696	9.0% $\pm 1.4\%$ 732
Pipe (tobacco)	3.9% $\pm 1.0\%$ 286	5.0% $\pm 1.6\%$ 377	3.5% $\pm 0.7\%$ 299	5.2% $\pm 1.0\%$ 444	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Bidi	4.9% $\pm 1.2\%$ 345	5.6% $\pm 1.5\%$ 420	3.3% $\pm 0.7\%$ 286	3.7% $\pm 0.8\%$ 316	BIDI SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	BIDI SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Kretek	3.2% $\pm 1.0\%$ 234	5.3% $\pm 1.6\%$ 399	3.7% $\pm 0.7\%$ 314	3.7% $\pm 0.8\%$ 311	KRETEK SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	KRETEK SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Smokeless Tobacco	3.5% $\pm 1.0\%$ 255	4.2% $\pm 1.5\%$ 309	2.7% $\pm 0.6\%$ 234	3.5% $\pm 0.8\%$ 296	3.3% $\pm 0.7\%$ 274	<u>5.6%</u> $\pm 1.2\%$ 459

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	22.5% $\pm 2.6\%$ 1,662	19.4% $\pm 2.0\%$ 1,511	15.5% $\pm 2.0\%$ 1,346	16.3% $\pm 2.0\%$ 1,421	16.5% $\pm 1.9\%$ 1,435	*14.5% $\pm 1.6\%$ 1,154
Minority Youth	23.1% $\pm 4.3\%$ 492	19.6% $\pm 4.2\%$ 438	14.8% $\pm 2.9\%$ 390	15.4% $\pm 2.6\%$ 354	19.4% $\pm 3.0\%$ 488	*15.4% $\pm 2.7\%$ 337

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	32.7% $\pm 6.0\%$ 322	25.2% $\pm 5.3\%$ 202	<u>39.2%</u> $\pm 6.4\%$ 247	35.2% $\pm 5.5\%$ 281	45.0% $\pm 6.2\%$ 253	38.0% $\pm 4.1\%$ 625
Minority Youth	37.2% $\pm 9.6\%$ 66	<u>15.7%</u> $\pm 10.3\%$ 26	<u>40.4%</u> $\pm 12.9\%$ 62	40.1% $\pm 13.4\%$ 79	51.3% $\pm 10.7\%$ 89	40.5% $\pm 7.4\%$ 190

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (\pm %) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2013 preceded by an asterisk (*).


CURRENT CIGARETTE SMOKING — School Type, Gender, and Minority Populations


Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Middle School (MS)	7.5% \pm 2.5% 235	8.5% \pm 2.1% 285	<u>3.2%</u> \pm 1.3% 114	4.3% \pm 1.8% 153	3.7% \pm 1.2% 139	5.5% \pm 2.0% 193
MS Females	6.1% \pm 2.8% 91	7.5% \pm 2.8% 127	<u>2.7%</u> \pm 1.5% 48	3.2% \pm 2.1% 56	3.4% \pm 1.4% 64	4.4% \pm 2.2% 78
MS Males	8.6% \pm 3.1% 141	9.2% \pm 2.9% 154	<u>3.7%</u> \pm 1.8% 66	5.3% \pm 2.3% 97	4.1% \pm 1.8% 76	6.7% \pm 2.5% 114
MS Minorities	9.3% \pm 4.2% 92	9.9% \pm 3.6% 99	4.1% \pm 2.4% 46	5.1% \pm 2.6% 48	3.1% \pm 1.8% 33	<u>8.1%</u> \pm 2.7% 80
High School (HS)	29.0% \pm 3.8% 1,129	26.0% \pm 4.5% 1,060	<u>15.4%</u> \pm 1.9% 756	17.2% \pm 2.4% 859	14.9% \pm 2.1% 724	*14.6% \pm 2.2% 626
HS Females	26.8% \pm 3.9% 524	30.4% \pm 6.7% 624	<u>15.5%</u> \pm 2.2% 387	15.9% \pm 2.9% 389	12.6% \pm 2.5% 313	*12.3% \pm 2.4% 267
HS Males	31.1% \pm 5.1% 596	21.6% \pm 5.5% 433	15.4% \pm 2.5% 369	18.5% \pm 3.2% 468	17.2% \pm 2.7% 406	*16.1% \pm 2.9% 340
HS Minorities	20.4% \pm 4.7% 201	22.9% \pm 7.4% 249	14.1% \pm 2.7% 199	16.7% \pm 4.0% 220	15.4% \pm 3.6% 213	*11.9% \pm 3.1% 133

St. Mary's County—Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Talbot County

Tobacco Use by Youth: 2000–2013

Maryland Department of Health and Mental Hygiene
Center for Tobacco Prevention and Control

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Any Tobacco Product (All Youth Populations)	28.3% $\pm 3.2\%$ 633	26.1% $\pm 3.1\%$ 606	23.5% $\pm 3.5\%$ 570	23.0% $\pm 2.9\%$ 547	21.4% $\pm 3.2\%$ 483	*15.8% $\pm 2.5\%$ 335
Any Tobacco Product (Minority Youth)	24.1% $\pm 4.3\%$ 146	25.6% $\pm 4.6\%$ 170	22.6% $\pm 5.2\%$ 171	26.7% $\pm 4.8\%$ 198	25.2% $\pm 4.8\%$ 196	17.1% $\pm 4.1\%$ 116
Cigarette	23.1% $\pm 3.1\%$ 494	18.1% $\pm 2.4\%$ 399	17.7% $\pm 3.0\%$ 411	17.3% $\pm 2.5\%$ 400	15.0% $\pm 2.4\%$ 331	*10.3% $\pm 2.0\%$ 203
Cigar	12.6% $\pm 2.0\%$ 276	12.1% $\pm 2.1\%$ 270	8.6% $\pm 1.6\%$ 200	13.3% $\pm 2.3\%$ 310	10.6% $\pm 2.1\%$ 237	*8.7% $\pm 1.8\%$ 183
Pipe (tobacco)	5.7% $\pm 1.5\%$ 127	6.9% $\pm 1.8\%$ 155	6.7% $\pm 1.6\%$ 160	8.4% $\pm 1.8\%$ 195	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2013
Bidi	5.6% $\pm 1.3\%$ 120	7.7% $\pm 1.9\%$ 173	6.9% $\pm 1.9\%$ 164	5.8% $\pm 1.5\%$ 134	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2013
Kretek	4.5% $\pm 1.2\%$ 98	6.6% $\pm 1.6\%$ 147	7.2% $\pm 1.8\%$ 172	5.6% $\pm 1.2\%$ 131	KRETEK SPECIFIC DATA NOT AVAILA- BLE FOR 2010	KRETEK SPECIFIC DATA NOT AVAILA- BLE FOR 2013
Smokeless Tobacco	4.1% $\pm 1.1\%$ 90	5.7% $\pm 1.2\%$ 125	6.0% $\pm 1.4\%$ 140	6.5% $\pm 1.5\%$ 146	6.1% $\pm 1.4\%$ 131	*7.6% $\pm 1.8\%$ 160

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	26.7% $\pm 3.2\%$ 598	20.7% $\pm 2.3\%$ 480	21.7% $\pm 3.2\%$ 525	19.3% $\pm 2.1\%$ 457	21.6% $\pm 2.7\%$ 487	*13.3% $\pm 2.2\%$ 266
Minority Youth	23.5% $\pm 5.0\%$ 142	17.9% $\pm 3.3\%$ 118	15.5% $\pm 3.8\%$ 116	17.7% $\pm 3.6\%$ 131	23.7% $\pm 4.1\%$ 184	15.3% $\pm 3.7\%$ 96

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	26.5% $\pm 6.0\%$ 84	33.6% $\pm 7.0\%$ 74	35.1% $\pm 6.3\%$ 100	29.5% $\pm 6.3\%$ 78	42.0% $\pm 9.3\%$ 68	*45.3% $\pm 6.5\%$ 192
Minority Youth	38.2% $\pm 11.8\%$ 27	26.8% $\pm 14.7\%$ 10	44.6% $\pm 14.7\%$ 28	32.7% $\pm 11.5\%$ 28	34.8% $\pm 15.3\%$ 17	48.7% $\pm 9.0\%$ 69

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (\pm %) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2013 preceded by an asterisk (*).


CURRENT CIGARETTE SMOKING — School Type, Gender, and Minority Populations


Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Middle School (MS)	9.3% $\pm 4.1\%$ 91	7.2% $\pm 2.7\%$ 72	5.9% $\pm 1.9\%$ 57	7.5% $\pm 3.1\%$ 70	4.0% $\pm 1.9\%$ 36	3.6% $\pm 1.6\%$ 30
MS Females	8.7% $\pm 4.9\%$ 41	6.4% $\pm 2.6\%$ 30	4.4% $\pm 1.9\%$ 22	6.9% $\pm 3.7\%$ 33	3.2% $\pm 2.2\%$ 14	2.6% $\pm 2.0\%$ 11
MS Males	10.0% $\pm 4.4\%$ 50	8.1% $\pm 3.3\%$ 42	7.5% $\pm 2.8\%$ 35	8.1% $\pm 3.8\%$ 38	4.8% $\pm 2.6\%$ 22	4.1% $\pm 2.4\%$ 18
MS Minorities	11.0% $\pm 6.2\%$ 31	8.1% $\pm 4.2\%$ 26	6.1% $\pm 4.0\%$ 19	9.5% $\pm 5.4\%$ 29	3.6% $\pm 2.4\%$ 11	4.6% $\pm 2.5\%$ 14
High School (HS)	34.5% $\pm 3.8\%$ 403	27.0% $\pm 3.4\%$ 327	26.2% $\pm 3.4\%$ 354	23.9% $\pm 3.6\%$ 330	22.4% $\pm 3.4\%$ 296	*15.4% $\pm 3.3\%$ 173
HS Females	32.5% $\pm 5.1\%$ 177	26.1% $\pm 4.7\%$ 153	25.2% $\pm 4.7\%$ 160	23.2% $\pm 5.3\%$ 160	19.4% $\pm 3.6\%$ 129	*15.0% $\pm 5.1\%$ 83
HS Males	36.1% $\pm 4.6\%$ 224	27.5% $\pm 4.0\%$ 170	26.8% $\pm 4.3\%$ 191	24.5% $\pm 3.7\%$ 169	25.5% $\pm 5.4\%$ 165	*14.3% $\pm 4.1\%$ 80
HS Minorities	22.5% $\pm 5.0\%$ 61	26.4% $\pm 5.4\%$ 77	19.9% $\pm 6.1\%$ 77	26.1% $\pm 5.8\%$ 106	25.8% $\pm 5.3\%$ 116	18.1% $\pm 5.5\%$ 60

Talbot County—Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Washington County

Tobacco Use by Youth: 2000–2013

Maryland Department of Health and Mental Hygiene
Center for Tobacco Prevention and Control

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Any Tobacco Product (All Youth Populations)	28.0% \pm 3.5% 2,770	22.4% \pm 2.7% 2,321	20.9% \pm 2.5% 2,319	20.0% \pm 2.0% 2,152	21.7% \pm 2.5% 2,457	*18.4% \pm 1.8% 2,045
Any Tobacco Product (Minority Youth)	29.5% \pm 6.6% 446	28.4% \pm 4.5% 464	21.8% \pm 3.4% 532	20.4% \pm 3.7% 421	27.3% \pm 4.1% 730	20.6% \pm 3.1% 531
Cigarette	22.9% \pm 3.2% 2,184	17.5% \pm 2.6% 1,754	14.9% \pm 2.1% 1,609	14.5% \pm 1.8% 1,535	14.2% \pm 2.0% 1,580	*11.5% \pm 1.4% 1,203
Cigar	9.9% \pm 1.6% 967	7.2% \pm 1.3% 735	8.5% \pm 1.3% 923	11.2% \pm 1.4% 1,182	10.5% \pm 1.5% 1,165	9.1% \pm 1.4% 994
Pipe (tobacco)	3.6% \pm 0.8% 358	4.0% \pm 0.9% 408	4.7% \pm 0.9% 518	6.0% \pm 1.1% 620	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Bidi	4.4% \pm 1.1% 427	4.3% \pm 0.9% 438	4.4% \pm 0.8% 482	3.8% \pm 0.8% 389	BIDI SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	BIDI SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Kretek	3.7% \pm 0.9% 364	3.4% \pm 0.9% 348	4.6% \pm 0.8% 508	4.0% \pm 0.9% 415	KRETEK SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	KRETEK SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Smokeless Tobacco	5.8% \pm 1.3% 563	4.6% \pm 1.1% 461	5.1% \pm 1.0% 553	6.2% \pm 0.9% 624	6.7% \pm 1.2% 709	*9.9% \pm 1.4% 1,091

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	26.0% \pm 2.9% 2,569	22.9% \pm 2.5% 2,373	22.3% \pm 2.5% 2,484	19.1% \pm 1.8% 2,054	21.1% \pm 2.3% 2,387	*17.0% \pm 1.6% 1,809
Minority Youth	27.6% \pm 5.5% 417	24.3% \pm 5.0% 398	20.3% \pm 3.5% 495	17.3% \pm 3.3% 359	25.6% \pm 3.7% 685	*18.9% \pm 2.9% 457

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	36.6% \pm 4.5% 622	40.2% \pm 5.9% 517	37.6% \pm 4.8% 434	34.3% \pm 5.1% 435	35.2% \pm 5.5% 382	38.8% \pm 3.3% 1,019
Minority Youth	47.2% \pm 14.9% 120	36.4% \pm 14.2% 74	44.2% \pm 12.1% 94	34.2% \pm 11.0% 83	40.4% \pm 9.7% 105	37.3% \pm 6.1% 246

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (\pm %) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2013 preceded by an asterisk (*).


CURRENT CIGARETTE SMOKING — School Type, Gender, and Minority Populations


Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Middle School (MS)	12.6% ± 3.9% 559	8.9% ± 2.7% 408	5.4% ± 1.5% 253	4.1% ± 1.6% 197	4.9% ± 1.7% 242	*6.2% ± 2.0% 298
MS Females	12.0% ± 4.8% 259	8.1% ± 3.0% 183	4.9% ± 2.1% 110	3.2% ± 1.5% 77	3.6% ± 1.8% 91	7.1% ± 3.1% 164
MS Males	13.2% ± 4.5% 301	9.6% ± 3.4% 222	5.9% ± 1.9% 143	4.9% ± 2.3% 120	6.0% ± 2.5% 144	*5.4% ± 2.0% 130
MS Minorities	10.9% ± 7.4% 83	16.7% ± 5.7% 142	6.8% ± 3.0% 80	4.9% ± 2.8% 50	6.3% ± 3.2% 76	9.7% ± 3.9% 109
High School (HS)	31.9% ± 3.1% 1,624	24.8% ± 3.3% 1,345	22.1% ± 2.3% 1,356	23.2% ± 2.8% 1,339	21.8% ± 2.6% 1,337	*15.9% ± 2.0% 907
HS Females	34.3% ± 4.3% 862	26.2% ± 4.4% 717	23.0% ± 2.7% 699	23.0% ± 3.3% 646	23.6% ± 3.6% 757	*14.7% ± 2.5% 424
HS Males	29.6% ± 3.7% 758	23.3% ± 4.0% 617	21.1% ± 3.0% 654	23.3% ± 3.4% 690	19.7% ± 2.7% 576	*16.7% ± 2.5% 468
HS Minorities	31.4% ± 6.5% 205	26.1% ± 6.4% 172	20.8% ± 3.7% 234	24.1% ± 4.6% 245	25.3% ± 3.9% 349	*17.2% ± 3.7% 214

Washington County—Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Wicomico County

Tobacco Use by Youth: 2000–2013

Maryland Department of Health and Mental Hygiene
Center for Tobacco Prevention and Control

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Any Tobacco Product (All Youth Populations)	27.0% \pm 3.3% 1,817	23.1% \pm 3.5% 1,554	<u>17.1%</u> \pm 2.0% 1,196	19.0% \pm 1.8% 1,298	22.5% \pm 2.6% 1,514	* <u>16.0%</u> \pm 1.8% 1,034
Any Tobacco Product (Minority Youth)	29.1% \pm 4.5% 673	21.4% \pm 4.0% 599	<u>16.8%</u> \pm 2.5% 555	18.7% \pm 2.2% 572	<u>24.5%</u> \pm 3.0% 801	* <u>16.7%</u> \pm 2.4% 511
Cigarette	22.4% \pm 3.1% 1,450	18.8% \pm 3.1% 1,214	<u>12.0%</u> \pm 1.6% 814	12.9% \pm 1.3% 865	15.2% \pm 2.0% 997	* <u>10.9%</u> \pm 1.4% 661
Cigar	12.7% \pm 2.0% 835	8.9% \pm 2.1% 577	7.9% \pm 1.3% 532	<u>12.4%</u> \pm 1.5% 823	13.5% \pm 1.9% 891	11.0% \pm 1.4% 693
Pipe (tobacco)	4.0% \pm 1.1% 267	5.4% \pm 1.7% 350	3.6% \pm 0.8% 247	<u>5.9%</u> \pm 1.1% 386	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Bidi	3.9% \pm 1.1% 256	4.3% \pm 1.4% 281	3.6% \pm 0.9% 243	4.0% \pm 0.8% 262	BIDI SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	BIDI SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Kretek	3.4% \pm 1.1% 229	3.3% \pm 1.2% 212	4.1% \pm 1.0% 276	3.5% \pm 0.9% 229	KRETEK SPECIFIC DATA NOT AVAIL- ABLE FOR 2010	KRETEK SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Smokeless Tobacco	2.5% \pm 0.8% 169	3.3% \pm 1.0% 212	3.3% \pm 0.7% 225	4.0% \pm 0.8% 258	5.1% \pm 1.0% 316	* <u>6.6%</u> \pm 1.2% 418

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	24.6% \pm 2.5% 1,656	20.9% \pm 3.2% 1,407	18.9% \pm 2.1% 1,332	18.6% \pm 1.8% 1,266	22.3% \pm 2.4% 1,497	* <u>14.2%</u> \pm 1.6% 876
Minority Youth	23.8% \pm 3.0% 552	<u>16.8%</u> \pm 3.8% 471	15.8% \pm 2.1% 526	17.1% \pm 2.1% 519	<u>23.4%</u> \pm 2.8% 768	* <u>14.3%</u> \pm 2.2% 415

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	34.3% \pm 5.7% 370	35.0% \pm 8.2% 284	40.6% \pm 5.9% 240	36.7% \pm 5.9% 248	33.1% \pm 5.5% 187	41.8% \pm 4.7% 563
Minority Youth	31.0% \pm 8.0% 116	36.1% \pm 13.6% 94	41.7% \pm 10.4% 91	44.7% \pm 8.8% 128	36.0% \pm 7.8% 93	44.0% \pm 6.3% 290

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (\pm %) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2013 preceded by an asterisk (*).


CURRENT CIGARETTE SMOKING — School Type, Gender, and Minority Populations

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Middle School (MS)	12.0% ± 3.7% 360	10.4% ± 3.2% 299	6.4% ± 1.7% 184	6.4% ± 1.6% 183	7.2% ± 2.6% 212	*4.3% ± 1.4% 126
MS Females	12.9% ± 4.7% 173	11.0% ± 4.4% 153	5.5% ± 2.4% 77	5.8% ± 2.0% 79	5.7% ± 2.5% 86	*3.8% ± 2.0% 54
MS Males	11.3% ± 4.1% 184	9.8% ± 3.6% 147	7.2% ± 2.7% 108	7.0% ± 2.7% 104	8.7% ± 3.4% 125	*4.7% ± 2.0% 69
MS Minorities	16.8% ± 5.4% 183	8.6% ± 3.0% 116	6.8% ± 2.4% 97	6.6% ± 2.4% 91	8.8% ± 3.5% 130	*5.7% ± 2.4% 79
High School (HS)	31.3% ± 3.9% 1,090	25.6% ± 4.4% 914	16.2% ± 1.9% 629	17.8% ± 1.9% 682	21.6% ± 2.6% 785	*16.9% ± 2.2% 535
HS Females	32.4% ± 5.1% 572	24.4% ± 5.7% 418	16.5% ± 2.5% 327	16.7% ± 2.5% 322	19.3% ± 3.3% 369	*14.7% ± 2.7% 240
HS Males	30.1% ± 5.0% 512	26.1% ± 4.9% 477	15.9% ± 2.6% 302	18.9% ± 2.6% 359	24.2% ± 3.0% 414	*18.9% ± 3.5% 287
HS Minorities	29.6% ± 4.6% 332	21.2% ± 5.8% 279	13.7% ± 2.8% 234	13.9% ± 2.4% 221	20.5% ± 2.7% 352	*14.7% ± 2.7% 215

Wicomico County—Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)





Worcester County

Tobacco Use by Youth: 2000–2013

Maryland Department of Health and Mental Hygiene
Center for Tobacco Prevention and Control

CURRENT USE OF TOBACCO PRODUCTS

Underage (<18) Public Middle and High School Youth (Combined)
Proportion (%) and Estimated Number of Youth

TOBACCO PRODUCTS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Any Tobacco Product (All Youth Populations)	25.2% \pm 3.5% 901	23.0% \pm 3.1% 849	19.3% \pm 2.4% 729	22.5% \pm 1.9% 784	23.9% \pm 2.5% 840	20.4% \pm 3.1% 594
Any Tobacco Product (Minority Youth)	23.1% \pm 4.5% 238	30.1% \pm 6.0% 339	19.7% \pm 3.1% 240	24.6% \pm 3.2% 258	26.2% \pm 3.1% 310	20.4% \pm 3.7% 177
Cigarette	19.9% \pm 3.1% 683	16.3% \pm 3.0% 575	13.7% \pm 2.0% 505	15.6% \pm 1.6% 533	16.1% \pm 2.1% 551	14.8% \pm 2.5% 406
Cigar	10.2% \pm 2.1% 356	11.0% \pm 2.2% 395	8.9% \pm 1.4% 328	13.9% \pm 1.4% 484	12.8% \pm 1.9% 449	10.6% \pm 2.0% 305
Pipe (tobacco)	4.7% \pm 1.2% 164	6.4% \pm 1.6% 228	4.4% \pm 0.8% 166	5.4% \pm 1.1% 186	PIPE SPECIFIC DATA NOT AVAILABLE FOR 2010	PIPE SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Bidi	5.2% \pm 1.3% 180	5.6% \pm 1.9% 201	3.9% \pm 0.8% 145	3.7% \pm 1.0% 129	BIDI SPECIFIC DATA NOT AVAILABLE FOR 2010	BIDI SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Kretek	4.1% \pm 1.2% 143	5.4% \pm 2.3% 193	4.0% \pm 0.8% 148	3.4% \pm 0.8% 118	KRETEK SPECIFIC DATA NOT AVAILA- BLE FOR 2010	KRETEK SPECIFIC DATA NOT AVAIL- ABLE FOR 2013
Smokeless Tobacco	3.7% \pm 1.1% 131	5.0% \pm 1.5% 176	4.3% \pm 0.9% 161	5.3% \pm 0.9% 181	6.7% \pm 1.2% 229	*8.8% \pm 2.0% 254

INITIATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

INITIATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	21.4% \pm 2.7% 765	19.2% \pm 2.5% 710	19.7% \pm 2.4% 743	22.5% \pm 1.9% 785	23.3% \pm 2.2% 817	19.7% \pm 2.9% 547
Minority Youth	17.6% \pm 3.7% 181	18.8% \pm 4.3% 211	17.3% \pm 3.0% 212	19.8% \pm 2.9% 207	24.6% \pm 2.9% 291	20.1% \pm 3.5% 161

CESSATION OF TOBACCO USE IN PAST YEAR

Underage (<18) Public Middle and High School Youth Combined, Proportion (%) and Estimated Number of Youth

CESSATION Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
All Youth	34.3% \pm 5.7% 370	35.0% \pm 8.2% 284	40.6% \pm 5.9% 240	36.7% \pm 5.9% 248	38.1% \pm 6.4% 149	36.4% \pm 5.1% 273
Minority Youth	42.7% \pm 14.3% 43	28.9% \pm 11.8% 28	38.2% \pm 13.0% 30	35.2% \pm 9.6% 38	43.7% \pm 9.3% 53	34.8% \pm 7.4% 77

- Estimates of prevalence are stated as a percentage (%) of the total relevant population.
- Confidence Intervals (\pm %) appear immediately following prevalence estimates.
- Statistically significant change from the previous survey year are underlined; change between 2000 and 2013 preceded by an asterisk (*).

CURRENT CIGARETTE SMOKING — School Type, Gender, and Minority Populations

Underage (<18) Public Middle and High School Youth
Proportion (%) and Estimated Number of Youth

YOUTH POPULATIONS Prevalence (%), CI, and Number Youth	2000 Baseline Data	2002	2006	2008	2010	2013
Middle School (MS)	9.0% $\pm 3.2\%$ 133	6.4% $\pm 1.4\%$ 100	5.0% $\pm 1.4\%$ 71	4.2% $\pm 1.2\%$ 58	5.8% $\pm 1.8\%$ 82	6.5% $\pm 3.5\%$ 71
MS Females	5.7% $\pm 2.9\%$ 40	6.0% $\pm 1.9\%$ 47	5.3% $\pm 1.7\%$ 38	3.4% $\pm 1.4\%$ 23	6.6% $\pm 2.3\%$ 48	8.4% $\pm 5.3\%$ 46
MS Males	12.2% $\pm 4.4\%$ 93	6.9% $\pm 2.2\%$ 53	4.7% $\pm 1.7\%$ 33	4.9% $\pm 1.8\%$ 34	4.7% $\pm 2.0\%$ 33	*4.4% $\pm 2.7\%$ 24
MS Minorities	8.1% $\pm 4.1\%$ 32	7.8% $\pm 2.3\%$ 36	6.3% $\pm 2.5\%$ 30	3.9% $\pm 2.0\%$ 16	7.6% $\pm 2.9\%$ 39	5.5% $\pm 2.9\%$ 19
High School (HS)	28.0% $\pm 3.9\%$ 550	24.0% $\pm 4.6\%$ 475	19.2% $\pm 2.4\%$ 433	23.3% $\pm 2.5\%$ 475	23.4% $\pm 3.1\%$ 468	*20.2% $\pm 3.1\%$ 335
HS Females	24.6% $\pm 5.2\%$ 229	23.0% $\pm 6.2\%$ 219	18.8% $\pm 2.7\%$ 208	20.3% $\pm 3.2\%$ 204	19.6% $\pm 3.4\%$ 205	19.0% $\pm 4.1\%$ 158
HS Males	31.1% $\pm 4.8\%$ 320	24.5% $\pm 5.6\%$ 249	19.6% $\pm 2.6\%$ 226	26.3% $\pm 3.3\%$ 271	27.4% $\pm 5.3\%$ 264	*21.0% $\pm 3.9\%$ 173
HS Minorities	20.5% $\pm 5.9\%$ 118	28.3% $\pm 8.1\%$ 170	16.9% $\pm 3.9\%$ 117	21.1% $\pm 3.8\%$ 125	21.5% $\pm 3.6\%$ 135	19.7% $\pm 4.1\%$ 90

Worcester County—Current Cigarette Smoking

(Underage Middle & High School Youth, by Grade)




Table of Contents

APPENDIX B—ADULT PREVALENCE, INITIATION, AND CESSATION

	<u>Pages</u>
Maryland	93
Allegany County	95
Anne Arundel County	97
Baltimore City	99
Baltimore County	101
Calvert County	103
Caroline County	105
Carroll County	107
Cecil County	109
Charles County	111
Dorchester County	113
Frederick County	115
Garrett County	117
Harford County	119
Howard County	121
Kent County	123
Montgomery County	125
Prince George’s County	127
Queen Anne’s County	129
Somerset County	131
St. Mary’s County	133
Talbot County	135
Washington County	137
Wicomico County	139
Worcester County	141

APPENDIX B—ADULT PREVALENCE, INITIATION, AND CESSATION

Statewide and Jurisdiction Data

Data for the State of Maryland's 24 major political subdivisions (23 counties and Baltimore City) are presented in detailed tables in this appendix. The jurisdictions are arranged in alphabetical order following the State of Maryland. For each jurisdiction, the data are presented in a two-page format intended to be printed double-sided with the jurisdiction name appearing at the top of the first page.

The following information is presented to help readers understand the data and any limitations or cautions regarding the data. Because the Behavioral Risk Factor Surveillance System (BRFSS) is a CDC instrument designed to produce statewide data, sample sizes for jurisdictions may be relatively small and cause the data to be suppressed if fewer than 50 people in a jurisdiction answered the question(s) underlying the data presented or if the relative standard error (standard error divided by the estimate) of the data was too great.

Historical statewide and jurisdiction estimates in this report may vary from those previously published. Recent changes in methodology used by the Maryland BRFSS website to estimate risk behavior may cause slight differences from previously published historical data. All data in this report is calculated using the revised methodology.

<u>Term Used</u>	<u>Meaning</u>
<i>BRFSS Data Not Available</i>	The BRFSS survey did not ask questions needed to report the data.
<i>BRFSS Data are Suppressed Due to Small Sample</i>	Less than 50 people in the jurisdiction's survey sample answered the question.
<i>BRFSS Data are Suppressed Due to Imprecise and Unreliable Estimates</i>	Although 50 or more people answered the question, the relative standard error was high enough to make the estimate so unreliable and imprecise that it had to be suppressed.

Reading the Data Block

19.1%	←	The weighted estimated proportion , i.e., the estimated percentage of smokers
11.4% - 26.8%	←	The weighted confidence interval , i.e., how precise the estimate is
3,076	←	The weighted estimated number , i.e. the estimated number of smokers

NOTE: If the weighted estimated proportion is preceded by a red asterisk ***19.1%**, then the relative standard error is sufficiently large that the data should be used with caution as it may be unreliable.

If the confidence interval is in black as in the example above, it is no greater than $\pm 7\%$.

If the confidence interval is in blue, then the interval is greater than $\pm 7\%$ but no more than $\pm 10\%$.

If the confidence interval is in red, then the interval is greater than $\pm 10\%$.



Maryland

Tobacco Use by Adults: 2000–2012

Maryland Department of Health and Mental Hygiene—Center for Tobacco Prevention and Control
(See Appendix—Statewide and Jurisdiction Data for information on how to use and interpret these tables)

Historical jurisdiction estimates in this report may vary from those previously published. A methodology change in 2011 was designed to improve estimates of risk behaviors. Previous BRFSS estimates cannot be compared to the 2011 (and thereafter) data because of the significant differences in weighting methodology.

CURRENT USE OF ANY TOBACCO PRODUCT

Adult (Age 18 and older) Estimated Current Use of One or More Tobacco Products (Estimated # Adults Currently Using Any Tobacco Product / # All Adults)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
All Adults	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		19.4% 18.2% - 20.7% 856,080

CURRENT USE OF SPECIFIC TOBACCO PRODUCTS

Adult (Age 18 and older) Estimated Current Use of Specific Tobacco Products (Estimated # Currently Using Specified Tobacco Product / # All Adults)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Cigarettes	20.5% 19.1% - 21.9% 799,273	21.9% 20.4% - 23.4% 879,623	19.5% 18.1% - 20.9% 808,759	17.7% 16.7% - 18.7% 746,347	14.9% 14.0% - 15.8% 641,382	15.2% 14.3% - 16.1% 654,019		16.2% 15.0% - 17.4% 708,885
Cigars	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		4.4% 3.6% - 5.2% 169,763
Smokeless Tobacco	1.4% 0.9% - 1.9% 55,199	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	2.1% 1.6% - 2.6% 89,545		2.0% 1.5% - 2.4% 14,164
All Other Tobacco Products	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		1.7% 1.2% - 2.2% 64,984

INITIATION AND CESSATION OF CIGARETTE SMOKING IN PAST YEAR

INITIATION—Percent of All Maryland Adults Who Report First Smoking a Cigarette During the Past Year (Est. # First Tried Cigarette Smoking Past Year / # All Adults)

CESSATION—Percent of All Maryland Adults Who Report Quitting Cigarette Smoking During the Past Year (Est. # Quit Smoking Cigarettes Past Year / # All Adults)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Initiation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		0.5% 0.2% - 0.8% 22,635
Cessation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		3.1% 2.6% - 3.7% 141,391

PREGNANT WOMEN — Cigarette Smoking At Any Time During Pregnancy

Maryland Residents Regardless of Where Delivery Occurred — Birth Certificate Data, 2012 — Percent = Number Reported Smoking / Total Number

Percentage Smoking (%) Number Smoking (#)	2000	2002	2004	2006	2008	2010	2012
Pregnant Women	9.2% 6,842	8.0% 5,877	7.4% 5,504	6.8% 5,261	6.6% 5,105	6.1% 4,584	7.4% 5,351

SELECTED POPULATIONS — Current Cigarette Smoking


(Estimated # In Selected Population Currently Smoking Cigarettes / # All Adults In the Selected Population)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Non-Hispanic Black	21.7% 18.5% - 25.0% 212,976	23.8% 20.1% - 27.5% 246,836	19.9% 16.4% - 23.4% 214,045	19.8% 17.1% - 22.5% 214,207	16.4% 14.1% - 18.7% 188,180	17.8% 15.6% - 20.0% 216,031		16.7% 14.2% - 19.2% 213,250
Non-Hispanic White	21.0% 19.3% - 22.7% 528,299	21.4% 19.7% - 23.1% 535,514	19.8% 18.1% - 21.5% 507,906	17.9% 16.8% - 19.0% 447,890	15.8% 14.8% - 16.8% 396,877	15.0% 14.0% - 16.1% 372,750		17.6% 16.1% - 19.1% 424,148
Hispanic	17.2% 10.0% - 24.4% 31,887	17.8% 8.8% - 26.8% 34,407	16.7% 8.4% - 25.0% 28,594	11.1% 5.6% - 16.6% 22,096	7.9% 3.9% - 11.9% 21,812	7.8% 3.7% - 11.9% 213,250		10.1% 5.2% - 14.9% 213,250
Female	19.1% 17.3% - 20.9% 387,768	18.7% 16.9% - 20.5% 396,305	16.8% 15.1% - 18.5% 364,906	16.5% 15.3% - 17.7% 364,354	13.8% 12.7% - 14.9% 312,743	13.4% 12.3% - 14.5% 301,289		14.9% 13.5% - 16.4% 343,167
Male	22.0% 19.6% - 24.24% 411,225	25.6% 23.1% - 28.1% 485,016	22.6% 20.2% - 25.0% 443,854	19.0% 17.4% - 20.6% 380,795	16.1% 14.6% - 17.6% 329,326	17.3% 15.8% - 18.9% 354,297		17.5% 15.7% - 19.4% 365,718
Highest Education: <= High School/GED	30.1% 27.2% - 33.0% 425,204	31.1% 27.9% - 34.3% 453,357	31.1% 27.6% - 34.7% 447,456	26.4% 23.9% - 29.0% 355,662	23.2% 20.8% - 25.6% 331,037	25.3% 22.4% - 28.2% 351,607		24.6% 22.2% - 27.0% 410,780
Highest Education: >= 4 Years of College	10.6% 8.8% - 12.4% 158,061	11.0% 9.2% - 12.8% 177,491	9.3% 7.7% - 10.9% 160,985	9.5% 8.4% - 10.7% 177,136	7.0% 6.0% - 8.0% 130,093	6.5% 5.5% - 7.5% 118,002		6.2% 5.1% - 7.3% 91,503
Household Income: < \$50,000 Annually	26.6% 24.1% - 29.0% 443,966	29.1% 26.1% - 32.1% 480,711	28.2% 25.0% - 31.4% 439,672	25.6% 23.2% - 28.0% 347,080	21.4% 19.2% - 23.5% 298,195	23.2% 20.6% - 25.9% 314,619		23.0% 20.7% - 25.4% 383,703
Household Income: >= \$50,000 Annually	16.2% 14.2% - 18.2% 270,909	16.5% 14.5% - 18.5% 297,085	14.0% 12.1% - 15.9% 287,878	14.3% 12.8% - 15.8% 319,529	11.3% 10.1% - 12.6% 269,778	10.7% 9.1% - 12.2% 240,376		11.6% 10.3% - 12.9% 251,936

% OF HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS

(Estimated # of Households with Resident Minor Children and a Resident Adult Smoker / # Households with Resident Minor Children)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Proportion with Adult Smokers	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		26.5% 24.8% - 28.2% 213,250



Allegany County

Tobacco Use by Adults: 2000–2012

Maryland Department of Health and Mental Hygiene—Center for Tobacco Prevention and Control
(See Appendix—Statewide and Jurisdiction Data for information on how to use and interpret these tables)

Historical jurisdiction estimates in this report may vary from those previously published. A methodology change in 2011 was designed to improve estimates of risk behaviors. Previous BRFSS estimates cannot be compared to the 2011 (and thereafter) data because of the significant differences in weighting methodology.

CURRENT USE OF ANY TOBACCO PRODUCT								
Adult (Ages 18 and older) Estimated Current Use of One or More Tobacco Products (Estimated # Adults Currently Using Any Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
All Adults	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		28.3% 17.9% - 38.7% 17,530
CURRENT USE OF SPECIFIC TOBACCO PRODUCTS								
Adult (Age 18 and older) Estimated Current Use of Specific Tobacco Products (Estimated # Currently Using Specified Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Cigarettes	25.8% 16.9% - 34.7 13,287	23.8% 15.6% - 32.1% 15,237	23.9% 14.7% - 33.0% 13,158	27.3% 18.1% - 36.5% 16,729	23.3% 16.3% - 30.2% 14,606	11.9 14.8% - 29.7% 14,066		24.4% 14.0% - 34.8% 14,981
Cigars	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE
Smokeless Tobacco	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	*5.9% 1.1% - 10.8% 4,028		*6.9% 1.8% - 12.0% 4,420
All Other Tobacco Products	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*0.3% 0.0% - 0.7% 165
INITIATION AND CESSATION OF CIGARETTE SMOKING IN PAST YEAR								
INITIATION—Percent of All Maryland Adults Who Report First Smoking a Cigarette During the Past Year (Est. # First Tried Cigarette Smoking Past Year / # All Adults) CESSATION—Percent of All Maryland Adults Who Report Quitting Cigarette Smoking During the Past Year (Est. # Quit Smoking Cigarettes Past Year / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Initiation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE
Cessation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE

PREGNANT WOMEN — Cigarette Smoking At Any Time During Pregnancy

Maryland Residents Regardless of Where Delivery Occurred— **Birth Certificate Data, 2012**—Percent = Number Reported Smoking / Total Number

Percentage Smoking (%) Number Smoking (#)	2000	2002	2004	2006	2008	2010	2012
Pregnant Women	16.8% 133	22.1% 157	24.8% 159	23.7% 166	24.7% 176	27.6% 203	26.7% 186

SELECTED POPULATIONS — Current Cigarette Smoking


(Estimated # In Selected Population Currently Smoking Cigarettes / # All Adults In the Selected Population)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Non-Hispanic Black	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE
Non-Hispanic White	22.7% 14.2% - 31.2% 11,229	24.0% 15.5% - 32.4% 13,159	24.8% 15.4% - 34.2% 13,158	28.8% 19.0% - 38.5% 16,049	21.7% 15.3% - 28.1% 12,335	22.1% 12.5% - 31.8% 13,117	25.2% 14.3% - 36.1% 13,591
Hispanic	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE
Female	23.3% 12.8% - 33.7% 6,809	30.7% 19.2% - 41.8% 10,167	28.7% 16.0% - 41.4% 8,421	22.1% 12.1% - 32.2% 7,482	21.7% 14.0% - 29.4% 7,303	24.1% 11.3% - 37.0% 9,538	26.3% 12.7% - 40.0% 7,716
Male	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	16.4% 4.7% - 28.2% 5,070	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	33.7% 17.6% - 49.9% 9,246	25.1% 13.0% - 37.1% 7,304	16.0% 5.0% - 27.1% 4,527	22.7% 6.8% - 38.5% 7,265
Highest Education: ≤ High School/GED	20.6% 16.7% - 40.4% 8,146	28.4% 17.0% - 39.8% 8,127	30.3% 15.8% - 44.8% 8,219	35.9% 22.1% - 49.8% 11,573	30.8% 19.4% - 42.2% 9,224	24.4% 12.3% - 36.5% 7,350	18.4% 8.7% - 28.2% 6,010
Highest Education: ≥ 4 Years of College	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	*20.8% 4.5% - 37.1% 2,671	*6.0% 0.4% - 11.7% 776	BRFSS DATA ARE SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES	*3.5% 0.5% - 6.6% 383
Household Income: < \$50,000 Annually	38.9% 26.2% - 51.7% 11,512	30.1% 19.1% - 41.2% 10,401	28.2% 15.1% - 41.2% 8,663	32.0% 18.1% - 45.9% 10,956	28.0% 17.9% - 38.1% 10,001	25.2% 13.1% - 37.3% 8,350	30.2% 16.6% - 44.7% 10,369
Household Income: ≥ \$50,000 Annually	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	*22.2% 8.6% - 35.7% 3,911	*16.6% 6.3% - 26.9% 3,205	BRFSS DATA ARE SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES	*8.7% 0.2% - 17.1% 1,511

% OF HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS

(Estimated # of Households with Resident Minor Children and a Resident Adult Smoker / # Households with Resident Minor Children)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Proportion with Adult Smokers	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	31.5% 14.2% - 48.8% 4,496



Anne Arundel County

Tobacco Use by Adults: 2000–2012

Maryland Department of Health and Mental Hygiene—Center for Tobacco Prevention and Control
(See Appendix—Statewide and Jurisdiction Data for information on how to use and interpret these tables)

Historical jurisdiction estimates in this report may vary from those previously published. A methodology change in 2011 was designed to improve estimates of risk behaviors. Previous BRFSS estimates cannot be compared to the 2011 (and thereafter) data because of the significant differences in weighting methodology.

CURRENT USE OF ANY TOBACCO PRODUCT								
Adult (Age 18 and older) Estimated Current Use of One or More Tobacco Products (Estimated # Adults Currently Using Any Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
All Adults	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		21.5% 17.3% - 25.7% 81,624
CURRENT USE OF SPECIFIC TOBACCO PRODUCTS								
Adult (Age 18 and older) Estimated Current Use of Specific Tobacco Products (Estimated # Currently Using Specified Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Cigarettes	24.8% 19.2% - 30.4% 88,630	23.7% 18.1% - 29.3% 90,944	20.1% 15.0% - 25.2% 80,514	17.0% 13.1% - 20.9% 61,288	13.7% 9.6% - 17.8% 52,114	15.3% 11.0% - 19.7% 57,566		18.1% 14.2% - 22.1% 68,450
Cigars	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		4.5% 2.2% - 6.7% 15,138
Smokeless Tobacco	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	*1.7% 0.3% - 3.0% 6,243		*1.5% 0.6% - 2.5% 5,923
All Other Tobacco Products	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*1.8% 0.3% - 3.3% 6,211
INITIATION AND CESSATION OF CIGARETTE SMOKING IN PAST YEAR								
INITIATION—Percent of All Maryland Adults Who Report First Smoking a Cigarette During the Past Year (Est. # First Tried Cigarette Smoking Past Year / # All Adults) CESSATION—Percent of All Maryland Adults Who Report Quitting Cigarette Smoking During the Past Year (Est. # Quit Smoking Cigarettes Past Year / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Initiation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA ARE SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES
Cessation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		6.6% 4.0% - 9.2% 25,576

PREGNANT WOMEN — Cigarette Smoking At Any Time During Pregnancy

Maryland Residents Regardless of Where Delivery Occurred— Birth Certificate Data, 2012—Percent = Number Reported Smoking / Total Number

Percentage Smoking (%) Number Smoking (#)	2000	2002	2004	2006	2008	2010	2012
Pregnant Women	11.1% 754	9.7% 656	8.7% 587	8.1% 576	7.7% 552	8.0% 566	7.5% 512

SELECTED POPULATIONS — Current Cigarette Smoking


(Estimated # In Selected Population Currently Smoking Cigarettes / # All Adults In the Selected Population)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Non-Hispanic Black	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	*16.1% 5.7% - 26.5% 7,123	*10.0% 1.9% - 18.0% 5,295	*22.6% 5.2% - 40.0% 12,488	*12.6% 3.2% - 22.1% 6,258
Non-Hispanic White	27.1% 20.5% - 33.6% 76,856	22.1% 16.5% - 27.7% 68,166	20.8% 15.1% - 26.5% 64,533	17.2% 13.0% - 21.5% 46,264	15.1% 10.2% - 20.1% 43,034	15.2% 10.8% - 19.6% 42,370	18.8% 14.3% - 23.2% 52,986
Hispanic	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE
Female	26.5% 19.9% - 34.0% 42,735	23.7% 16.8% - 30.5% 45,430	21.7% 14.8% - 28.5% 46,304	16.6% 12.2% - 21.1% 33,074	13.2% 7.2% - 19.2% 27,205	16.0% 10.6% - 21.5% 30,488	18.5% 13.1% - 23.9% 38,547
Male	23.4% 15.4% - 31.4% 45,895	23.7% 14.9% - 32.6% 45,515	18.3% 10.7% - 26.0% 34,209	17.4% 10.8% - 24.1% 28,214	14.3% 9.0% - 19.6% 24,909	14.6% 7.7% - 21.5% 27,078	17.7% 11.9% - 23.6% 29,903
Highest Education: <= High School/GED	33.0% 22.8% - 43.2% 39,652	35.1% 25.0% - 45.1% 50,472	32.0% 21.0% - 42.9% 41,325	28.2% 18.9% - 37.5% 26,546	19.0% 9.8% - 28.3% 23,487	26.6% 16.5% - 36.6% 38,542	24.4% 17.7% - 31.0% 35,681
Highest Education: >= 4 Years of College	16.9% 9.2% - 24.7% 24,792	11.4% 5.6% - 17.3% 16,730	9.1% 3.9% - 14.2% 14,930	8.6% 4.9% - 12.2% 13,295	10.1% 5.9% - 14.3% 16,144	5.9% 3.0% - 8.8% 10,262	4.3% 2.3% - 6.3% 5,050
Household Income: < \$50,000 Annually	30.9% 21.5% - 40.4% 43,039	33.8% 23.2% - 44.3% 43,513	38.9% 27.1% - 50.7% 47,242	19.9% 12.2% - 27.5% 18,102	18.6% 10.6% - 26.6% 17,978	32.2% 20.7% - 43.8% 31,750	22.7% 14.8% - 30.6% 27,793
Household Income: >= \$50,000 Annually	21.2% 13.8% - 28.6% 31,305	15.2% 9.3% - 21.1% 27,834	12.1% 7.4% - 16.8% 27,748	15.0% 10.3% - 19.8% 33,418	11.1% 7.4% - 14.8% 26,435	8.1% 4.7% - 11.4% 17,966	14.7% 10.1% - 19.3% 30,114

% OF HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS

(Estimated # of Households with Resident Minor Children and a Resident Adult Smoker / # Households with Resident Minor Children)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Proportion with Adult Smokers	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	30.7% 23.1% - 38.4% 48,196



Baltimore City

Tobacco Use by Adults: 2000–2012

Maryland Department of Health and Mental Hygiene—Center for Tobacco Prevention and Control
(See Appendix—Statewide and Jurisdiction Data for information on how to use and interpret these tables)

Historical jurisdiction estimates in this report may vary from those previously published. A methodology change in 2011 was designed to improve estimates of risk behaviors. Previous BRFSS estimates cannot be compared to the 2011 (and thereafter) data because of the significant differences in weighting methodology.

CURRENT USE OF ANY TOBACCO PRODUCT

Adult (Age 18 and older) Estimated Current Use of One or More Tobacco Products (Estimated # Adults Currently Using Any Tobacco Product / # All Adults)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
All Adults	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		24.2% 19.1% - 29.3% 96,402

CURRENT USE OF SPECIFIC TOBACCO PRODUCTS

Adult (Age 18 and older) Estimated Current Use of Specific Tobacco Products (Estimated # Currently Using Specified Tobacco Product / # All Adults)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Cigarettes	29.0% 23.2% - 34.7% 126,781	26.0% 19.3% - 32.7% 123,614	28.3% 21.0% - 35.6% 137,459	27.0% 21.6% - 32.4% 112,289	21.9% 17.6% - 26.2% 88,599	25.5% 19.6% - 31.3% 93,651		21.2% 16.4% - 26.0% 84,398
Cigars	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		5.7% 2.5% - 8.8% 19,716
Smokeless Tobacco	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	*2.7% 1.0% - 4.3% 9,855		*2.5% 0.3% - 4.7% 10,102
All Other Tobacco Products	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*2.2% 0.0% - 4.7% 7,658

INITIATION AND CESSATION OF CIGARETTE SMOKING IN PAST YEAR

INITIATION—Percent of All Maryland Adults Who Report First Smoking a Cigarette During the Past Year (Est. # First Tried Cigarette Smoking Past Year / # All Adults)

CESSATION—Percent of All Maryland Adults Who Report Quitting Cigarette Smoking During the Past Year (Est. # Quit Smoking Cigarettes Past Year / # All Adults)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Initiation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*1.3% 0.0% - 3.1% 5,282
Cessation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*3.7% 1.2% - 6.1% 14,892

PREGNANT WOMEN — Cigarette Smoking At Any Time During Pregnancy

Maryland Residents Regardless of Where Delivery Occurred — Birth Certificate Data, 2012 — Percent = Number Reported Smoking / Total Number

Percentage Smoking (%) Number Smoking (#)	2000	2002	2004	2006	2008	2010	2012
Pregnant Women	14.9% 1,435	13.9% 1,254	12.0% 1,106	10.3% 1,009	10.3% 1,025	11.6% 1,036	10.4% 945

SELECTED POPULATIONS — Current Cigarette Smoking


(Estimated # In Selected Population Currently Smoking Cigarettes / # All Adults In the Selected Population)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Non-Hispanic Black	30.0% 21.6% - 38.3% 77,550	28.4% 18.1% - 38.8% 76,345	30.2% 19.0% - 41.3% 79,500	31.6% 23.9% - 39.2% 82,722	25.5% 19.4% - 31.7% 61,702	30.8% 22.7% - 39.0% 68,460	25.2% 18.2% - 32.2% 58,845
Non-Hispanic White	32.9% 24.6% - 41.2% 47,377	22.9% 15.0% - 30.9% 36,395	26.8% 16.8% - 36.8% 48,196	18.4% 12.2% - 24.5% 22,117	16.9% 10.9% - 22.8% 22,861	16.6% 8.9% - 24.3% 19,471	18.8% 11.5% - 26.0% 22,084
Hispanic	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE
Female	26.0% 19.5% - 32.6% 63,542	22.2% 15.0% - 29.5% 58,671	18.2% 12.4% - 24.1% 48,263	24.0% 18.1% - 29.8% 55,567	20.6% 16.1% - 25.2% 48,615	21.0% 15.3% - 26.7% 47,920	16.3% 11.6% - 21.0% 36,769
Male	32.7% 22.5% - 42.8% 63,239	30.7% 18.9% - 42.5% 64,943	40.4% 26.9% - 53.9% 89,196	30.8% 21.2% - 40.3% 56,722	23.8% 15.7% - 31.8% 39,984	32.8% 21.2% - 44.3% 45,731	27.6% 18.5% - 36.7% 47,629
Highest Education: ≤ High School/GED	34.0% 25.0% - 42.9% 74,916	33.5% 22.4% - 44.5% 80,485	43.0% 31.0% - 54.9% 104,871	32.1% 23.1% - 41.1% 61,282	31.6% 23.9% - 39.3% 57,257	37.7% 28.1% - 47.3% 64,824	30.5% 22.0% - 39.0% 56,042
Highest Education: ≥ 4 Years of College	16.6% 7.7% - 25.5% 18,075	*9.6% 3.6% - 15.5% 12,945	*9.8% 3.8% - 15.7% 15,883	16.3% 9.4% - 23.2% 20,252	8.7% 4.6% - 12.8% 11,416	*9.4% 2.7% - 16.0% 10,518	11.4% 4.8% - 17.9% 11,735
Household Income: < \$50,000 Annually	33.4% 25.9% - 41.0% 95,012	32.3% 23.8% - 40.9% 96,464	35.3% 25.3% - 45.4% 108,340	31.4% 24.0% - 38.8% 69,680	27.6% 21.2% - 34.0% 60,559	31.2% 22.6% - 39.7% 61,211	27.0% 19.6% - 34.5% 60,119
Household Income: ≥ \$50,000 Annually	20.9% 9.9% - 32.0% 17,284	*10.6% 2.9% - 18.2% 10,897	16.3% 7.0% - 25.5% 20,785	24.5% 14.2% - 34.9% 31,218	14.0% 8.3% - 19.8% 19,679	*12.8% 5.2% - 20.5% 14,454	14.8% 8.0% - 21.7% 17,851

% OF HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS

(Estimated # of Households with Resident Minor Children and a Resident Adult Smoker / # Households with Resident Minor Children)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Proportion with Adult Smokers	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	31.6% 21.3% - 41.8% 36,345



Baltimore County

Tobacco Use by Adults: 2000–2012

Maryland Department of Health and Mental Hygiene—Center for Tobacco Prevention and Control
(See Appendix—Statewide and Jurisdiction Data for information on how to use and interpret these tables)

Historical jurisdiction estimates in this report may vary from those previously published. A methodology change in 2011 was designed to improve estimates of risk behaviors. Previous BRFSS estimates cannot be compared to the 2011 (and thereafter) data because of the significant differences in weighting methodology.

CURRENT USE OF ANY TOBACCO PRODUCT								
Adult (Age 18 and older) Estimated Current Use of One or More Tobacco Products (Estimated # Adults Currently Using Any Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
All Adults	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		22.8% 19.2% - 26.3% 166,242
CURRENT USE OF SPECIFIC TOBACCO PRODUCTS								
Adult (Age 18 and older) Estimated Current Use of Specific Tobacco Products (Estimated # Currently Using Specified Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Cigarettes	22.6% 18.3% - 26.8% 135,002	24.4% 19.8% - 28.9% 142,386	22.9% 18.3% - 27.3% 133,225	20.1% 16.6% - 23.6% 122,230	16.9% 13.8% - 19.9% 111,038	15.6% 11.9% - 19.3% 103,135		19.0% 15.7% - 22.4% 138,639
Cigars	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		4.9% 2.8% - 6.9% 32,194
Smokeless Tobacco	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	*2.4% 0.7% - 4.1% 15,732		*1.8% 0.6% - 3.0% 13,260
All Other Tobacco Products	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*1.8% 0.6% - 3.0% 11,754
INITIATION AND CESSATION OF CIGARETTE SMOKING IN PAST YEAR								
INITIATION—Percent of All Maryland Adults Who Report First Smoking a Cigarette During the Past Year (Est. # First Tried Cigarette Smoking Past Year / # All Adults) CESSATION—Percent of All Maryland Adults Who Report Quitting Cigarette Smoking During the Past Year (Est. # Quit Smoking Cigarettes Past Year / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Initiation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*0.7% 0.0% - 1.6% 22,635
Cessation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		3.1% 2.0% - 4.1% 22,991

PREGNANT WOMEN — Cigarette Smoking At Any Time During Pregnancy

Maryland Residents Regardless of Where Delivery Occurred— Birth Certificate Data, 2012—Percent = Number Reported Smoking / Total Number

Percentage Smoking (%) Number Smoking (#)	2000	2002	2004	2006	2008	2010	2012
Pregnant Women	9.6% 904	8.8% 791	8.6% 809	7.8% 776	7.3% 748	9.0% 894	8.1% 777

SELECTED POPULATIONS — Current Cigarette Smoking

(Estimated # In Selected Population Currently Smoking Cigarettes / # All Adults In the Selected Population)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Non-Hispanic Black	25.1% 14.8% - 35.5% 32,246	23.3% 12.1% - 34.5% 30,675	23.0% 12.6% - 33.4% 35,834	16.3% 9.0% - 23.5% 27,953	17.3% 11.6% - 23.0% 33,305	19.7% 12.5% - 26.9% 33,434	16.4% 10.4% - 22.4% 39,738
Non-Hispanic White	22.6% 17.6% - 27.4% 97,526	25.7% 20.5% - 31.0% 97,979	21.9% 16.9% - 26.9% 78,790	21.3% 17.6% - 25.1% 78,034	17.6% 13.7% - 21.5% 68,431	12.4% 9.8% - 14.9% 50,223	22.7% 18.3% - 27.0% 89,914
Hispanic	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE
Female	21.1% 16.2% - 25.9% 65,684	17.3% 12.6% - 21.9% 57,101	18.6% 13.6% - 23.6% 56,225	18.6% 14.7% - 22.4% 62,478	15.2% 11.6% - 18.6% 54,932	13.2% 10.2% - 16.3% 46,029	18.5% 14.0% - 23.0% 80,467
Male	24.2% 17.1% - 31.3% 69,317	33.6% 25.4% - 41.8% 85,286	27.6% 19.7% - 35.4% 77,000	22.0% 15.8% - 28.3% 59,752	18.9% 13.5% - 24.3% 56,107	18.2% 11.4% - 25.1% 57,107	19.8% 14.8% - 24.8% 58,172
Highest Education: ≤ High School/GED	33.3% 25.5% - 41.0% 79,819	33.0% 24.3% - 41.6% 69,888	33.2% 23.7% - 42.7% 65,349	28.7% 21.5% - 36.0% 61,486	23.7% 17.8% - 29.6% 52,611	25.5% 17.1% - 33.8% 60,994	31.2% 24.6% - 37.8% 91,131
Highest Education: ≥ 4 Years of College	11.6% 6.8% - 16.5% 23,385	11.7% 6.7% - 16.7% 25,333	15.4% 9.7% - 21.1% 36,051	9.7% 6.4% - 13.0% 22,459	9.1% 5.3% - 12.8% 24,845	6.9% 4.6% - 9.3% 18,116	8.8% 5.3% - 12.4% 19,562
Household Income: < \$50,000 Annually	27.9% 21.2% - 34.5% 78,557	29.7% 22.1% - 37.2% 75,439	30.2% 22.2% - 38.2% 76,884	23.9% 18.1% - 29.7% 54,061	23.4% 17.7% - 29.2% 55,274	22.8% 17.1% - 28.5% 50,492	26.8% 20.6% - 32.9% 83,834
Household Income: ≥ \$50,000 Annually	16.9% 11.0% - 22.9% 38,578	19.8% 13.4% - 26.2% 48,894	17.7% 11.9% - 23.5% 43,545	18.7% 13.8% - 23.6% 50,851	13.3% 9.5% - 17.2% 43,806	11.3% 5.3% - 17.2% 37,792	13.6% 9.7% - 17.5% 44,923

% OF HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS

(Estimated # of Households with Resident Minor Children and a Resident Adult Smoker / # Households with Resident Minor Children)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Proportion with Adult Smokers	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	32.9% 26.1% - 39.6% 90,479



Calvert County

Tobacco Use by Adults: 2000–2012

Maryland Department of Health and Mental Hygiene—Center for Tobacco Prevention and Control
(See Appendix—Statewide and Jurisdiction Data for information on how to use and interpret these tables)

Historical jurisdiction estimates in this report may vary from those previously published. A methodology change in 2011 was designed to improve estimates of risk behaviors. Previous BRFSS estimates cannot be compared to the 2011 (and thereafter) data because of the significant differences in weighting methodology.

CURRENT USE OF ANY TOBACCO PRODUCT								
Adult (Age 18 and older) Estimated Current Use of One or More Tobacco Products (Estimated # Adults Currently Using Any Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
All Adults	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		27.2% 19.5% - 34.8% 18,800
CURRENT USE OF SPECIFIC TOBACCO PRODUCTS								
Adult (Age 18 and older) Estimated Current Use of Specific Tobacco Products (Estimated # Currently Using Specified Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Cigarettes	31.1% 20.3% - 41.9% 17,949	23.0% 15.9% - 30.9% 15,264	21.7% 12.8% - 30.6% 12,918	18.9% 12.8% - 24.9% 9,543	20.0% 13.1% - 26.8% 14,028	14.5% 7.8% - 21.3% 9,388		18.9% 13.1% - 24.6% 13,014
Cigars	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES
Smokeless Tobacco	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	*2.4% 0.0% - 5.4% 1,569		*1.1% 0.2% - 2.0% 780
All Other Tobacco Products	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES
INITIATION AND CESSATION OF CIGARETTE SMOKING IN PAST YEAR								
INITIATION—Percent of All Maryland Adults Who Report First Smoking a Cigarette During the Past Year (Est. # First Tried Cigarette Smoking Past Year / # All Adults) CESSATION—Percent of All Maryland Adults Who Report Quitting Cigarette Smoking During the Past Year (Est. # Quit Smoking Cigarettes Past Year / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Initiation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Cessation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*3.1% 0.4% - 6.6% 2,462

PREGNANT WOMEN — Cigarette Smoking At Any Time During Pregnancy

Maryland Residents Regardless of Where Delivery Occurred— Birth Certificate Data, 2012—Percent = Number Reported Smoking / Total Number

Percentage Smoking (%) Number Smoking (#)	2000	2002	2004	2006	2008	2010	2012
Pregnant Women	14.2% 145	12.7% 129	11.8% 118	12.7% 128	11.5% 110	15.3% 138	13.8% 126

SELECTED POPULATIONS — Current Cigarette Smoking

(Estimated # In Selected Population Currently Smoking Cigarettes / # All Adults In the Selected Population)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Non-Hispanic Black	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Non-Hispanic White	33.3% 21.6% - 45.0% 15,088	24.4% 15.8% - 32.9% 14,680	24.2% 14.4% - 34.0% 12,738	18.9% 12.3% - 25.5% 7,477	19.0% 11.7% - 26.2% 10,944	16.6% 8.3% - 25.0% 8,374	16.7% 11.0% - 22.5% 8,669
Hispanic	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Female	30.2% 17.2% - 43.1% 7,573	20.0% 10.6% - 29.5% 6,616	25.1% 11.7% - 38.6% 7,689	23.7% 15.1% - 32.2% 6,216	16.6% 9.5% - 23.7% 6,472	17.8% 7.6% - 28.1% 6,848	18.6% 11.0% - 26.2% 7,001
Male	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	25.9% 13.3% - 38.5% 8,648	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*13.7% 5.5% - 21.8% 3,327	24.2% 11.7% - 36.7% 7,555	*9.7% 3.7% - 15.7% 2,540	19.2% 10.6% - 27.8% 6,013
Highest Education: ≤ High School/GED	38.7% 23.2% - 54.2% 11,241	30.0% 17.1% - 42.8% 8,881	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	26.4% 13.3% - 39.5% 4,116	27.2% 14.2% - 40.1% 7,129	*14.6% 4.1% - 25.0% 2,844	23.9% 14.5% - 33.3% 7,171
Highest Education: ≥ 4 Years of College	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*9.3% 2.7% - 15.9% 1,706	*4.9% 1.0% - 8.8% 1,119	*8.6% 1.8% - 15.4% 1,869	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES
Household Income: < \$50,000 Annually	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	29.5% 15.0% - 44.0% 3,315	33.7% 16.9% - 50.5% 6,382	*20.7% 3.7% - 37.8% 3,398	22.4% 12.0% - 32.8% 4,467
Household Income: ≥ \$50,000 Annually	*19.7% 4.7% - 34.8% 5,725	20.9% 11.0% - 30.9% 8,834	21.5% 10.1% - 32.8% 8,008	15.2% 8.4% - 22.1% 5,187	14.2% 7.3% - 21.1% 6,508	11.7% 4.7% - 18.7% 4,553	18.3% 10.6% - 26.1% 8,084

% OF HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS

(Estimated # of Households with Resident Minor Children and a Resident Adult Smoker / # Households with Resident Minor Children)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Proportion with Adult Smokers	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	34.4% 21.4% - 47.4% 9,246



Caroline County

Tobacco Use by Adults: 2000–2012

Maryland Department of Health and Mental Hygiene—Center for Tobacco Prevention and Control
(See Appendix—Statewide and Jurisdiction Data for information on how to use and interpret these tables)

Historical jurisdiction estimates in this report may vary from those previously published. A methodology change in 2011 was designed to improve estimates of risk behaviors. Previous BRFSS estimates cannot be compared to the 2011 (and thereafter) data because of the significant differences in weighting methodology.

CURRENT USE OF ANY TOBACCO PRODUCT								
Adult (Age 18 and older) Estimated Current Use of One or More Tobacco Products (Estimated # Adults Currently Using Any Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
All Adults	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		27.6% 14.1% - 41.1% 7,727
CURRENT USE OF SPECIFIC TOBACCO PRODUCTS								
Adult (Age 18 and older) Estimated Current Use of Specific Tobacco Products (Estimated # Currently Using Specified Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Cigarettes	24.8% 11.4% - 38.3% 7,390	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	26.4% 11.8% - 40.9% 7,634	25.5% 15.1% - 35.9% 6,373	19.0% 11.2% - 26.8% 5,057	29.9% 17.0% - 42.7% 8,495		26.2% 12.7% - 39.6% 7,302
Cigars	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES
Smokeless Tobacco	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	*2.2% 0.0% - 4.3% 627		*1.6% 0.0% - 3.2% 452
All Other Tobacco Products	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES
INITIATION AND CESSATION OF CIGARETTE SMOKING IN PAST YEAR								
INITIATION—Percent of All Maryland Adults Who Report First Smoking a Cigarette During the Past Year (Est. # First Tried Cigarette Smoking Past Year / # All Adults) CESSATION—Percent of All Maryland Adults Who Report Quitting Cigarette Smoking During the Past Year (Est. # Quit Smoking Cigarettes Past Year / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Initiation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Cessation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*0.7% 0.5% - 1.4% 199

PREGNANT WOMEN — Cigarette Smoking At Any Time During Pregnancy

Maryland Residents Regardless of Where Delivery Occurred— **Birth Certificate Data, 2012**—Percent = Number Reported Smoking / Total Number

Percentage Smoking (%) Number Smoking (#)	2000	2002	2004	2006	2008	2010	2012
Pregnant Women	17.0% 69	15.1% 58	15.9% 74	12.8% 61	12.8% 64	17.1% 74	17.1% 63

SELECTED POPULATIONS — Current Cigarette Smoking

(Estimated # In Selected Population Currently Smoking Cigarettes / # All Adults In the Selected Population)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Non-Hispanic Black	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Non-Hispanic White	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	22.1% 14.3% - 30.0% 4,169	17.8% 9.1% - 26.6% 3,697	24.5% 15.7% - 33.3% 4,864	29.5% 13.7% - 45.4% 6,451
Hispanic	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Female	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	23.7% 13.8% - 33.7% 3,085	16.5% 8.7% - 24.2% 2,098	28.5% 15.6% - 41.3% 3,386	15.8% 7.1% - 24.6% 1,903
Male	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	27.4% 8.8% - 45.9% 3,288	*21.4% 8.4% - 34.5% 2,958	*30.9% 11.0% - 50.8% 5,109	*33.9% 10.3% - 57.6% 5,399
Highest Education: ≤ High School/GED	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	34.3% 18.4% - 50.1% 4,825	29.4% 16.3% - 42.5% 3,917	45.6% 27.3% - 64.0% 6,994	*32.6% 12.4% - 52.8% 6,022
Highest Education: ≥ 4 Years of College	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES
Household Income: < \$50,000 Annually	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	32.2% 15.3% - 49.2% 4,327	24.0% 11.2% - 36.8% 2,724	41.6% 21.3% - 61.9% 6,126	*27.7% 10.8% - 44.6% 4,538
Household Income: ≥ \$50,000 Annually	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*14.2% 4.2% - 24.2% 1,181	*15.9% 4.1% - 27.7% 1,944	*11.9% 3.2% - 20.6% 1,280	*7.5% 2.4% - 12.7% 531

% OF HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS

(Estimated # of Households with Resident Minor Children and a Resident Adult Smoker / # Households with Resident Minor Children)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Proportion with Adult Smokers	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE



Carroll County

Tobacco Use by Adults: 2000–2012

Maryland Department of Health and Mental Hygiene—Center for Tobacco Prevention and Control
(See Appendix—Statewide and Jurisdiction Data for information on how to use and interpret these tables)

Historical jurisdiction estimates in this report may vary from those previously published. A methodology change in 2011 was designed to improve estimates of risk behaviors. Previous BRFSS estimates cannot be compared to the 2011 (and thereafter) data because of the significant differences in weighting methodology.

CURRENT USE OF ANY TOBACCO PRODUCT								
Adult (Age 18 and older) Estimated Current Use of One or More Tobacco Products (Estimated # Adults Currently Using Any Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
All Adults	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		23.2% 15.6% - 30.8% 29,043
CURRENT USE OF SPECIFIC TOBACCO PRODUCTS								
Adult (Age 18 and older) Estimated Current Use of Specific Tobacco Products (Estimated # Currently Using Specified Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Cigarettes	14.6% 7.2% - 22.0% 15,323	22.5% 13.2% - 31.9% 26,267	22.0% 13.1% - 31.0% 27,587	12.8% 7.7% - 17.9% 20,539	12.0% 7.3% - 16.7% 14,424	21.9% 13.2% - 30.6% 35,549		17.9% 11.1% - 24.7% 22,302
Cigars	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*4.1% 0.1% - 8.1% 4,684
Smokeless Tobacco	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES		*1.4% 0.2% - 2.8% 1,786
All Other Tobacco Products	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES
INITIATION AND CESSATION OF CIGARETTE SMOKING IN PAST YEAR								
INITIATION—Percent of All Maryland Adults Who Report First Smoking a Cigarette During the Past Year (Est. # First Tried Cigarette Smoking Past Year / # All Adults) CESSATION—Percent of All Maryland Adults Who Report Quitting Cigarette Smoking During the Past Year (Est. # Quit Smoking Cigarettes Past Year / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Initiation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Cessation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*1.3% 0.1% - 2.5% 1,657

PREGNANT WOMEN — Cigarette Smoking At Any Time During Pregnancy

Maryland Residents Regardless of Where Delivery Occurred— Birth Certificate Data, 2012—Percent = Number Reported Smoking / Total Number

Percentage Smoking (%) Number Smoking (#)	2000	2002	2004	2006	2008	2010	2012
Pregnant Women	11.1% 211	11.2% 213	9.6% 191	10.2% 191	10.6% 186	13.1% 210	10.6% 170

SELECTED POPULATIONS — Current Cigarette Smoking

(Estimated # In Selected Population Currently Smoking Cigarettes / # All Adults In the Selected Population)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Non-Hispanic Black	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Non-Hispanic White	15.7% 7.8% - 23.6% 15,323	21.4% 12.0% - 30.9% 23,977	25.6% 15.7% - 35.5% 27,587	13.6% 7.8% - 19.4% 17,418	12.7% 7.6% - 17.8% 13,554	22.1% 12.8% - 31.4% 31,886	17.5% 10.3% - 27.7% 20,207
Hispanic	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Female	18.6% 7.6% - 29.5% 11,365	22.8% 10.2% - 35.3% 16,592	*17.5% 6.8% - 28.1% 11,416	11.7% 6.3% - 17.2% 10,385	11.9% 6.2% - 17.7% 7,888	20.4% 10.6% - 30.2% 17,137	16.7% 9.3% - 24.2% 9,920
Male	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	27.0% 12.4% - 41.6% 16,171	*14.1% 4.8% - 23.4% 10,155	*12.1% 4.3% - 19.9% 6,537	*23.5% 9.0% - 38.0% 18,412	*19.0% 7.7% - 30.2% 12,383
Highest Education: ≤ High School/GED	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	26.6% 12.4% - 40.9% 15,608	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	18.9% 9.9% - 28.0% 4,456	16.0% 7.8% - 24.2% 7,306	37.0% 21.7% - 52.3% 27,817	27.1% 12.5% - 41.7% 12,307
Highest Education: ≥ 4 Years of College	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*2.3% 0.0% - 4.8% 1,404	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES	*5.9% 0.5% - 11.3% 1,946
Household Income: < \$50,000 Annually	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	26.4% 12.9% - 39.9% 12,229	*13.7% 4.9% - 22.2% 4,349	24.3% 11.2% - 37.5% 11,280	24.3% 10.1% - 38.4% 12,013
Household Income: ≥ \$50,000 Annually	*9.9% 1.0% - 18.8% 6,332	*18.9% 6.9% - 30.9% 10,983	*18.5% 7.4% - 29.5% 12,768	*6.3% 1.8% - 10.7% 5,956	12.4% 6.0% - 18.7% 9,427	*18.3% 7.3% - 29.2% 14,366	10.9% 4.8% - 17.0% 6,574

% OF HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS

(Estimated # of Households with Resident Minor Children and a Resident Adult Smoker / # Households with Resident Minor Children)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Proportion with Adult Smokers	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	41.4% 24.4% - 58.3% 17,048



Cecil County

Tobacco Use by Adults: 2000–2012

Maryland Department of Health and Mental Hygiene—Center for Tobacco Prevention and Control
(See Appendix—Statewide and Jurisdiction Data for information on how to use and interpret these tables)

Historical jurisdiction estimates in this report may vary from those previously published. A methodology change in 2011 was designed to improve estimates of risk behaviors. Previous BRFSS estimates cannot be compared to the 2011 (and thereafter) data because of the significant differences in weighting methodology.

CURRENT USE OF ANY TOBACCO PRODUCT								
Adult (Age 18 and older) Estimated Current Use of One or More Tobacco Products (Estimated # Adults Currently Using Any Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
All Adults	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		26.1% 16.3% - 35.8% 20,515
CURRENT USE OF SPECIFIC TOBACCO PRODUCTS								
Adult (Age 18 and older) Estimated Current Use of Specific Tobacco Products (Estimated # Currently Using Specified Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Cigarettes	26.7% 18.7% - 34.7% 16,364	29.9% 20.6% - 39.3% 18,884	27.8% 18.7% - 36.9% 20,690	24.6% 18.7% - 30.5% 23,868	22.4% 13.9% - 30.9% 15,644	25.4% 18.1% - 32.6% 17,737		23.0% 13.4% - 32.5% 18,089
Cigars	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*2.9% 0.4% - 5.3% 1,577
Smokeless Tobacco	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	*2.3% 0.2% - 4.5% 1,621		*1.2% 0.3% - 5.1% 2,138
All Other Tobacco Products	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		4.3% 0.0% - 9.2% 2,323
INITIATION AND CESSATION OF CIGARETTE SMOKING IN PAST YEAR								
INITIATION—Percent of All Maryland Adults Who Report First Smoking a Cigarette During the Past Year (Est. # First Tried Cigarette Smoking Past Year / # All Adults) CESSATION—Percent of All Maryland Adults Who Report Quitting Cigarette Smoking During the Past Year (Est. # Quit Smoking Cigarettes Past Year / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Initiation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Cessation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		3.2% 0.3% - 6.0% 2,513

PREGNANT WOMEN — Cigarette Smoking At Any Time During Pregnancy

Maryland Residents Regardless of Where Delivery Occurred— **Birth Certificate Data, 2012**—Percent = Number Reported Smoking / Total Number

Percentage Smoking (%) Number Smoking (#)	2000	2002	2004	2006	2008	2010	2012
Pregnant Women	23.3% 265	18.2% 211	19.7% 239	19.7% 267	22.8% 291	21.5% 254	21.2% 246

SELECTED POPULATIONS — Current Cigarette Smoking

(Estimated # In Selected Population Currently Smoking Cigarettes / # All Adults In the Selected Population)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Non-Hispanic Black	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Non-Hispanic White	25.7% 17.5% - 33.9% 14,368	28.6% 18.9% - 38.2% 16,280	27.1% 18.0% - 36.3% 17,629	25.9% 19.7% - 32.2% 21,469	23.2% 14.1% - 32.3% 14,771	25.4% 17.5% - 33.2% 15,624	23.9% 12.9% - 34.8% 15,963
Hispanic	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Female	27.4% 17.4% - 37.4% 9,595	28.0% 16.8% - 39.3% 8,715	25.6% 13.7% - 37.5% 8,720	24.0% 15.9% - 32.0% 11,694	24.6% 12.5% - 36.7% 9,163	31.0% 20.2% - 41.9% 10,724	21.7% 10.6% - 32.7% 8,985
Male	25.8% 12.6% - 39.0% 6,770	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	29.6% 16.0% - 43.2% 11,970	25.2% 16.5% - 33.9% 12,174	19.9% 8.2% - 31.6% 6,480	19.9% 10.8% - 29.0% 7,013	*24.5% 8.1% - 40.9% 9,103
Highest Education: ≤ High School/GED	38.6% 25.7% - 51.5% 11,480	36.6% 22.1% - 51.1% 10,166	47.6% 32.9% - 62.4% 15,310	36.2% 26.1% - 46.3% 14,878	27.9% 12.5% - 43.4% 9,309	34.6% 23.3% - 45.8% 10,043	29.7% 14.2% - 45.3% 12,634
Highest Education: ≥ 4 Years of College	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*13.6% 5.1% - 21.1% 3,693	*9.9% 3.3% - 16.6% 1,965	*9.6% 2.2% - 17.0% 2,152	*13.1% 4.4% - 21.9% 1,748
Household Income: < \$50,000 Annually	38.3% 24.6% - 52.0% 11,608	41.4% 25.7% - 57.0% 11,446	35.6% 20.9% - 50.3% 13,247	36.9% 26.3% - 47.6% 12,900	32.0% 18.4% - 47.2% 7,242	31.8% 17.5% - 46.2% 7,107	*26.2% 9.5% - 42.9% 9,769
Household Income: ≥ \$50,000 Annually	*17.7% 7.0% - 28.4% 3,891	24.8% 13.3% - 36.2% 7,439	25.2% 12.9% - 37.5% 7,229	17.8% 10.2% - 25.5% 8,914	15.1% 8.0% - 22.1% 5,262	21.1% 12.1% - 30.1% 7,811	14.0% 5.8% - 22.2% 4,361

% OF HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS

(Estimated # of Households with Resident Minor Children and a Resident Adult Smoker / # Households with Resident Minor Children)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Proportion with Adult Smokers	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	31.7% 14.2% - 49.2% 10,813



Charles County Tobacco Use by Adults: 2000–2012

Maryland Department of Health and Mental Hygiene—Center for Tobacco Prevention and Control
(See Appendix—Statewide and Jurisdiction Data for information on how to use and interpret these tables)

Historical jurisdiction estimates in this report may vary from those previously published. A methodology change in 2011 was designed to improve estimates of risk behaviors. Previous BRFSS estimates cannot be compared to the 2011 (and thereafter) data because of the significant differences in weighting methodology.

CURRENT USE OF ANY TOBACCO PRODUCT

Adult (Age 18 and older) Estimated Current Use of One or More Tobacco Products (Estimated # Adults Currently Using Any Tobacco Product / # All Adults)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
All Adults	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		20.5% 13.5% - 27.4% 27,840

CURRENT USE OF SPECIFIC TOBACCO PRODUCTS

Adult (Age 18 and older) Estimated Current Use of Specific Tobacco Products (Estimated # Currently Using Specified Tobacco Product / # All Adults)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Cigarettes	26.3% 19.2% - 33.4% 21,262	24.4% 16.6% - 32.2% 24,082	17.8% 10.4% - 25.3% 16,586	18.3% 12.2% - 24.4% 19,814	18.2% 11.4% - 25.0% 18,553	14.0% 9.4% - 18.7% 16,158		19.3% 12.3% - 26.3% 15,379
Cigars	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*2.3% 0.3% - 4.4% 124
Smokeless Tobacco	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	*1.8% 0.0% - 3.9% 2,043		*0.8% 0.06% - 1.5% 1,086
All Other Tobacco Products	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*0.2% 0.0% - 0.4% 215

INITIATION AND CESSATION OF CIGARETTE SMOKING IN PAST YEAR

INITIATION—Percent of All Maryland Adults Who Report First Smoking a Cigarette During the Past Year (Est. # First Tried Cigarette Smoking Past Year / # All Adults)

CESSATION—Percent of All Maryland Adults Who Report Quitting Cigarette Smoking During the Past Year (Est. # Quit Smoking Cigarettes Past Year / # All Adults)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Initiation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Cessation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*2.5% 0.7% - 4.2% 3,392

PREGNANT WOMEN — Cigarette Smoking At Any Time During Pregnancy

Maryland Residents Regardless of Where Delivery Occurred— Birth Certificate Data, 2012—Percent = Number Reported Smoking / Total Number

Percentage Smoking (%) Number Smoking (#)	2000	2002	2004	2006	2008	2010	2012
Pregnant Women	13.3% 232	11.1% 195	10.7% 194	8.7% 169	7.0% 134	8.7% 157	8.3% 160

SELECTED POPULATIONS — Current Cigarette Smoking


(Estimated # In Selected Population Currently Smoking Cigarettes / # All Adults In the Selected Population)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Non-Hispanic Black	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*16.7% 2.9% - 30.6% 5,490	*8.9% 2.4% - 15.3% 2,909	*7.3% 0.9% - 13.7% 2,861	*18.6% 4.9% - 32.2% 8,594
Non-Hispanic White	30.3% 21.5% - 39.0% 16,833	32.6% 22.6% - 42.6% 18,857	15.4% 7.9% - 22.9% 8,249	21.0% 13.8% - 28.3% 13,234	20.5% 13.4% - 27.7% 10,108	19.3% 12.4% - 26.2% 11,140	22.7% 13.4% - 31.9% 16,907
Hispanic	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Female	22.5% 13.6% - 31.5% 9,365	24.4% 14.1% - 34.6% 10,973	20.1% 11.4% - 28.8% 9,386	15.4% 9.4% - 21.3% 7,230	13.2% 7.6% - 18.9% 6,628	11.1% 6.0% - 16.3% 5,898	8.9% 4.4% - 13.5% 6,128
Male	30.3% 19.1% - 41.4% 11,897	24.4% 13.2% - 35.9% 13,109	*15.5% 3.3% - 27.8% 7,200	20.5% 10.8% - 30.3% 12,584	23.1% 11.2% - 34.9% 11,925	16.6% 9.1% - 24.2% 10,260	30.0% 17.6% - 42.5% 19,890
Highest Education: ≤ High School/GED	33.2% 21.6% - 44.9% 10,971	23.8% 11.6% - 35.9% 8,313	*24.2% 8.3% - 40.2% 8,231	32.1% 18.7% - 45.5% 11,783	23.8% 10.6% - 37.0% 10,473	28.8% 17.9% - 39.8% 9,920	26.7% 14.6% - 38.7% 15,060
Highest Education: ≥ 4 Years of College	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*9.6% 0.6% - 18.6% 2,962	*8.1% 1.2% - 15.0% 2,664	*11.4% 3.2% - 19.6% 3,783	*11.1% 4.4% - 17.9% 4,090	*3.7% 0.6% - 6.8% 1,491	*3.0% 0.0% - 5.9% 896
Household Income: < \$50,000 Annually	25.4% 15.3% - 35.6% 8,885	29.1% 15.7% - 42.5% 11,762	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	35.6% 17.2% - 54.0% 7,740	*24.6% 5.8% - 43.4% 6,472	22.4% 10.8% - 34.0% 6,053	25.9% 13.8% - 38.1% 12,116
Household Income: ≥ \$50,000 Annually	23.6% 13.5% - 33.7% 9,330	21.0% 10.8% - 31.1% 10,503	16.4% 8.7% - 24.1% 9,340	13.4% 7.5% - 19.2% 9,823	15.1% 8.7% - 21.5% 9,833	11.3% 5.7% - 16.9% 8,323	17.3% 7.2% - 27.3% 12,982

% OF HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS

(Estimated # of Households with Resident Minor Children and a Resident Adult Smoker / # Households with Resident Minor Children)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Proportion with Adult Smokers	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	31.4% 18.4% - 44.5% 17,600



Dorchester County

Tobacco Use by Adults: 2000–2012

Maryland Department of Health and Mental Hygiene—Center for Tobacco Prevention and Control
(See Appendix—Statewide and Jurisdiction Data for information on how to use and interpret these tables)

Historical jurisdiction estimates in this report may vary from those previously published. A methodology change in 2011 was designed to improve estimates of risk behaviors. Previous BRFSS estimates cannot be compared to the 2011 (and thereafter) data because of the significant differences in weighting methodology.

CURRENT USE OF ANY TOBACCO PRODUCT

Adult (Age 18 and older) Estimated Current Use of One or More Tobacco Products (Estimated # Adults Currently Using Any Tobacco Product / # All Adults)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
All Adults	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		16.9% 7.8% - 26.0% 4,694

CURRENT USE OF SPECIFIC TOBACCO PRODUCTS

Adult (Age 18 and older) Estimated Current Use of Specific Tobacco Products (Estimated # Currently Using Specified Tobacco Product / # All Adults)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Cigarettes	*27.4% 10.9% - 43.8% 6,769	33.3% 17.3% - 49.4% 10,384	28.4% 15.5% - 41.4% 8,430	20.7% 13.3% - 28.1% 5,058	19.7% 11.0% - 28.3% 4,485	13.9% 7.0% - 20.8% 3,136		16.3% 7.1% - 25.5% 4,389
Cigars	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*0.5% 0.0% - 1.1% 124
Smokeless Tobacco	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	*1.7% 0.0% - 4.4% 392		0.2% 0.0% - 0.5% 65
All Other Tobacco Products	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*0.7% 0.0% - 1.9% 181

INITIATION AND CESSATION OF CIGARETTE SMOKING IN PAST YEAR

INITIATION—Percent of All Maryland Adults Who Report First Smoking a Cigarette During the Past Year (Est. # First Tried Cigarette Smoking Past Year / # All Adults)

CESSATION—Percent of All Maryland Adults Who Report Quitting Cigarette Smoking During the Past Year (Est. # Quit Smoking Cigarettes Past Year / # All Adults)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Initiation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Cessation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*0.9% 0.1% - 1.8% 276

PREGNANT WOMEN — Cigarette Smoking At Any Time During Pregnancy

Maryland Residents Regardless of Where Delivery Occurred— **Birth Certificate Data, 2012**—Percent = Number Reported Smoking / Total Number

Percentage Smoking (%) Number Smoking (#)	2000	2002	2004	2006	2008	2010	2012
Pregnant Women	18.6% 61	16.6% 51	13.2% 48	13.1% 53	14.4% 65	19.9% 76	11.8% 49

SELECTED POPULATIONS — Current Cigarette Smoking


(Estimated # In Selected Population Currently Smoking Cigarettes / # All Adults In the Selected Population)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Non-Hispanic Black	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES
Non-Hispanic White	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	24.8% 15.9% - 33.7% 4,040	17.4% 8.7% - 26.0% 2,801	*12.5% 4.8% - 20.2% 1,743	23.1% 10.1% - 36.2% 3,062
Hispanic	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Female	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	20.2% 11.2% - 29.3% 2,723	13.3% 6.4% - 20.2% 1,440	*14.6% 5.7% - 23.4% 1,812	*15.8% 4.8% - 26.9% 2,239
Male	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	21.3% 9.1% - 33.6% 2,335	25.4% 10.6% - 40.2% 3,045	*13.2% 2.3% - 24.0% 1,323	*16.7% 1.8% - 31.6% 2,149
Highest Education: <= High School/GED	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	21.9% 10.9% - 32.8% 2,968	24.1% 11.1% - 37.0% 2,952	*15.7% 6.0% - 25.5% 1,842	*14.2% 4.8% - 23.6% 2,471
Highest Education: >= 4 Years of College	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*10.4% 1.2% - 19.7% 586	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*5.9% 0.6% - 11.2% 307
Household Income: < \$50,000 Annually	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	18.8% 8.4% - 29.1% 1,939	28.9% 14.0% - 43.8% 2,695	*16.8% 6.6% - 27.1% 1,790	*12.6% 1.7% - 23.5% 2,109
Household Income: >= \$50,000 Annually	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*17.3% 6.5% - 28.0% 1,282	*15.7% 3.1% - 28.3% 1,539	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*8.8% 1.3% - 16.3% 606

% OF HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS

(Estimated # of Households with Resident Minor Children and a Resident Adult Smoker / # Households with Resident Minor Children)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Proportion with Adult Smokers	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE



Frederick County

Tobacco Use by Adults: 2000–2012

Maryland Department of Health and Mental Hygiene—Center for Tobacco Prevention and Control
(See Appendix—Statewide and Jurisdiction Data for information on how to use and interpret these tables)

Historical jurisdiction estimates in this report may vary from those previously published. A methodology change in 2011 was designed to improve estimates of risk behaviors. Previous BRFSS estimates cannot be compared to the 2011 (and thereafter) data because of the significant differences in weighting methodology.

CURRENT USE OF ANY TOBACCO PRODUCT								
Adult (Age 18 and older) Estimated Current Use of One or More Tobacco Products (Estimated # Adults Currently Using Any Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
All Adults	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		19.7% 13.5% - 26.0% 32,315
CURRENT USE OF SPECIFIC TOBACCO PRODUCTS								
Adult (Age 18 and older) Estimated Current Use of Specific Tobacco Products (Estimated # Currently Using Specified Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Cigarettes	23.3% 17.1% - 29.5% 32,186	18.7% 13.3% - 24.0% 26,100	24.6% 18.2% - 30.9% 39,136	18.7% 14.1% - 23.3% 32,087	13.9% 10.3% - 17.5% 23,304	12.0% 8.1% - 15.9% 19,405		16.5% 10.6% - 22.4% 26,870
Cigars	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*6.0% 1.5% - 10.5% 9,114
Smokeless Tobacco	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	*2.6% 0.9% - 4.2% 4,122		BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES
All Other Tobacco Products	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES
INITIATION AND CESSATION OF CIGARETTE SMOKING IN PAST YEAR								
INITIATION—Percent of All Maryland Adults Who Report First Smoking a Cigarette During the Past Year (Est. # First Tried Cigarette Smoking Past Year / # All Adults) CESSATION—Percent of All Maryland Adults Who Report Quitting Cigarette Smoking During the Past Year (Est. # Quit Smoking Cigarettes Past Year / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Initiation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Cessation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*1.7% 0.5% - 2.8% 2,743

PREGNANT WOMEN — Cigarette Smoking At Any Time During Pregnancy

Maryland Residents Regardless of Where Delivery Occurred— Birth Certificate Data, 2012—Percent = Number Reported Smoking / Total Number

Percentage Smoking (%) Number Smoking (#)	2000	2002	2004	2006	2008	2010	2012
Pregnant Women	12.1% 351	9.2% 276	8.6% 254	7.5% 231	7.4% 219	9.9% 281	9.0% 248

SELECTED POPULATIONS — Current Cigarette Smoking

(Estimated # In Selected Population Currently Smoking Cigarettes / # All Adults In the Selected Population)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Non-Hispanic Black	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Non-Hispanic White	20.0% 14.7% - 25.3% 23,366	19.1% 13.5% - 24.6% 22,378	24.8% 18.1% - 31.6% 34,277	17.9% 13.8% - 21.9% 26,271	16.1% 11.9% - 20.4% 21,914	13.5% 8.8% - 18.2% 17,445	14.0% 9.0% - 19.0% 17,838
Hispanic	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Female	23.8% 16.4% - 31.3% 15,817	15.3% 9.3% - 21.3% 11,334	24.3% 16.4% - 32.2% 18,655	16.4% 12.0% - 20.9% 14,628	14.8% 10.3% - 19.2% 12,426	11.5% 6.2% - 16.8% 8,933	13.7% 6.1% - 21.3% 11,534
Male	22.8% 13.0% - 32.5% 16,368	22.5% 13.2% - 31.7% 14,765	24.8% 15.0% - 34.6% 20,572	21.1% 12.9% - 29.2% 17,459	13.0% 7.4% - 18.6% 10,878	12.5% 6.8% - 18.3% 10,472	19.5% 10.5% - 28.5% 15,336
Highest Education: ≤ High School/GED	32.7% 19.3% - 46.0% 14,579	27.6% 17.4% - 37.9% 11,789	41.7% 29.3% - 54.2% 25,151	26.1% 16.6% - 35.7% 15,256	19.3% 12.4% - 26.2% 11,214	22.6% 13.0% - 32.2% 12,093	24.9% 11.5% - 38.3% 14,265
Highest Education: ≥ 4 Years of College	14.5% 7.2% - 21.8% 8,891	*8.8% 2.5% - 15.1% 4,786	12.0% 5.2% - 18.8% 7,696	12.4% 7.1% - 17.7% 8,555	*8.4% 3.4% - 13.4% 5,999	*3.3% 1.2% - 5.4% 2,389	*10.2% 2.8% - 17.5% 5,832
Household Income: < \$50,000 Annually	26.4% 17.9% - 35.4% 12,252	23.8% 13.3% - 34.2% 10,177	42.1% 28.9% - 55.3% 18,609	29.4% 20.1% - 38.7% 10,663	19.0% 10.6% - 27.3% 8,140	*19.4% 7.8% - 30.9% 7,883	23.9% 10.1% - 37.6% 13,082
Household Income: ≥ \$50,000 Annually	21.0% 13.1% - 28.9% 16,039	14.4% 8.2% - 20.7% 11,691	14.4% 8.6% - 20.2% 13,649	17.5% 11.6% - 23.5% 20,586	13.2% 8.6% - 17.8% 13,975	10.0% 6.4% - 13.6% 9,631	12.1% 6.1% - 18.0% 11,097

% OF HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS

(Estimated # of Households with Resident Minor Children and a Resident Adult Smoker / # Households with Resident Minor Children)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Proportion with Adult Smokers	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	23.4% 11.4% - 35.4% 14,507



Garrett County

Tobacco Use by Adults: 2000–2012

Maryland Department of Health and Mental Hygiene—Center for Tobacco Prevention and Control
(See Appendix—Statewide and Jurisdiction Data for information on how to use and interpret these tables)

Historical jurisdiction estimates in this report may vary from those previously published. A methodology change in 2011 was designed to improve estimates of risk behaviors. Previous BRFSS estimates cannot be compared to the 2011 (and thereafter) data because of the significant differences in weighting methodology.

CURRENT USE OF ANY TOBACCO PRODUCT								
Adult (Age 18 and older) Estimated Current Use of One or More Tobacco Products (Estimated # Adults Currently Using Any Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
All Adults	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		23.7% 14.4% - 32.9% 5,317
CURRENT USE OF SPECIFIC TOBACCO PRODUCTS								
Adult (Age 18 and older) Estimated Current Use of Specific Tobacco Products (Estimated # Currently Using Specified Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Cigarettes	20.2% 6.5% - 33.9% 4,748	19.9% 8.1% - 31.7% 5,123	23.1% 10.6% - 35.7% 6,136	16.8% 10.6% - 23.0% 4,886	18.0% 11.4% - 24.6% 5,137	19.0% 9.9% - 28.1% 5,038		20.8% 11.4% - 30.2% 4,662
Cigars	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*5.1% 1.2% - 8.4% 1,010
Smokeless Tobacco	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	*6.0% 1.0% - 11.0% 1,590		*4.0% 0.7% - 6.4% 887
All Other Tobacco Products	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*2.6% 0.0% - 5.8% 515
INITIATION AND CESSATION OF CIGARETTE SMOKING IN PAST YEAR								
INITIATION—Percent of All Maryland Adults Who Report First Smoking a Cigarette During the Past Year (Est. # First Tried Cigarette Smoking Past Year / # All Adults) CESSATION—Percent of All Maryland Adults Who Report Quitting Cigarette Smoking During the Past Year (Est. # Quit Smoking Cigarettes Past Year / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Initiation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Cessation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*2.3% 0.2% - 4.5% 578

PREGNANT WOMEN — Cigarette Smoking At Any Time During Pregnancy

Maryland Residents Regardless of Where Delivery Occurred— **Birth Certificate Data, 2012**—Percent = Number Reported Smoking / Total Number

Percentage Smoking (%) Number Smoking (#)	2000	2002	2004	2006	2008	2010	2012
Pregnant Women	16.8% 56	17.0% 51	20.9% 66	16.2% 48	17.0% 47	16.9% 47	17.4% 50

SELECTED POPULATIONS — Current Cigarette Smoking


(Estimated # In Selected Population Currently Smoking Cigarettes / # All Adults In the Selected Population)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Non-Hispanic Black	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Non-Hispanic White	*20.7% 6.7% - 34.7% 4,748	20.3% 8.3% - 32.3% 5,123	25.5% 12.1% - 38.9% 6,136	17.4% 10.8% - 23.9% 4,656	19.2% 12.2% - 26.2% 5,025	15.5% 8.6% - 22.4% 3,805	20.9% 12.3% - 29.4% 4,523
Hispanic	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Female	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	21.2% 12.7% - 29.7% 2,911	17.8% 10.1% - 25.5% 3,102	11.5% 5.3% - 17.7% 1,761	*20.1% 5.7% - 34.5% 2,442
Male	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*12.8% 3.9% - 21.7% 1,976	*18.3% 6.2% - 30.3% 2,036	*29.4% 11.2% - 47.5% 3,276	21.6% 8.9% - 34.3% 2,220
Highest Education: ≤ High School/GED	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	20.7% 11.3% - 30.1% 3,219	19.7% 10.8% - 28.7% 3,086	*21.0% 7.2% - 34.8% 3,007	26.6% 14.7% - 38.4% 3,170
Highest Education: ≥ 4 Years of College	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES	*10.3% 0.6% - 20.0% 733	*7.6% 1.1% - 14.1% 334
Household Income: < \$50,000 Annually	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	24.6% 14.6% - 34.6% 3,690	14.9% 8.4% - 21.5% 2,312	24.7% 11.0% - 38.4% 3,504	26.3% 11.7% - 40.9% 3,209
Household Income: ≥ \$50,000 Annually	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*10.6% 1.1% - 20.1% 962	*16.3% 4.7% - 28.0% 1,545	*16.0% 2.5% - 29.4% 1,502	*18.4% 3.9% - 32.8% 1,453

% OF HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS

(Estimated # of Households with Resident Minor Children and a Resident Adult Smoker / # Households with Resident Minor Children)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Proportion with Adult Smokers	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	20.8% 9.5% - 32.2% 947



Harford County

Tobacco Use by Adults: 2000–2012

Maryland Department of Health and Mental Hygiene—Center for Tobacco Prevention and Control
(See Appendix—Statewide and Jurisdiction Data for information on how to use and interpret these tables)

Historical jurisdiction estimates in this report may vary from those previously published. A methodology change in 2011 was designed to improve estimates of risk behaviors. Previous BRFSS estimates cannot be compared to the 2011 (and thereafter) data because of the significant differences in weighting methodology.

CURRENT USE OF ANY TOBACCO PRODUCT								
Adult (Age 18 and older) Estimated Current Use of One or More Tobacco Products (Estimated # Adults Currently Using Any Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
All Adults	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		24.4% 18.8% - 30.1% 49,272
CURRENT USE OF SPECIFIC TOBACCO PRODUCTS								
Adult (Age 18 and older) Estimated Current Use of Specific Tobacco Products (Estimated # Currently Using Specified Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Cigarettes	16.5% 10.1% - 22.9% 31,558	20.0% 12.5% - 27.5% 29,626	15.9% 9.1% - 22.1% 28,006	18.1% 12.4% - 23.4% 36,475	20.8% 14.3% - 27.3% 40,879	19.9% 13.1% - 28.7% 35,944		21.7% 16.3% - 27.1% 43,602
Cigars	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*3.7% 0.8% - 6.5% 6,606
Smokeless Tobacco	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	*1.9% 0.0% - 4.0% 3,488		*1.6% 0.3% - 3.0% 3,423
All Other Tobacco Products	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*0.8% 0.0% - 1.8% 1,504
INITIATION AND CESSATION OF CIGARETTE SMOKING IN PAST YEAR								
INITIATION—Percent of All Maryland Adults Who Report First Smoking a Cigarette During the Past Year (Est. # First Tried Cigarette Smoking Past Year / # All Adults) CESSATION—Percent of All Maryland Adults Who Report Quitting Cigarette Smoking During the Past Year (Est. # Quit Smoking Cigarettes Past Year / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Initiation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Cessation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*2.2% 0.8% - 3.7% 4,600

PREGNANT WOMEN — Cigarette Smoking At Any Time During Pregnancy

Maryland Residents Regardless of Where Delivery Occurred— Birth Certificate Data, 2012—Percent = Number Reported Smoking / Total Number

Percentage Smoking (%) Number Smoking (#)	2000	2002	2004	2006	2008	2010	2012
Pregnant Women	13.1% 386	12.4% 360	10.1% 301	11.3% 342	9.1% 271	10.4% 281	10.8% 286

SELECTED POPULATIONS — Current Cigarette Smoking

(Estimated # In Selected Population Currently Smoking Cigarettes / # All Adults In the Selected Population)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Non-Hispanic Black	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*27.5% 10.3% - 44.6% 7,882
Non-Hispanic White	13.2% 6.9% - 19.6% 21,197	22.3% 13.8% - 30.8% 26,907	15.1% 8.1% - 22.1% 22,267	18.8% 12.8% - 24.8% 28,180	22.3% 15.0% - 29.6% 34,071	22.2% 14.5% - 30.0% 33,616	22.4% 16.4% - 28.4% 33,397
Hispanic	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Female	16.7% 8.5% - 24.9% 17,774	17.6% 9.1% - 26.1% 13,537	16.7% 7.4% - 25.9% 16,369	19.2% 11.7% - 26.6% 17,705	20.6% 12.9% - 28.2% 19,577	14.2% 7.7% - 20.8% 11,875	22.8% 14.9% - 30.6% 24,223
Male	*16.2% 6.0% - 26.3% 13,785	22.7% 10.1% - 35.3% 16,090	*14.9% 4.9% - 24.8% 11,637	17.2% 8.7% - 25.7% 18,770	20.9% 10.6% - 31.3% 21,302	24.8% 13.8% - 35.9% 24,069	20.5% 13.3% - 27.8% 19,379
Highest Education: ≤ High School/GED	23.5% 11.6% - 35.3% 18,648	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*18.1% 5.2% - 30.9% 10,151	23.9% 13.2% - 34.6% 18,125	24.7% 12.9% - 36.5% 17,988	37.4% 23.0% - 51.7% 23,055	34.8% 24.2% - 45.5% 26,325
Highest Education: ≥ 4 Years of College	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES	*9.2% 2.0% - 16.3% 5,385	*17.5% 7.0% - 28.0% 10,473	*7.4% 1.9% - 12.8% 5,993	*8.7% 2.8% - 14.5% 6,042	*6.1% 0.5% - 11.6% 3,900	*6.1% 1.1% - 11.1% 3,692
Household Income: < \$50,000 Annually	20.7% 10.9% - 30.5% 18,836	31.2% 17.5% - 44.9% 20,166	28.2% 14.5% - 42.0% 18,854	40.0% 26.2% - 53.9% 22,882	19.1% 9.4% - 28.9% 12,422	37.6% 23.4% - 51.9% 21,435	33.7% 22.2% - 45.1% 23,178
Household Income: ≥ \$50,000 Annually	*12.9% 2.9% - 22.9% 10,072	*12.0% 4.3% - 19.7% 8,313	*9.5% 2.8% - 16.2% 8,707	11.9% 6.2% - 17.5% 13,533	19.2% 10.3% - 28.1% 19,406	11.9% 9.1% - 12.2% 11,912	14.3% 8.7% - 20.0% 15,190

% OF HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS

(Estimated # of Households with Resident Minor Children and a Resident Adult Smoker / # Households with Resident Minor Children)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Proportion with Adult Smokers	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	41.6% 30.3% - 53.0% 31,005



Howard County

Tobacco Use by Adults: 2000–2012

Maryland Department of Health and Mental Hygiene—Center for Tobacco Prevention and Control
(See Appendix—Statewide and Jurisdiction Data for information on how to use and interpret these tables)

Historical jurisdiction estimates in this report may vary from those previously published. A methodology change in 2011 was designed to improve estimates of risk behaviors. Previous BRFSS estimates cannot be compared to the 2011 (and thereafter) data because of the significant differences in weighting methodology.

CURRENT USE OF ANY TOBACCO PRODUCT

Adult (Age 18 and older) Estimated Current Use of One or More Tobacco Products (Estimated # Adults Currently Using Any Tobacco Product / # All Adults)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
All Adults	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		13.0% 8.4% - 17.6% 28,642

CURRENT USE OF SPECIFIC TOBACCO PRODUCTS

Adult (Age 18 and older) Estimated Current Use of Specific Tobacco Products (Estimated # Currently Using Specified Tobacco Product / # All Adults)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Cigarettes	12.6% 6.8% - 18.4% 21,005	21.2% 13.0% - 29.4% 38,437	12.9% 6.2% - 19.7% 24,991	12.6% 8.2% - 16.9% 29,226	6.5% 3.1% - 9.9% 15,859	8.8% 4.2% - 13.5% 20,796		9.5% 5.3% - 13.7% 20,777
Cigars	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*4.4% 0.4% - 8.3% 8,750
Smokeless Tobacco	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES		*3.7% 0.3% - 7.14% 8,296
All Other Tobacco Products	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES

INITIATION AND CESSATION OF CIGARETTE SMOKING IN PAST YEAR

INITIATION—Percent of All Maryland Adults Who Report First Smoking a Cigarette During the Past Year (Est. # First Tried Cigarette Smoking Past Year / # All Adults)

CESSATION—Percent of All Maryland Adults Who Report Quitting Cigarette Smoking During the Past Year (Est. # Quit Smoking Cigarettes Past Year / # All Adults)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Initiation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Cessation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*1.7% 0.0% - 3.6% 3,732

PREGNANT WOMEN — Cigarette Smoking At Any Time During Pregnancy

Maryland Residents Regardless of Where Delivery Occurred— Birth Certificate Data, 2012—Percent = Number Reported Smoking / Total Number

Percentage Smoking (%) Number Smoking (#)	2000	2002	2004	2006	2008	2010	2012
Pregnant Women	3.3% 119	3.3% 117	3.3% 116	2.9% 99	2.2% 74	3.1% 104	3.5% 121

SELECTED POPULATIONS — Current Cigarette Smoking


(Estimated # In Selected Population Currently Smoking Cigarettes / # All Adults In the Selected Population)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Non-Hispanic Black	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*10.9% 1.8% - 19.9% 3,776
Non-Hispanic White	13.4% 7.2% - 19.6% 17,532	23.9% 14.2% - 33.7% 32,884	*8.3% 2.7% - 13.9% 11,719	12.5% 7.3% - 17.8% 18,708	7.6% 3.5% - 11.7% 11,719	*11.0% 4.3% - 17.6% 16,556	8.4% 4.4% - 12.4% 11,354
Hispanic	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Female	12.7% 6.0% - 19.4% 11,995	20.4% 9.7% - 31.1% 17,912	*10.8% 2.4% - 19.3% 11,745	9.9% 5.4% - 14.5% 12,126	4.4% 2.1% - 6.7% 4,821	*10.0% 2.6% - 17.4% 12,393	9.3% 4.9% - 13.7% 10,127
Male	*12.5% 2.3% - 22.7% 9,011	21.9% 9.7% - 34.1% 20,524	*15.6% 4.8% - 26.5% 13,246	15.5% 7.9% - 23.0% 17,100	*8.1% 2.3% - 13.8% 11,038	*7.5% 2.4% - 12.7% 8,403	*9.7% 2.5% - 16.8% 10,650
Highest Education: ≤ High School/GED	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*25.0% 3.4% - 46.6% 8,838	*8.4% 2.1% - 14.7% 3,834
Highest Education: ≥ 4 Years of College	*4.3% 0.4% - 8.1% 4,519	*14.1% 5.2% - 23.1% 15,462	*6.6% 1.8% - 11.4% 7,584	9.6% 4.9% - 14.2% 15,038	*3.3% 1.0% - 5.7% 5,009	*5.4% 2.2% - 8.5% 7,986	*3.8% 1.3% - 6.4% 4,427
Household Income: < \$50,000 Annually	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*26.0% 10.6% - 41.4% 10,403	*18.0% 2.5% - 33.6% 5,974	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES	*14.1% 4.1% - 24.1% 6,370
Household Income: ≥ \$50,000 Annually	*11.8% 4.6% - 19.0% 13,727	18.4% 9.4% - 27.4% 21,280	*4.5% 0.6% - 8.3% 6,202	11.2% 6.5% - 15.8% 18,131	*5.0% 1.8% - 8.2% 9,472	7.0% 9.1% - 12.2% 10,676	9.7% 4.1% - 15.2% 136,686

% OF HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS

(Estimated # of Households with Resident Minor Children and a Resident Adult Smoker / # Households with Resident Minor Children)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Proportion with Adult Smokers	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	12.7% 6.0% - 19.4% 10,751



Kent County

Tobacco Use by Adults: 2000–2012

Maryland Department of Health and Mental Hygiene—Center for Tobacco Prevention and Control
(See Appendix—Statewide and Jurisdiction Data for information on how to use and interpret these tables)

Historical jurisdiction estimates in this report may vary from those previously published. A methodology change in 2011 was designed to improve estimates of risk behaviors. Previous BRFSS estimates cannot be compared to the 2011 (and thereafter) data because of the significant differences in weighting methodology.

CURRENT USE OF ANY TOBACCO PRODUCT								
Adult (Age 18 and older) Estimated Current Use of One or More Tobacco Products (Estimated # Adults Currently Using Any Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
All Adults	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		21.7% 6.2% - 37.2% 3,960
CURRENT USE OF SPECIFIC TOBACCO PRODUCTS								
Adult (Age 18 and older) Estimated Current Use of Specific Tobacco Products (Estimated # Currently Using Specified Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Cigarettes	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	19.1% 11.4% - 26.8% 3,076	13.7% 7.7% - 19.6% 2,768	14.0% 7.2% - 20.9% 1,903		*19.2% 3.7% - 34.7% 3,511
Cigars	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		2.5% 0.0% - 5.1% 397
Smokeless Tobacco	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES		*1.9% 0.0% - 4.1% 354
All Other Tobacco Products	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*0.5% 0.0% - 1.2% 82
INITIATION AND CESSATION OF CIGARETTE SMOKING IN PAST YEAR								
INITIATION—Percent of All Maryland Adults Who Report First Smoking a Cigarette During the Past Year (Est. # First Tried Cigarette Smoking Past Year / # All Adults)								
CESSATION—Percent of All Maryland Adults Who Report Quitting Cigarette Smoking During the Past Year (Est. # Quit Smoking Cigarettes Past Year / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Initiation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Cessation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*1.0% 0.0% - 2.1% 187

PREGNANT WOMEN — Cigarette Smoking At Any Time During Pregnancy

Maryland Residents Regardless of Where Delivery Occurred— Birth Certificate Data, 2012—Percent = Number Reported Smoking / Total Number

Percentage Smoking (%) Number Smoking (#)	2000	2002	2004	2006	2008	2010	2012
Pregnant Women	21.2% 43	16.6% 26	16.3% 33	19.0% 36	13.7% 30	18.1% 30	15.8% 28

SELECTED POPULATIONS — Current Cigarette Smoking

(Estimated # In Selected Population Currently Smoking Cigarettes / # All Adults In the Selected Population)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Non-Hispanic Black	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Non-Hispanic White	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	17.4% 9.5% - 25.4% 2,377	15.4% 8.7% - 22.2% 2,498	14.8% 6.8% - 22.8% 1,640	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES
Hispanic	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Female	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	21.8% 11.7% - 31.9% 1,595	21.4% 12.0% - 30.9% 2,251	*13.0% 5.0% - 21.8% 1,028	*28.3% 1.8% - 54.8% 2,933
Male	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*16.8% 5.6% - 28.0% 1,481	*5.3% 0.5% - 10.1% 517	*15.4% 3.4% - 27.5% 874	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES
Highest Education: <= High School/GED	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	26.9% 12.5% - 41.4% 1,591	*14.2% 5.0% - 23.3% 1,289	*9.8% 1.7% - 17.9% 501	*35.1% 8.2% - 62.1% 2,932
Highest Education: >= 4 Years of College	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES	*10.5% 2.1% - 18.9% 260	*14.4% 1.7% - 27.2% 800	*2.1% 0.0% - 4.7% 118
Household Income: < \$50,000 Annually	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	16.9% 8.4% - 25.3% 1,284	*15.1% 5.8% - 24.3% 1,335	*12.3% 3.9% - 20.7% 640	*33.0% 1.8% - 64.2% 3,016
Household Income: >= \$50,000 Annually	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*22.3% 8.0% - 36.6% 1,454	*17.1% 5.9% - 28.4% 1,311	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*6.7% 0.5% - 12.8% 456

% OF HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS

(Estimated # of Households with Resident Minor Children and a Resident Adult Smoker / # Households with Resident Minor Children)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Proportion with Adult Smokers	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE



Montgomery County

Tobacco Use by Adults: 2000–2012

Maryland Department of Health and Mental Hygiene—Center for Tobacco Prevention and Control
(See Appendix—Statewide and Jurisdiction Data for information on how to use and interpret these tables)

Historical jurisdiction estimates in this report may vary from those previously published. A methodology change in 2011 was designed to improve estimates of risk behaviors. Previous BRFSS estimates cannot be compared to the 2011 (and thereafter) data because of the significant differences in weighting methodology.

CURRENT USE OF ANY TOBACCO PRODUCT								
Adult (Age 18 and older) Estimated Current Use of One or More Tobacco Products (Estimated # Adults Currently Using Any Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
All Adults	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		10.9% 8.2% - 13.7% 81,828
CURRENT USE OF SPECIFIC TOBACCO PRODUCTS								
Adult (Age 18 and older) Estimated Current Use of Specific Tobacco Products (Estimated # Currently Using Specified Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Cigarettes	13.2% 10.2% - 16.2% 85,839	16.2% 12.1% - 20.4% 115,251	8.1% 5.1% - 11.1% 55,470	9.2% 6.9% - 11.5% 70,439	8.1% 5.4% - 10.9% 57,657	7.8% 5.3% - 10.3% 55,245		7.5% 5.5% - 9.6% 55,857
Cigars	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*3.5% 1.3% - 5.6% 20,964
Smokeless Tobacco	*1.4% 0.4% - 2.4% 9,104	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	*1.1% 0.4% - 1.7% 7,564		*1.3% 0.5% - 2.2% 6,624
All Other Tobacco Products	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		1.1% 0.4% - 1.8% 6,624
INITIATION AND CESSATION OF CIGARETTE SMOKING IN PAST YEAR								
INITIATION—Percent of All Maryland Adults Who Report First Smoking a Cigarette During the Past Year (Est. # First Tried Cigarette Smoking Past Year / # All Adults) CESSATION—Percent of All Maryland Adults Who Report Quitting Cigarette Smoking During the Past Year (Est. # Quit Smoking Cigarettes Past Year / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Initiation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*0.7% 0.0% - 1.5% 5,761
Cessation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		3.3% 1.8% - 4.9% 26,531

PREGNANT WOMEN — Cigarette Smoking At Any Time During Pregnancy

Maryland Residents Regardless of Where Delivery Occurred — Birth Certificate Data, 2012 — Percent = Number Reported Smoking / Total Number

Percentage Smoking (%) Number Smoking (#)	2000	2002	2004	2006	2008	2010	2012
Pregnant Women	2.5% 327	1.3% 168	1.0% 142	0.7% 95	0.6% 80	1.6% 219	1.6% 213

SELECTED POPULATIONS — Current Cigarette Smoking

(Estimated # In Selected Population Currently Smoking Cigarettes / # All Adults In the Selected Population)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Non-Hispanic Black	19.1% 8.7% - 29.5% 11,667	*19.6% 5.6% - 33.6% 24,779	*7.2% 0.5% - 13.9% 6,976	*4.4% 0.6% - 8.3% 4,655	*18.7% 5.1% - 32.3% 19,590	*11.4% 4.2% - 18.6% 13,340	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES
Non-Hispanic White	13.3% 9.9% - 16.7% 60,646	12.9% 9.3% - 16.5% 55,042	8.6% 5.6% - 11.5% 37,612	12.2% 8.8% - 15.5% 55,487	7.4% 5.0% - 9.8% 29,677	7.4% 4.3% - 10.4% 29,518	9.1% 6.2% - 12.0% 32,182
Hispanic	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES
Female	11.5% 8.0% - 15.0% 37,837	13.6% 8.0% - 19.3% 46,544	4.2% 2.4% - 6.0% 14,746	9.0% 6.2% - 11.8% 34,375	7.0% 4.1% - 10.0% 25,151	5.1% 2.6% - 7.5% 18,506	6.9% 4.0% - 9.7% 22,861
Male	14.9% 9.9% - 19.8% 48,003	18.7% 12.6% - 24.7% 68,707	12.2% 6.6% - 17.9% 40,725	9.4% 5.7% - 13.1% 36,064	9.2% 4.6% - 13.8% 32,506	10.8% 6.3% - 15.2% 36,739	8.1% 5.1% - 11.1% 32,996
Highest Education: <= High School/GED	28.5% 17.3% - 39.7% 30,043	30.7% 18.5% - 42.9% 37,393	*6.9% 0.8% - 13.0% 8,331	12.9% 6.4% - 19.5% 15,201	*19.0% 6.1% - 31.9% 20,676	*7.1% 1.7% - 12.5% 8,803	8.4% 3.8% - 13.0% 15,958
Highest Education: >= 4 Years of College	7.9% 5.1% - 10.7% 31,424	10.8% 5.9% - 15.8% 49,619	6.5% 3.9% - 9.0% 28,252	7.0% 4.4% - 9.7% 34,660	3.7% 2.4% - 5.0% 17,822	6.9% 4.0% - 9.9% 29,669	4.9% 2.7% - 7.2% 19,254
Household Income: < \$50,000 Annually	19.4% 12.1% - 26.7% 35,541	21.2% 11.4% - 31.0% 42,214	8.0% 2.8% - 13.2% 12,624	15.1% 9.1% - 21.0% 23,923	9.5% 5.0% - 13.9% 12,225	12.9% 7.2% - 18.5% 18,645	7.8% 3.9% - 11.7% 16,499
Household Income: >= \$50,000 Annually	12.3% 8.7% - 15.9% 44,717	14.5% 9.9% - 19.2% 59,378	*7.8% 3.8% - 11.8% 34,179	8.1% 5.4% - 10.8% 40,105	6.5% 4.2% - 8.7% 31,389	7.5% 4.0% - 11.0% 31,732	7.7% 5.1% - 10.4% 34,304

% OF HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS

(Estimated # of Households with Resident Minor Children and a Resident Adult Smoker / # Households with Resident Minor Children)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Proportion with Adult Smokers	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	15.2% 9.7% - 20.8% 41,554



Prince George's County

Tobacco Use by Adults: 2000–2012

Maryland Department of Health and Mental Hygiene—Center for Tobacco Prevention and Control
(See Appendix—Statewide and Jurisdiction Data for information on how to use and interpret these tables)

Historical jurisdiction estimates in this report may vary from those previously published. A methodology change in 2011 was designed to improve estimates of risk behaviors. Previous BRFSS estimates cannot be compared to the 2011 (and thereafter) data because of the significant differences in weighting methodology.

CURRENT USE OF ANY TOBACCO PRODUCT								
Adult (Age 18 and older) Estimated Current Use of One or More Tobacco Products (Estimated # Adults Currently Using Any Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
All Adults	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		16.6% 13.1% - 20.1% 103,805
CURRENT USE OF SPECIFIC TOBACCO PRODUCTS								
Adult (Age 18 and older) Estimated Current Use of Specific Tobacco Products (Estimated # Currently Using Specified Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Cigarettes	16.5% 12.8% - 20.2% 95,997	18.3% 13.7% - 23.0% 103,727	18.2% 13.6% - 22.9% 109,349	17.2% 13.4% - 21.0% 92,157	12.0% 9.2% - 14.8% 75,290	13.6% 10.1% - 17.1% 88,535		14.2% 10.8% - 17.5% 88,296
Cigars	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*4.7% 2.6% - 6.7% 23,783
Smokeless Tobacco	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	*1.4% 0.1% - 2.7% 9,176		*1.3% 0.2% - 2.3% 7,991
All Other Tobacco Products	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*0.6% 0.0% - 1.2% 2,834
INITIATION AND CESSATION OF CIGARETTE SMOKING IN PAST YEAR								
INITIATION—Percent of All Maryland Adults Who Report First Smoking a Cigarette During the Past Year (Est. # First Tried Cigarette Smoking Past Year / # All Adults) CESSATION—Percent of All Maryland Adults Who Report Quitting Cigarette Smoking During the Past Year (Est. # Quit Smoking Cigarettes Past Year / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Initiation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES
Cessation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		2.2% 1.0% - 3.4% 13,903

PREGNANT WOMEN — Cigarette Smoking At Any Time During Pregnancy

Maryland Residents Regardless of Where Delivery Occurred— Birth Certificate Data, 2012—Percent = Number Reported Smoking / Total Number

Percentage Smoking (%) Number Smoking (#)	2000	2002	2004	2006	2008	2010	2012
Pregnant Women	3.6% 447	2.5% 316	1.9% 237	1.4% 177	1.7% 213	2.4% 290	2.4% 281

SELECTED POPULATIONS — Current Cigarette Smoking


(Estimated # In Selected Population Currently Smoking Cigarettes / # All Adults In the Selected Population)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Non-Hispanic Black	15.1% 10.6% - 19.6% 53,030	21.5% 15.0% - 28.1% 68,684	16.7% 10.7% - 22.7% 62,951	17.7% 12.6% - 22.8% 58,529	9.7% 6.5% - 12.8% 36,872	14.3% 10.2% - 18.5% 64,751	13.9% 10.0% - 17.7% 53,114
Non-Hispanic White	20.1% 13.0% - 27.1% 31,227	13.1% 7.7% - 18.5% 19,786	22.9% 13.9% - 31.9% 32,764	16.1% 10.6% - 21.6% 18,458	20.4% 12.7% - 28.0% 26,005	12.9% 7.3% - 18.5% 12,677	15.5% 8.6% - 22.4% 18,452
Hispanic	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*16.2% 1.8% - 30.6% 13,051
Female	14.8% 10.4% - 19.2% 45,618	13.5% 9.2% - 17.8% 42,791	18.1% 12.5% - 23.8% 56,823	14.0% 10.2% - 17.8% 39,148	11.9% 8.2% - 15.5% 39,826	9.9% 6.4% - 13.4% 33,189	11.8% 8.2% - 15.4% 38,113
Male	18.5% 12.4% - 24.6% 50,379	24.6% 15.8% - 33.3% 60,936	18.4% 10.9% - 25.8% 52,525	20.7% 14.0% - 27.4% 53,009	12.1% 7.8% - 16.4% 35,464	17.5% 11.4% - 23.5% 55,346	16.8% 11.0% - 22.5% 50,183
Highest Education: ≤ High School/GED	*25.3% 17.4% - 33.1% 51,213	25.2% 17.1% - 33.3% 52,286	25.7% 16.5% - 34.9% 57,818	19.3% 13.0% - 25.5% 36,470	17.1% 11.1% - 23.0% 39,992	17.7% 9.9% - 25.5% 34,955	25.2% 17.7% - 32.6% 58,242
Highest Education: ≥ 4 Years of College	11.5% 6.1% - 17.0% 21,474	*8.4% 3.0% - 13.7% 18,193	*6.0% 2.4% - 9.5% 12,981	11.5% 6.7% - 16.2% 23,660	7.8% 4.7% - 10.9% 18,808	5.6% 2.8% - 8.5% 14,076	5.1% 3.0% - 7.2% 9,468
Household Income: < \$50,000 Annually	19.4% 13.4% - 25.4% 46,791	21.6% 13.6% - 29.7% 53,584	20.5% 13.4% - 27.7% 42,699	22.5% 15.9% - 29.0% 42,876	14.9% 9.7% - 20.1% 34,940	16.4% 8.7% - 24.2% 33,577	22.4% 15.6% - 29.3% 52,946
Household Income: ≥ \$50,000 Annually	16.9% 11.4% - 22.4% 44,076	16.4% 10.8% - 22.0% 39,399	18.0% 11.2% - 24.9% 55,693	15.8% 10.1% - 21.5% 42,337	10.7% 7.2% - 14.3% 36,014	11.3% 7.5% - 15.0% 38,700	8.8% 6.0% - 11.5% 26,947

% OF HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS

(Estimated # of Households with Resident Minor Children and a Resident Adult Smoker / # Households with Resident Minor Children)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Proportion with Adult Smokers	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	21.4% 14.7% - 28.2% 48,264



Queen Anne's County

Tobacco Use by Adults: 2000–2012

Maryland Department of Health and Mental Hygiene—Center for Tobacco Prevention and Control
(See Appendix—Statewide and Jurisdiction Data for information on how to use and interpret these tables)

Historical jurisdiction estimates in this report may vary from those previously published. A methodology change in 2011 was designed to improve estimates of risk behaviors. Previous BRFSS estimates cannot be compared to the 2011 (and thereafter) data because of the significant differences in weighting methodology.

CURRENT USE OF ANY TOBACCO PRODUCT								
Adult (Age 18 and older) Estimated Current Use of One or More Tobacco Products (Estimated # Adults Currently Using Any Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
All Adults	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		19.3% 10.6% - 28.0% 5,970
CURRENT USE OF SPECIFIC TOBACCO PRODUCTS								
Adult (Age 18 and older) Estimated Current Use of Specific Tobacco Products (Estimated # Currently Using Specified Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Cigarettes	29.0% 15.8% - 42.1% 7,426	*13.6% 3.0% - 24.2% 3,891	*13.6% 3.3% - 24.0% 3,963	21.6% 15.3% - 27.9% 7,222	19.0% 12.4% - 25.7% 7,350	16.0% 9.6% - 22.4% 6,216		17.2% 8.6% - 25.7% 5,307
Cigars	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*2.1% 0.4% - 3.9% 603
Smokeless Tobacco	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	*4.5% 0.1% - 8.8% 1,740		BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES
All Other Tobacco Products	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*0.6% 0.0% - 1.5% 176
INITIATION AND CESSATION OF CIGARETTE SMOKING IN PAST YEAR								
INITIATION—Percent of All Maryland Adults Who Report First Smoking a Cigarette During the Past Year (Est. # First Tried Cigarette Smoking Past Year / # All Adults) CESSATION—Percent of All Maryland Adults Who Report Quitting Cigarette Smoking During the Past Year (Est. # Quit Smoking Cigarettes Past Year / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Initiation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Cessation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*1.3% 0.5% - 2.2% 419

PREGNANT WOMEN — Cigarette Smoking At Any Time During Pregnancy

Maryland Residents Regardless of Where Delivery Occurred— **Birth Certificate Data, 2012**—Percent = Number Reported Smoking / Total Number

Percentage Smoking (%) Number Smoking (#)	2000	2002	2004	2006	2008	2010	2012
Pregnant Women	15.6% 78	10.2% 54	10.8% 55	8.9% 46	6.2% 33	10.9% 53	10.2% 45

SELECTED POPULATIONS — Current Cigarette Smoking

(Estimated # In Selected Population Currently Smoking Cigarettes / # All Adults In the Selected Population)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Non-Hispanic Black	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Non-Hispanic White	26.4% 12.8% - 40.0% 6,079	*13.5% 4.0% - 23.0% 3,288	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	24.3% 17.2% - 31.4% 6,779	17.9% 11.4% - 24.5% 5,594	16.7% 9.9% - 23.6% 5,985	17.0% 7.5% - 26.5% 4,602
Hispanic	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Female	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	25.2% 17.5% - 32.8% 4,518	18.0% 10.9% - 25.1% 3,339	14.5% 6.5% - 22.5% 2,519	10.7% 5.6% - 15.9% 1,318
Male	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	17.4% 7.4% - 27.5% 2,704	20.0% 9.1% - 30.9% 4,011	17.2% 7.6% - 26.8% 3,698	*21.4% 7.5% - 35.5% 3,989
Highest Education: <= High School/GED	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	33.2% 21.5% - 44.9% 3,656	27.0% 15.2% - 38.8% 4,823	21.5% 8.9% - 34.1% 3,061	29.2% 13.4% - 45.0% 3,529
Highest Education: >= 4 Years of College	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*10.0% 3.3% - 16.7% 1,349	*8.6% 2.1% - 15.1% 1,054	*11.0% 2.9% - 19.0% 1,491	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES
Household Income: < \$50,000 Annually	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	29.7% 16.2% - 43.2% 2,620	33.7% 18.6% - 48.8% 3,951	*24.4% 9.4% - 39.5% 2,259	43.0% 24.8% - 61.3% 3,525
Household Income: >= \$50,000 Annually	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	19.6% 11.8% - 27.4% 4,238	12.3% 6.5% - 18.1% 2,513	13.4% 5.9% - 20.9% 3,230	*5.2% 1.9% - 8.5% 1,061

% OF HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS

(Estimated # of Households with Resident Minor Children and a Resident Adult Smoker / # Households with Resident Minor Children)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Proportion with Adult Smokers	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	23.6% 10.1% - 37.2% 3,477



Somerset County

Tobacco Use by Adults: 2000–2012

Maryland Department of Health and Mental Hygiene—Center for Tobacco Prevention and Control
(See Appendix—Statewide and Jurisdiction Data for information on how to use and interpret these tables)

Historical jurisdiction estimates in this report may vary from those previously published. A methodology change in 2011 was designed to improve estimates of risk behaviors. Previous BRFSS estimates cannot be compared to the 2011 (and thereafter) data because of the significant differences in weighting methodology.

CURRENT USE OF ANY TOBACCO PRODUCT								
Adult (Age 18 and older) Estimated Current Use of One or More Tobacco Products (Estimated # Adults Currently Using Any Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
All Adults	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		34.5% 16.0% - 53.0% 5,087
CURRENT USE OF SPECIFIC TOBACCO PRODUCTS								
Adult (Age 18 and older) Estimated Current Use of Specific Tobacco Products (Estimated # Currently Using Specified Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Cigarettes	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES	30.5% 20.7% - 40.3% 4,911		34.8% 16.0% - 53.0% 5,050
Cigars	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		13.5% 6.7% - 20.3% 1,842
Smokeless Tobacco	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES		*1.3% 0.0% - 2.9% 201
All Other Tobacco Products	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES
INITIATION AND CESSATION OF CIGARETTE SMOKING IN PAST YEAR								
INITIATION—Percent of All Maryland Adults Who Report First Smoking a Cigarette During the Past Year (Est. # First Tried Cigarette Smoking Past Year / # All Adults) CESSATION—Percent of All Maryland Adults Who Report Quitting Cigarette Smoking During the Past Year (Est. # Quit Smoking Cigarettes Past Year / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Initiation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Cessation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*2.8% 0.0% - 16.6% 2,259

PREGNANT WOMEN — Cigarette Smoking At Any Time During Pregnancy

Maryland Residents Regardless of Where Delivery Occurred— **Birth Certificate Data, 2012**—Percent = Number Reported Smoking / Total Number

Percentage Smoking (%) Number Smoking (#)	2000	2002	2004	2006	2008	2010	2012
Pregnant Women	20.9% 57	17.0% 44	16.5% 44	16.1% 45	14.8% 41	17.3% 47	13.3% 35

SELECTED POPULATIONS — Current Cigarette Smoking

(Estimated # In Selected Population Currently Smoking Cigarettes / # All Adults In the Selected Population)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Non-Hispanic Black	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Non-Hispanic White	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	30.8% 19.0% - 42.6% 3,031	25.6% 12.2% - 38.9% 2,567	*27.9% 8.9% - 46.9% 2,966	41.4% 18.8% - 64.1% 4,631
Hispanic	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Female	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*13.5% 3.3% - 23.8% 854	*14.4% 5.1% - 23.6% 995	*17.8% 5.6% - 30.0% 1,407	*38.7% 12.0% - 65.4% 3,998
Male	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Highest Education: ≤ High School/GED	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	37.6% 22.1% - 53.2% 2,603	32.7% 16.4% - 48.9% 3,042	44.4% 21.5% - 67.3% 3,893	*35.5% 13.3% - 57.6% 2,784
Highest Education: ≥ 4 Years of College	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Household Income: < \$50,000 Annually	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	31.7% 17.0% - 46.3% 3,288	37.1% 21.9% - 52.4% 3,486	41.4% 18.2% - 64.5% 3,755	*41.4% 10.9% - 71.9% 3,636
Household Income: ≥ \$50,000 Annually	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE

% OF HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS

(Estimated # of Households with Resident Minor Children and a Resident Adult Smoker / # Households with Resident Minor Children)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Proportion with Adult Smokers	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE



St. Mary's County

Tobacco Use by Adults: 2000–2012

Maryland Department of Health and Mental Hygiene—Center for Tobacco Prevention and Control
(See Appendix—Statewide and Jurisdiction Data for information on how to use and interpret these tables)

Historical jurisdiction estimates in this report may vary from those previously published. A methodology change in 2011 was designed to improve estimates of risk behaviors. Previous BRFSS estimates cannot be compared to the 2011 (and thereafter) data because of the significant differences in weighting methodology.

CURRENT USE OF ANY TOBACCO PRODUCT								
Adult (Age 18 and older) Estimated Current Use of One or More Tobacco Products (Estimated # Adults Currently Using Any Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
All Adults	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		23.4% 14.7% - 32.0% 20,585
CURRENT USE OF SPECIFIC TOBACCO PRODUCTS								
Adult (Age 18 and older) Estimated Current Use of Specific Tobacco Products (Estimated # Currently Using Specified Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Cigarettes	20.2% 12.4% - 27.9% 10,609	31.3% 21.5% - 41.0% 19,090	29.5% 20.0% - 39.0% 20,699	23.4% 15.9% - 30.9% 16,339	22.3% 15.6% - 29.0% 17,757	15.9% 10.8% - 20.9% 13,530		20.3% 11.8% - 28.8% 17,800
Cigars	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*4.4% 1.2% - 7.7% 3,340
Smokeless Tobacco	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	*3.7% 0.1% - 7.4% 3,179		*1.4% 0.0% - 3.0% 1,225
All Other Tobacco Products	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		0.4% 0.0% - 0.8% 268
INITIATION AND CESSATION OF CIGARETTE SMOKING IN PAST YEAR								
INITIATION—Percent of All Maryland Adults Who Report First Smoking a Cigarette During the Past Year (Est. # First Tried Cigarette Smoking Past Year / # All Adults) CESSATION—Percent of All Maryland Adults Who Report Quitting Cigarette Smoking During the Past Year (Est. # Quit Smoking Cigarettes Past Year / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Initiation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Cessation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		1.8% 0.5% - 3.0% 1,593

PREGNANT WOMEN — Cigarette Smoking At Any Time During Pregnancy

Maryland Residents Regardless of Where Delivery Occurred— **Birth Certificate Data, 2012**—Percent = Number Reported Smoking / Total Number

Percentage Smoking (%) Number Smoking (#)	2000	2002	2004	2006	2008	2010	2012
Pregnant Women	13.0% 158	12.3% 165	13.5% 195	9.9% 148	8.8% 127	12.8% 185	12.2% 169

SELECTED POPULATIONS — Current Cigarette Smoking

(Estimated # In Selected Population Currently Smoking Cigarettes / # All Adults In the Selected Population)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Non-Hispanic Black	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Non-Hispanic White	18.5% 11.1% - 25.9% 7,628	30.0% 20.0% - 39.9% 14,687	30.9% 20.5% - 41.4% 18,154	21.7% 14.3% - 29.0% 10,895	22.8% 15.3% - 30.4% 14,342	18.2% 12.1% - 24.4% 12,056	23.5% 13.5% - 33.5% 16,660
Hispanic	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Female	23.6% 13.6% - 33.5% 5,957	24.6% 13.0% - 36.3% 7,107	22.3% 12.3% - 32.4% 7,363	26.0% 16.5% - 35.4% 9,359	19.9% 13.3% - 26.5% 8,240	18.3% 11.1% - 25.5% 7,385	19.1% 9.0% - 29.1% 8,671
Male	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	37.2% 22.1% - 52.3% 11,983	35.9% 20.9% - 51.0% 13,336	20.7% 9.0% - 32.4% 6,980	24.8% 13.1% - 36.6% 9,516	13.7% 6.6% - 20.7% 6,144	*21.6% 7.7% - 35.6% 9,129
Highest Education: ≤ High School/GED	24.9% 12.8% - 36.9% 5,951	36.8% 20.8% - 52.8% 9,753	51.4% 35.4% - 67.5% 14,729	27.1% 15.1% - 39.1% 6,956	26.0% 16.3% - 35.7% 7,466	21.3% 12.7% - 29.9% 6,492	*22.5% 8.3% - 36.6% 9,390
Highest Education: ≥ 4 Years of College	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*12.0% 2.8% - 21.3% 2,614	*13.3% 1.8% - 24.8% 4,193	*4.8% 0.9% - 8.6% 1,303	*6.3% 2.2% - 10.3% 1,239
Household Income: < \$50,000 Annually	28.8% 14.9% - 42.6% 6,563	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	38.2% 22.3% - 54.1% 11,501	24.8% 12.7% - 35.9% 4,885	25.0% 13.1% - 36.9% 5,613	17.9% 7.7% - 28.1% 3,997	*20.4% 7.3% - 33.5% 4,892
Household Income: ≥ \$50,000 Annually	*12.2% 3.8% - 20.5% 2,984	20.6% 9.1% - 32.1% 6,312	23.8% 12.0% - 35.6% 8,634	22.3% 10.8% - 31.4% 8,236	22.8% 13.4% - 32.2% 10,619	13.4% 7.7% - 19.1% 6,776	20.8% 8.9% - 32.6% 10,410

% OF HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS

(Estimated # of Households with Resident Minor Children and a Resident Adult Smoker / # Households with Resident Minor Children)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Proportion with Adult Smokers	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	41.2% 24.3% - 58.1% 13,392



Talbot County

Tobacco Use by Adults: 2000–2012

Maryland Department of Health and Mental Hygiene—Center for Tobacco Prevention and Control
(See Appendix—Statewide and Jurisdiction Data for information on how to use and interpret these tables)

Historical jurisdiction estimates in this report may vary from those previously published. A methodology change in 2011 was designed to improve estimates of risk behaviors. Previous BRFSS estimates cannot be compared to the 2011 (and thereafter) data because of the significant differences in weighting methodology.

CURRENT USE OF ANY TOBACCO PRODUCT								
Adult (Age 18 and older) Estimated Current Use of One or More Tobacco Products (Estimated # Adults Currently Using Any Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
All Adults	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		21.5% 7.3% - 35.2% 6,776
CURRENT USE OF SPECIFIC TOBACCO PRODUCTS								
Adult (Age 18 and older) Estimated Current Use of Specific Tobacco Products (Estimated # Currently Using Specified Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Cigarettes	20.1% 9.5% - 30.8% 4,625	20.7% 9.2% - 32.2% 5,630	*15.0% 3.8% - 26.1% 4,167	16.2% 10.6% - 21.8% 4,564	12.6% 7.5% - 17.7% 3,208	13.6% 7.3% - 19.8% 4,382		*15.0% 4.3% - 25.8% 4,739
Cigars	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES
Smokeless Tobacco	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES		BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES
All Other Tobacco Products	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*1.0% 0.0% - 2.3% 322
INITIATION AND CESSATION OF CIGARETTE SMOKING IN PAST YEAR								
INITIATION—Percent of All Maryland Adults Who Report First Smoking a Cigarette During the Past Year (Est. # First Tried Cigarette Smoking Past Year / # All Adults) CESSATION—Percent of All Maryland Adults Who Report Quitting Cigarette Smoking During the Past Year (Est. # Quit Smoking Cigarettes Past Year / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Initiation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Cessation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES

PREGNANT WOMEN — Cigarette Smoking At Any Time During Pregnancy

Maryland Residents Regardless of Where Delivery Occurred— Birth Certificate Data, 2012—Percent = Number Reported Smoking / Total Number

Percentage Smoking (%) Number Smoking (#)	2000	2002	2004	2006	2008	2010	2012
Pregnant Women	14.4% 53	10.2% 34	8.0% 31	10.1% 36	10.6% 41	10.1% 36	11.8% 38

SELECTED POPULATIONS — Current Cigarette Smoking

(Estimated # In Selected Population Currently Smoking Cigarettes / # All Adults In the Selected Population)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Non-Hispanic Black	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Non-Hispanic White	21.5% 10.3% - 32.8% 4,625	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	15.1% 9.7% - 20.5% 3,394	13.2% 7.4% - 19.0% 2,682	13.0% 7.0% - 19.0% 3,288	*18.1% 4.8% - 31.4% 4,650
Hispanic	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Female	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	17.5% 10.8% - 24.1% 2,592	8.6% 3.9% - 13.3% 1,202	12.7% 6.4% - 19.0% 2,367	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES
Male	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*14.8% 5.6% - 24.0% 1,972	17.5% 8.0% - 27.0% 2,006	*14.7% 2.9% - 26.5% 2,015	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES
Highest Education: <= High School/GED	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	19.4% 8.7% - 30.1% 2,051	15.6% 6.8% - 24.3% 1,451	*9.5% 2.4% - 16.6% 1,200	*18.4% 1.5% - 35.3% 2,366
Highest Education: >= 4 Years of College	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*11.6% 3.9% - 19.4% 1,413	*7.3% 0.5% - 14.1% 815	*9.6% 1.7% - 17.5% 1,279	*5.2% 0.8% - 9.5% 437
Household Income: < \$50,000 Annually	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	22.2% 11.7% - 32.7% 2,349	23.2% 12.5% - 33.9% 1,940	*11.5% 2.0% - 21.0% 1,303	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES
Household Income: >= \$50,000 Annually	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*13.1% 5.2% - 21.0% 1,770	*9.5% 2.6% - 16.4% 1,184	*15.3% 5.4% - 25.2% 2,389	*19.3% 2.1% - 36.5% 2,497

% OF HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS

(Estimated # of Households with Resident Minor Children and a Resident Adult Smoker / # Households with Resident Minor Children)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Proportion with Adult Smokers	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES



Washington County

Tobacco Use by Adults: 2000–2012

Maryland Department of Health and Mental Hygiene—Center for Tobacco Prevention and Control
(See Appendix—Statewide and Jurisdiction Data for information on how to use and interpret these tables)

Historical jurisdiction estimates in this report may vary from those previously published. A methodology change in 2011 was designed to improve estimates of risk behaviors. Previous BRFSS estimates cannot be compared to the 2011 (and thereafter) data because of the significant differences in weighting methodology.

CURRENT USE OF ANY TOBACCO PRODUCT

Adult (Age 18 and older) Estimated Current Use of One or More Tobacco Products (Estimated # Adults Currently Using Any Tobacco Product / # All Adults)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
All Adults	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		20.8% 14.0% - 27.6% 21,800

CURRENT USE OF SPECIFIC TOBACCO PRODUCTS

Adult (Age 18 and older) Estimated Current Use of Specific Tobacco Products (Estimated # Currently Using Specified Tobacco Product / # All Adults)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Cigarettes	23.6% 17.2% - 30.0% 23,026	24.0% 17.9% - 30.1% 26,037	24.4% 16.9% - 31.8% 24,701	23.3% 18.0% - 28.5% 25,221	25.7% 19.5% - 31.9% 30,793	18.3% 13.2% - 23.5% 19,969		18.0% 11.3% - 24.7% 18,809
Cigars	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*2.6% 0.4% - 4.9% 2,708
Smokeless Tobacco	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	*5.2% 1.4% - 9.0% 5,675		*1.9% 0.4% - 3.4% 2,095
All Other Tobacco Products	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*2.9% 0.6% - 5.3% 2,991

INITIATION AND CESSATION OF CIGARETTE SMOKING IN PAST YEAR

INITIATION—Percent of All Maryland Adults Who Report First Smoking a Cigarette During the Past Year (Est. # First Tried Cigarette Smoking Past Year / # All Adults)

CESSATION—Percent of All Maryland Adults Who Report Quitting Cigarette Smoking During the Past Year (Est. # Quit Smoking Cigarettes Past Year / # All Adults)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Initiation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Cessation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*1.6% 0.4% - 2.8% 1,719

PREGNANT WOMEN — Cigarette Smoking At Any Time During Pregnancy

Maryland Residents Regardless of Where Delivery Occurred— **Birth Certificate Data, 2012**—Percent = Number Reported Smoking / Total Number

Percentage Smoking (%) Number Smoking (#)	2000	2002	2004	2006	2008	2010	2012
Pregnant Women	19.1% 305	18.1% 306	15.6% 266	16.3% 310	16.3% 298	19.9% 351	21.3% 376

SELECTED POPULATIONS — Current Cigarette Smoking


(Estimated # In Selected Population Currently Smoking Cigarettes / # All Adults In the Selected Population)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Non-Hispanic Black	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Non-Hispanic White	24.4% 17.6% - 31.1% 21,311	24.6% 18.2% - 30.9% 25,328	25.5% 17.5% - 33.4% 23,616	22.7% 17.4% - 28.0% 21,907	22.9% 17.0% - 28.8% 23,933	20.3% 14.7% - 26.0% 19,969	15.9% 10.2% - 21.6% 14,424
Hispanic	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Female	22.4% 13.8% - 31.1% 11,350	22.8% 15.1% - 30.5% 13,037	22.2% 13.9% - 30.6% 11,884	25.7% 19.2% - 32.2% 13,967	18.4% 12.6% - 24.3% 11,034	19.6% 13.0% - 26.2% 11,703	13.9% 8.9% - 18.9% 8,010
Male	24.9% 15.4% - 34.4% 11,676	25.3% 15.5% - 35.0% 13,000	26.7% 14.2% - 39.2% 12,817	20.9% 12.6% - 29.1% 11,254	33.0% 22.4% - 43.6% 19,758	16.8% 8.7% - 24.9% 8,266	22.9% 10.0% - 35.9% 10,799
Highest Education: ≤ High School/GED	28.5% 19.4% - 37.6% 14,704	29.1% 20.2% - 36.0% 16,747	28.5% 17.2% - 39.9% 14,368	37.2% 27.9% - 46.4% 16,873	30.4% 21.2% - 39.6% 18,362	24.7% 16.7% - 32.7% 14,141	24.6% 14.1% - 35.1% 14,595
Highest Education: ≥ 4 Years of College	*19.0% 4.3% - 33.6% 4,459	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*8.5% 3.0% - 13.9% 2,569	*10.9% 3.6% - 18.2% 2,820	BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES	*3.7% 0.4% - 7.0% 856
Household Income: < \$50,000 Annually	22.7% 14.6% - 30.9% 12,200	33.8% 24.6% - 43.0% 19,865	22.0% 13.1% - 31.0% 9,683	26.0% 18.3% - 33.8% 12,287	35.5% 25.2% - 45.8% 19,889	26.6% 17.8% - 35.5% 12,238	28.6% 15.2% - 42.0% 12,740
Household Income: ≥ \$50,000 Annually	*21.3% 7.8% - 34.8% 5,595	16.0% 6.8% - 25.3% 5,519	22.2% 10.7% - 33.7% 8,708	20.3% 12.5% - 28.1% 9,267	18.1% 10.7% - 25.6% 8,614	12.3% 5.6% - 19.1% 5,728	*6.7% 2.6% - 10.8% 3,190

% OF HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS

(Estimated # of Households with Resident Minor Children and a Resident Adult Smoker / # Households with Resident Minor Children)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Proportion with Adult Smokers	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	30.0% 16.7% - 43.3% 12,818



Wicomico County

Tobacco Use by Adults: 2000–2012

Maryland Department of Health and Mental Hygiene—Center for Tobacco Prevention and Control
(See Appendix—Statewide and Jurisdiction Data for information on how to use and interpret these tables)

Historical jurisdiction estimates in this report may vary from those previously published. A methodology change in 2011 was designed to improve estimates of risk behaviors. Previous BRFSS estimates cannot be compared to the 2011 (and thereafter) data because of the significant differences in weighting methodology.

CURRENT USE OF ANY TOBACCO PRODUCT								
Adult (Age 18 and older) Estimated Current Use of One or More Tobacco Products (Estimated # Adults Currently Using Any Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
All Adults	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		26.4% 15.5% - 37.3% 17,134
CURRENT USE OF SPECIFIC TOBACCO PRODUCTS								
Adult (Age 18 and older) Estimated Current Use of Specific Tobacco Products (Estimated # Currently Using Specified Tobacco Product / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Cigarettes	25.0% 15.1% - 34.8% 17,777	25.5% 17.1% - 33.9% 14,420	23.7% 14.7% - 32.8% 15,736	22.2% 15.6% - 28.7% 13,972	17.3% 11.1% - 23.5% 12,488	22.7% 15.3% - 30.3% 16,344		20.3% 11.7% - 24.7% 13,166
Cigars	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES
Smokeless Tobacco	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	*1.4% 0.1% - 2.8% 1,043		*1.8% 0.2% - 3.3% 1,167
All Other Tobacco Products	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES
INITIATION AND CESSATION OF CIGARETTE SMOKING IN PAST YEAR								
INITIATION—Percent of All Maryland Adults Who Report First Smoking a Cigarette During the Past Year (Est. # First Tried Cigarette Smoking Past Year / # All Adults) CESSATION—Percent of All Maryland Adults Who Report Quitting Cigarette Smoking During the Past Year (Est. # Quit Smoking Cigarettes Past Year / # All Adults)								
Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Initiation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Cessation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*3.8% 1.0% - 6.6% 2,521

PREGNANT WOMEN — Cigarette Smoking At Any Time During Pregnancy

Maryland Residents Regardless of Where Delivery Occurred — Birth Certificate Data, 2012 — Percent = Number Reported Smoking / Total Number

Percentage Smoking (%) Number Smoking (#)	2000	2002	2004	2006	2008	2010	2012
Pregnant Women	14.4% 169	15.1% 175	14.8% 172	13.3% 177	15.5% 212	16.4% 205	13.7% 166

SELECTED POPULATIONS — Current Cigarette Smoking


(Estimated # In Selected Population Currently Smoking Cigarettes / # All Adults In the Selected Population)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Non-Hispanic Black	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*31.0% 12.7% - 49.3% 5,187	*26.9% 3.2% - 50.7% 5,155
Non-Hispanic White	23.8% 14.1% - 33.5% 11,618	24.9% 15.7% - 34.1% 11,391	24.1% 13.6% - 34.5% 12,509	22.9% 15.8% - 29.9% 11,254	16.2% 9.9% - 22.4% 7,841	22.8% 14.3% - 31.3% 10,811	*19.0% 13.1% - 24.9% 7,558
Hispanic	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE
Female	17.5% 8.7% - 26.2% 6,418	18.1% 9.4% - 26.7% 6,028	*14.6% 6.0% - 23.3% 5,528	17.4% 11.1% - 23.7% 6,639	15.6% 9.6% - 21.7% 5,217	17.4% 9.2% - 25.5% 6,752	19.5% 12.0% - 26.9% 5,706
Male	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	29.5% 16.3% - 42.8% 7,333	18.7% 8.4% - 29.1% 7,271	29.1% 16.1% - 42.1% 9,592	*21.1% 6.1% - 36.0% 7,460
Highest Education: ≤ High School/GED	*27.7% 11.4% - 44.0% 9,841	31.0% 17.3% - 44.7% 8,268	34.6% 17.3% - 51.8% 9,145	30.3% 18.5% - 42.1% 7,276	27.7% 15.2% - 40.2% 8,181	31.1% 18.4% - 43.9% 8,532	24.0% 13.9% - 34.2% 7,521
Highest Education: ≥ 4 Years of College	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*17.6% 4.3% - 30.9% 4,097	*10.5% 3.2% - 17.9% 2,460	*5.6% 1.0% - 10.2% 1,390	*10.0% 1.8% - 18.2% 2,122	*6.7% 2.3% - 11.2% 1,193
Household Income: < \$50,000 Annually	26.8% 15.5% - 38.1% 10,026	33.8% 21.4% - 46.3% 10,456	28.2% 15.8% - 40.6% 10,682	29.0% 18.8% - 39.2% 8,685	22.3% 11.6% - 33.0% 7,518	21.4% 11.96% - 30.9% 7,412	30.9% 14.3% - 47.5% 8,805
Household Income: ≥ \$50,000 Annually	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA IS SUPPRESSED DUE TO SMALL SAMPLE	*16.8% 6.9% - 26.7% 4,154	13.5% 6.0% - 21.1% 4,044	22.9% 10.3% - 35.5% 6,260	*11.7% 4.4% - 18.9% 3,457

% OF HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS

(Estimated # of Households with Resident Minor Children and a Resident Adult Smoker / # Households with Resident Minor Children)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Proportion with Adult Smokers	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	27.6% 13.9% - 41.4% 5,681



Worcester County

Tobacco Use by Adults: 2000–2012

Maryland Department of Health and Mental Hygiene—Center for Tobacco Prevention and Control
(See Appendix—Statewide and Jurisdiction Data for information on how to use and interpret these tables)

Historical jurisdiction estimates in this report may vary from those previously published. A methodology change in 2011 was designed to improve estimates of risk behaviors. Previous BRFSS estimates cannot be compared to the 2011 (and thereafter) data because of the significant differences in weighting methodology.

CURRENT USE OF ANY TOBACCO PRODUCT

Adult (Age 18 and older) Estimated Current Use of One or More Tobacco Products (Estimated # Adults Currently Using Any Tobacco Product / # All Adults)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
All Adults	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		8.6% 3.1% - 14.0% 3,169

CURRENT USE OF SPECIFIC TOBACCO PRODUCTS

Adult (Age 18 and older) Estimated Current Use of Specific Tobacco Products (Estimated # Currently Using Specified Tobacco Product / # All Adults)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Cigarettes	23.7% 13.5% - 33.8% 10,200	30.5% 19.5% - 41.5% 11,771	20.1% 10.5% - 29.7% 8,750	21.6% 13.0% - 30.2% 8,252	17.2% 7.8% - 26.7% 6,797	11.5% 4.5% - 18.5% 4,488		*7.7% 2.4% - 13.1% 2,854
Cigars	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		*0.8% 0.0% - 1.8% 266
Smokeless Tobacco	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	*2.8% 0.0% - 5.6% 1,085		*1.6% 0.0% - 3.6% 580
All Other Tobacco Products	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA ARE SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES

INITIATION AND CESSATION OF CIGARETTE SMOKING IN PAST YEAR

INITIATION—Percent of All Maryland Adults Who Report First Smoking a Cigarette During the Past Year (Est. # First Tried Cigarette Smoking Past Year / # All Adults)

CESSATION—Percent of All Maryland Adults Who Report Quitting Cigarette Smoking During the Past Year (Est. # Quit Smoking Cigarettes Past Year / # All Adults)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010		2012
Initiation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE
Cessation	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE		6.8% 0.0% - 14.6% 2,522

PREGNANT WOMEN — Cigarette Smoking At Any Time During Pregnancy

Maryland Residents Regardless of Where Delivery Occurred— **Birth Certificate Data, 2012**—Percent = Number Reported Smoking / Total Number

Percentage Smoking (%) Number Smoking (#)	2000	2002	2004	2006	2008	2010	2012
Pregnant Women	16.9% 84	15.4% 70	14.9% 67	16.1% 75	14.2% 68	14.1% 59	13.4% 61

SELECTED POPULATIONS — Current Cigarette Smoking

(Estimated # In Selected Population Currently Smoking Cigarettes / # All Adults In the Selected Population)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Non-Hispanic Black	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE
Non-Hispanic White	26.0% 14.0% - 38.0% 8,426	28.0% 15.4% - 40.6% 8,962	18.1% 8.2% - 28.0% 7,002	19.4% 11.7% - 27.2% 6,221	19.3% 11.7% - 26.9% 6,473	11.8% 5.1% - 18.5% 3,607	4.7% 0.9% - 8.5% 1,239
Hispanic	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE
Female	18.7% 6.0% - 31.4% 3,552	30.0% 15.0% - 45.0% 7,032	20.3% 7.3% - 33.3% 4,858	20.6% 11.0% - 30.2% 4,004	15.9% 8.1% - 23.8% 3,816	9.0% 2.1% - 15.9% 1,690	6.7% 1.5% - 11.9% 1,148
Male	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	22.6% 10.2% - 35.0% 4,248	19.2% 6.6% - 31.8% 2,980	13.8% 3.8% - 23.8% 2,798	8.6% 1.2% - 16.0% 1,707
Highest Education: ≤ High School/GED	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	30.4% 14.7% - 46.0% 4,181	*24.2% 2.3% - 46.1% 3,210	*13.7% 4.7% - 22.8% 1,835	*7.1% 2.3% - 11.8% 1,341
Highest Education: ≥ 4 Years of College	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	*12.2% 2.8% - 21.6% 1,371	BRFSS DATA ARE SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES	BRFSS DATA ARE SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES	*2.0% 0.0% - 4.0% 222
Household Income: < \$50,000 Annually	29.0% 15.2% - 42.8% 7,668	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	20.5% 10.1% - 30.8% 2,997	*28.4% 9.9% - 46.9% 4,771	*20.5% 6.7% - 34.2% 3,594	*13.0% 2.8% - 23.3% 2,577
Household Income: ≥ \$50,000 Annually	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE	*18.4% 6.0% - 30.8% 3,296	*11.1% 3.0% - 19.1% 1,775	BRFSS DATA ARE SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES	BRFSS DATA ARE SUPPRESSED DUE TO IMPRECISE AND UNRELIABLE ESTIMATES

% OF HOUSEHOLDS WITH MINOR CHILDREN THAT HAVE RESIDENT ADULT SMOKERS

(Estimated # of Households with Resident Minor Children and a Resident Adult Smoker / # Households with Resident Minor Children)

Estimated Prevalence (%) Confidence Interval (%-%) Estimated Prevalence (#)	2000	2002	2004	2006	2008	2010	2012
Proportion with Adult Smokers	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA NOT AVAILABLE	BRFSS DATA ARE SUPPRESSED DUE TO SMALL SAMPLE

APPENDIX C—WHERE TO FIND MORE INFORMATION

Center for Tobacco Prevention and Control

See: http://phpa.dhmh.maryland.gov/ohpetup/SitePages/tob_home.aspx

Maryland Department of Health and Mental Hygiene

Center for Tobacco Prevention and Control

Dawn Berkowitz, Director, 410-767-5529

MDQuit.org

See: <http://mdquit.org>

Maryland's Resource Center for Quitting Use and Initiation of Tobacco

At the University of Maryland Baltimore County

Dr. Carlo DiClemente, Director, 410-455-3628

Legal Resource Center for Public Health Policy

See: <http://www.law.umaryland.edu/programs/tobacco/index.html>

Legal support to communities, community groups, employers, local governments, and state legislators and agencies interested in reducing the negative health consequences of tobacco use

Kathleen Hoke, Director, 410-706-0842

Maryland's Local Health Departments

See <http://mdquit.org/maryland-county-health-departments>

The Health Consequences of Smoking—50 Years of Progress: A Report of the Surgeon General, 2014

See: <http://www.surgeongeneral.gov/library/reports/50-years-of-progress/index.html>